village of the Bagar south-east of Sirsa, but it was found impossible to supply it with water for irrigation purposes. A proposal to bring a kharif-irrigation branch from the main line of the Western Jamna Canal through the Hissar district into the south-east end of the Sirsa district was referred to the Secretary of State in 1871, but was set aside on account of want of funds. It is now again being discussed and may some day be realised.

Area dependent on the local rainfall.

narily irrigated either from wells or by the floods of the Satlaj and Ghaggar, which depend on the melting of the snows on the distant Himálaya or the rainfall on the lower ranges; and 94 per cent of the cultivation depends on the scanty and uncertain local rainfall. The area returned at the present Settlement measurements as ordinarily cultivated with the aid of the rainfall alone (bárání is 9,77,502 acres, but the rainfall is rarely so general and so favourable as to allow of nearly the whole of this area being actually cultivated. The areas actually cultivated with rain alone during the past seven harvests have been as follows:—

Agricultural year.	Area cultivated by rain alone in.					
	Kharif.	Rabí.	Whole year.			
1879-80 1880-81 1881-82 1882-83	Not observed. 713,967 756,552 702,523	256,267 123,481 225,846 300,118	837,448 981,898 1,002,641			

But the last two years have been exceptionally favourable years, and even in 1880 the seed-time of the kharif was unusually promising, and the statistics of these three years do not adequately show the fluctuations to which the rain-cultivation is liable. In 1877-78 the area returned as cultivated was less than 600,000 acres, and there is reason to believe that even this was an exaggeration, and that the area actually sown in that year was very much less than six lakhs. So uncertain is the rainfall and so much is the cultivation dependent on it that it would be nothing extraordinary in a bad year for half the cultivated land in the district to lie unsown. Moreover, unless the subsequent rains are favourable, a large area sown produces no crop at all, and a still larger area produces no grain, only straw. For instance, in the kharif of 1880 although 713,967 acres were actually sown, only 351,513 acres produced a grain crop, so that more than half the area sown produced nothing but straw, and the greater part of the remainder produced very little grain; and again in the kharif of 1882, half the

area sown in tahsil Dabwali was returned as having produced no

grain.

The statement of annual rainfall shows how variable is the Uncertainty of the rainfall. quantity of rain which annually falls at each of the recording stations, and it is to be remembered that the showers are very partial, and that often one village gets rain when its neighbour gets none. But it is not so much upon the quantity of rain that falls as on its seasonableness that the success of failure of the crops depends. The rains usually set in about the end of June, and for a favourable kharif seedtime a heavy downpour is wanted then or in the beginning of July. If the rain be then good and general a large area is sown with kharif crops, and to bring them to maturity showers of rain are wanted at intervals up to September. If during this period no rain falls for a month or so, the crops dry up and produce no grain, and sometimes not even straw. If the early rains are scanty, only a small area is sown, and unless the seed is in the ground before the middle of August, it comes to nothing. It may then be said that the area of land sown for the kharif varies with the amount of rain that falls in June and July, and that the outturn per acre varies according to the seasonableness of the showers that fall in the following months. For the rabi erops there should be good rain in August or September to moisten the land thoroughly for the sowing, and one or two showers are necessary in December, January or February, to save the crop from drying up. The people estimate the amount of rain that falls by the number of finger-breadths (ungal) it has sunk into the ground—a down-pour that penetrates into the ground 100 ungal being considered perfection. If there be already moisture in the ground, it is sufficient that the rain from above should sink to the moisture below (vattar se vattar milgal, or al se al milgai). The peasants say that if it rains for 24 hours incessantly, it is a sure sign that there has been rain for a distance of 100 kos (150 miles). Even if there be good rain in June and July, the rainfall of August and September is very important, for on it depend both the ripening of the kharif crop and the sowing of the rabí. Sometimes, even in this district, there is too much rain. Heavy rain sometimes washes the seed out of the sandy soil, or drowns it on harder ground; and an inopportune shower when the bajra is blossoming prevents the ear from being fertilised. The rice-crop also is injured by heavy rain. More than once the fever that follows heavy rains has so much weakened the scanty population of the district, as appreciably to lessen the area cultivated. Inopportune rains in October may injure the ripening kharif crops, or interfere with the rabi sowings, and in some soils where there is kankar not far below the surface, heavy showers in the cold weather kill the sprouting rabi.

Account of some good and phenomena of some of the good and bad years.

Account of some good and years through which the district has passed.

In the good year 1872-73, the monthly rainfall

at the three tahsils was as follows :-

	Month.		Sirsá.	Dabwáli.	Fázilká.
			Maria San	Party Law Park	
April *			· · · 2	3	:1
May			9 *	3	1.1
June	***]	2:2	-6	1
July	14160 F-17.7		7.2	2.	6.
August	***		4.6	5.5	•3
September			1.6	1.2	4.3
October	***			1	•
November	***	!			Alterial No. of the State of th
December	•••		1.6	6	.5
	***		1.6		
January	•••		******		******
ebruary	***		******		
March	•••		******	-8	••••
	TOTAL	[18:3	10.8	12.5

The heavy rainfall in June, July and August gave a good seed-time for the kharif, and the rains in September brought it to maturity. The harvest was an exceptionally good one, though in places it suffered from the depredations of locusts from Bikaner. The fair rains of August and September moistened the ground thoroughly for the rabi sowings, the rain in December fell just when it was wanted, and the rabi was a good crop. It may be noted that except at Sirsá the rainfall was actually below the average, but it was so seasonable (especially the comparatively slight fall in December) that the harvests were good. Although the district had suffered from bad seasons for four years, since the famine of 1868-69, these good crops enabled the people to pay in this year (including balances of former years) more than one and a half times the annual revenue demand.

In the good year 1875-76 the rainfall was as follows:-

	Month.		Sirsá.	Dabwáli.	Fázilká.
April	er eren var var en	944	*****	1	******
May			-4	3	-2
June		***	:3	1.	•4
July	The second secon		4-9	4.6	3.4
August			3.	1.3	16.6
September		***	9.5	8-1	7.1
October		1	-6	1	*****
November	The same has been		******	*****	*****
December	***	***	******	*****	******
January	•••		******		*****
February		***	100 may 40 115	-2	.8
March			7	1.1	3
	* TOTAL		19-4	16.8	28-1

The total rainfall for the year was above the average everywhere. The good rain in July and August enabled the peasants to sow a large extent of land for the kharif, and the heavy fall in September suited both seasons, and encouraged them to sow an unusually large area for the rabi also. Although little rain fell in October and none in November, December or January, the subsoil which had been well soaked in September remained moist and the rabi outturn was good. The balances which still hung over the district were nearly all cleared off.

In the good year 1878-79 the rainfall was as follows :-

	Month.		Sirsá.	Dabwálí.	Fázilká.	
April			3.4	1.3	0.9	
May			3.4	2.2	1.2	
June			1.7	0.5	0.6	
July	•••		3.5	4.3	69	
August	•••		9.7	7.3	7.7	
September	•••		0.1	0.8	•••••	
October						
November	•••					
December		:	1.0			
January						
February		•••		0.3	•••••	
March		•••	6.6	0.7	- 0.6	
To	TAL		23.4	17:4	17-8	

The total rainfall was everywhere much above the average. The rainfall in April and May was very exceptional, as usually in those months hardly any rain falls. In consequence of this early heavy rain there was a great deal of fever in the district, but these early rains and the good fall in July and August gave an excellent kharif; and the ground was so thoroughly soaked in August that the rabi sowings were very extensive, and although after September no rain fell at Dabwali until February and at Fazilka till-March, the rabi harvest was a bumper everywhere and almost all the balances which had accrued in the previous bad year 1877-78 were paid up.

'In the famine year 1860-61 the rainfall was as follows :-

Month.	Sirsá.	Dabwálí.	Fázilká.
May	*1.1	0.6	0.1
June	1.4	0.8	0.4
July *	2.7	2.5	2.2
August	1.6	0.6	8.3
September		•	•••••
October			
November			
December			
January	1.1	0.9	2.6
February			
March		0.2	
April	0.6		0.1
TOTAL	8.5	5.6	13.7

The total rainfall, except at Fázilká, was much below the average. The comparatively scanty rains of June and July did not sufficiently moisten the ground, and there being little rain in August at Sirsá and Dabwálí and none at all throughout the district in the following four months, the kharíf crops dried up and the rabí could not be sown, for the rain in January was too late to help the rabí sowings. Here then, owing to the failure of rain in August and September, both kharíf and rabí crops were lost. The harvests of the previous three years had been poor, and the distress in the district was very severe until the good rains of 1861 came to relieve it. The revenue fell into arrears which were not paid up for several years.

In the famine year 1868-69 the rainfall was as follows:-

Month.		Sirsá.	Dabwálí.	Fázilká	
April			0.1	0.3	0.5
May			0.4	0.3	0.6
T.			4.0	0.8	0.4
July			1.3	2.8	5.6
August			0.3	0.3	0.1
September			0.5	2.3	
October	•••		*****	0.2	*****
THE RESIDENCE OF THE PROPERTY	•••		*****		0.2
		***	1.0	1.3	0.8
February	•••		0.4	0.4	. 0.1
March	•••	***	1.8	2.2	2.7
Тот	AL		9.8	10.7	11.0

The total rainfall was below the average everywhere. The rains in June and July were exceptionally light, and there was very little in August, so that what kharif crops were sown dried up, and the September rain was too light to restore them: so that the kharif was a failure. Owing to the absence of good rain in August, September and October the ground was not in a fit state for the rabi sowings, and there being as usual no rain in November and but little in December there was no rabi crop to benefit from the rain of January and February. Here again the want of rain in August and September ruined both the kharif and the rabi. The previous year's harvests had failed in part of the district and the distress was great. Balances were incurred which were not finally paid off for several years. The importance of the later autumn rains was shown next year (1869) when again the kharif crop seemed to be gone, but good rain in the beginning of September thoroughly revived it.

In the scarcity year 1877-78 the rainfall was as follows:-

, , , , , , , , , , , , , , , , , , ,	Ionth.		Sirsá.	Dabwálí.	Fázilká
April			0-8	3.4	0.9
May	***		0.6	1.1	1.5
June			2.2	2.0	0.9
July	Value of the second		2:2	2.4	3.3
August	• • •		0.1	0.2	0.9
September	•••		2.1	2.7	3.5
October			0.1	0.1	0.6
November			0.7	1.2	0.8
December	100		3.2	0.6	0.9
January			0.1	0.1	
February			0.5	0.4	0.6
March		•••		0.5	0.4
To	TAL		12.6	14.7	14.3

Although, except at Sirsá, the total rainfall was actually above the average, it was too much spread over the year. The failure of rain in August caused the kharif crop to dry up, and the rain in September was neither sufficient to restore it nor to enable the people to sow the rabi well. Both harvests were much below the average, but the previous two years had been good; consequently there was comparatively little distress felt and the balances were small.

Nature of the rainfall in District Officers, the seasons have varied from year to vear, chiefly as regards the crops dependent on the local rainfall:—

Yı	EAR.	CHARACTER OF	
Sambat.	A. D.	SEASON.	Remarks.
1904	1847-48	Fair.	
1905	1848-49	Famine.	
1906	1849-50	Fair.	
1907	1850-51	Good	Kharif failed in part of the district.
1908	1851-52	Poor	Kharif a failure. Rabi very scanty.
1909	1852-53	Good	II in and not all and
1910	1853-54	Poor.	
1911	1854-55	Poor.	
1912	1855-56	Fair.	
1913	1856-57	Excellent.	
1914	1857-58	Fair	Small area sown because of the mu
1915	1858-59	Poor.	
1916	1859-60	Poor	Kharif failed. Rabi below average.
1917	1860-61	Famine	Both harvests entirely failed.
1918	1861-62	Fair.	
1919	1862-63	Good	Both harvests good.
1920	1863-64	Fair.	
1921	1864-65	Poor	Much land unploughed.
1922	1865-66	Good	Towards Fázilká excellent.
1923	1866-67	Poor	Harvest less than half the average.
1924	1867-68	Poor	Fair along north-east; a failure t south-west.
1925	1868-69	Famine	Both harvests failed.
1926	1869-70	Excellent.	
1927	1870-71	Poor.	
1928	1871-72	Poor.	
1929	1872-73	Good	Injury done by locusts.
1930	1873-74	Poor	Rain failed towards end of season.
1931	1874-75	Poor	Winter rains failed.
1932	1875-76	Excellent	Both harvests good.
1933	1876-77	Fair	Kharif suffered from hot wind.
1934	1877-78	Searcity	Kharif dried up. Rabi poor.
1935	1878-79	Excellent	The state of the s
1936	1879-80	Fair	T1 12 D-17 1
1937	1880-81	Poor	Kharif dried up. Rabi little sown.
1938	1881-82	Excellent	Kharif and rabi both good.
1939	1882-83	Fair	TOTAL

It appears that in the last 36 years the seasons have been in-

5	years		***	Excellent.
5	years			Good.
9	years	17.000		Fair.
13	years .	5		Poor.
4	years			Scarcity and Famine.

In this statement I have given a general estimate of the nature of each season for the district as a whole. The enquiries were made in more than a hundred villages throughout the different assessment circles, and in each year there were many exceptions to the general rule. In the very best years some villages got little produce and in the very worst year some had good rain. There are always patches of country which fare better or worse than the tract generally.

Crops grown on unirri- 166. The crops cultivated by rain alone

gated lands. in 1881-82 were as follows:-

KHARIF	•	Rabi.			
Crop.	Area in Acres.	Crop.		Area in Acres	
Jawar (alone and with pulses) Bajra (alone and with pulses) Moth Mung Gwar Miscellaneous	135,513 528,715 36,847 3,292 23,549 28,574	Wheat Wheat and Gram Gram Barley and Gram Barley Sarson and Tara Miscellaneous	::	11,010 1,329 22,349 109,177 77,517 2,048 1,916	
TOTAL	756,552 •	· Total	***	235,346	

The staple crops are bájra mixed with pulses in the kharíf, and barley and gram in the rabí.

167. The cultivation of the kharif crops on the ordinary rainland is a very simple affair. When a sufficient System of cultivation. and opportune shower of rain falls, each man goes out to the fields with his plough and his camels, bullocks or donkeys, and ploughs and sows as much land as he can before the moisture leaves the ground. It is not necessary to give the land a preliminary ploughing; indeed sometimes prairie-land previously uncultivated is sown at the first ploughing. A drill of bamboo is attached to the plough and sowing begun at once, the ploughman feeding the drill with the hand with which he holds the plough, from a bag of seed slung at his back. The plough, drill and harness cost altogether little more than one rupee. With an eighty-rupee camel a man can plough and sow two or even three acres in a day; with a pair of twenty-five rupee bullocks an acre or more; with one bullock three-fourths of an acre; with a pair of donkeys still less. Camels are used chiefly by the Bagris along the south-west of the district, and in hot weather plough only by night; bullocks by the Sikh Jats along the north-east, and by the Musalmans along the centre of the district and in the Ghaggar and Satlaj valleys; donkeys by their Kumhar (potter) owners. When bullocks are scarce

after a drought, the women sometimes draw the plough. About three ser of bajra is sown per acre generally mixed with moth, mung, pumpkin (kakri) and water-melon (matira); jawar takes from 8 to 18 sers per acre and generally has moth or mung mixed with it. Gwar is sown both with these crops and separately, about seven ser of seed to the acre. Til is sown broadcast one to three sers per acre. The hard clay soil of the Sotar valley requires to be ploughed once before sowing; a pair of bullocks in the first ploughing get over half an acre a day and then sow about an acre a day. The seed can be sown only while the moisture remains in the ground, and there is at such times a great demand for ploughs, a man with his camel and plough sometimes getting Re. 1-4 or Re. 1-8 for a day's work. Such favourable times are so few that in one season one plough can sow only about 20 acres for the kharif. Often in light sandy soils the seed gets washed or blown out of the ground or covered with light sand, and the sowing has to be done over again; or the first sowing dries up, and a shower of rain encourages the peasant to sow again. Otherwise he has simply to wait until the sun and rain give him a crop or none. When the first ears begin to ripen, sometimes before the end of September, the women pluck enough to give fresh grain for the day's meal, and when harvesting begins in earnest the whole family often camp out in the fields for the time, living on the fresh grain and the juicy water-melons. If there is much grain to reap, the peasant cuts off the ears first and threshes them with the help of his oxen, and afterwards cuts down the straw at his leisure. Sometimes February comes in before all the bajfa is threshed, and the straw cut and stacked.

More pains are taken with the cultivation of the rabi crop. Sometimes the land is turned over in the early part of the rainy season and exposed to sun and rain. It generally gets at least two ploughings, sometimes as many as four, before it is sown; and where necessary the soil is pulverised and levelled by the schaga or clod-crusher, a plank or log on which the ploughman stands while his oxen draw it over the field. Then the ground is ploughed again, and the seed drilled in at the same time through a drill attached to the plough, the furrows being made much closer than for the kharif. About 25 sers of wheat are sown to the acre, and about 22 of barley, gram and oilseeds, the barley and gram being sown either separately or mixed, and the oilseed (sarson or tara) being generally sown in lines among the other crops. The cultivator must make the most of the favourable seedtime in October and November, and usually a plough can sow only eight or ten acres for the rabi. If the kharif has failed, or the first rabí sowing has dried up, a good shower in December encourages the peasant to sow again in hopes of a crop; but such late sowings (pichheti or kanaujai) seldom produce anything but a little straw. The busiest times of the year are in October and November when the kharif is being cut and the rabi sown, and in March and April when the rabi is being reaped, as all the rabi crops ripen much about the same time, and if the ripe crop is allowed to stand long uncut it is apt to be shaken

by the wind or injured by hail or animals. The easiest times of the year are the hottest months (May and June) and the coldest months (December to February). The Sikhs with their fine bullocks devote most of their attention to the rabi (Hárí); the Bágris with their camels have hitherto cultivated little but the kharif (Sáwaní), the only crop known on the Bíkáner prairies; it is only of late years that, copying their neighbours the Sikhs, they have begun to cultivate the rabi, and they do not yet make a thoroughly good job of it. But the cultivation of the rabi is rapidly extending among them, and in the last few years some villages have begun to sow a rabi crop which had never sown it before.

In the case of land dependent on rain only, it is usual to sow one crop each year, thus allowing the land to lie fallow half the year; for instance, a field sown with bájra in the kharif will lie fallow in the following rabí and be again cultivated in the next kharíf. No such land bears two crops in one agricultural year, that is, land sown with a kharif crop is invariably left fallow in the following rabí, unless the kharíf crop has totally failed. Some of the Sikh peasants, when they wish to change the crop, allow the land to lie fallow for a whole year instead of for six months only, that is, land which has borne a kharif crop is allowed to lie fallow for the following rabí and kharif and then sown with a rabí crop, to be followed immediately by a kharif. But this course is not often kept up, and generally speaking land best suited for a kharif crop is always sown with kharif, and land best suited for a rabi crop is always sown for the rabi. The trouble and expense required in growing a kharif crop are much less than for the rabi; a preliminary ploughing is hardly necessary, the seed costs much less and the produce is less valuable and requires less care and watching; moreover, at the time when the kharif operations are in progress, there is generally plenty of water in the hollows, and grass everywhere available for the plough-bullocks. It is therefore common to sow a kharif crop first in land newly-broken up, and to sow only kharif crops in land situated far from the village. On the other hand, light loam does not produce a good kharif crop after it has been under cultivation for some years, but when well worked up produces a fair rabi erop with little rain, the seed is more valuable and the produce requires more care and protection, and at the season of rabi operations the outlying ponds are often dry and grass is scarce. It is therefore usual to find that the land nearer the village, which is generally the land that has been longest cultivated, is always sown with rabí; and in some of the older Sikh villages a circle of land close to the village rarely produces anything but rabi, the land farthest away from the village usually produces kharif crops only, and the belt between these produces sometimes kharif and sometimes rabí. This is however only a tendency and is often overruled by other considerations. Some sandy fields, especially the land in the drainage-channels which cross the district, rarely produce anything but rabi; and sometimes the question whether a peasant will sow kharif or rabi in a particular field is determined by his personal conveni-

ence at the time rather than by the nature of the soil or regard for any particular course of cropping. As already noted, the Bágrís often sow no rabí at all in any circumstances, while the Sikh, whatever be the nature of the soil, sows as much rabí as will keep him and his cattle fully employed. But most of all, these arrangements are subject to be modified by the distribution of the rainfall. If the rain fails altogether, nothing at all is sown, and the land gets a whole year's rest. If the rain is favourable for the kharif sowings, as much land is sown for the kharif as there is time and means to sow. If the kharif fails and the rains promise well for the rabi, as much land as possible is sown with a rabi crop, and sometimes the stunted kharif crop is ploughed up to make way for a rabi. Thus the area sown, and still more the area harvested, for each crop varies very much with the rainfall: for instance, in 1877-8, 36 per cent. of the whole cultivation was rabí, and 1880-81 only 17 per cent. It is not common to give fallow land many ploughings, as the rests it gets owing to failure of rain seem sufficient to keep up its crop-bearing powers. Rotation of crops is almost unknown, and the same crop is sown year after year in the same land. Much land produces bájra year after year, and much produces barley and gram without any change. The only crop not usually repeated on rain-land is jawar which is considered to exhaust the land; and if a kharif crop is taken after jawar it is generally bajra, moth or til; but sometimes land which has borne a jawar crop gets a rest for a year and is then sown with rabi. When light land is cultivated for the first time, it is sometimes sown with barley alone to begin with, and afterwards barley and gram. In the rice-kunds on the Ghaggar a crop of rice is taken every year in which the floods are sufficient to allow of its being sown; it is only rarely that wheat or gram is sown in rice-land, and the only rest the land gets is when the floods fail. The wheat and gram lands also, both on the Ghaggar and Satlaj, bear the same crop year after year when there is a sufficient flood. Lands irrigated from wells when manured sometimes give two crops in the same agricultural year. On such lands after sugarcane a fallow of six months is given, and then a crop of wheat is taken. Maize also is followed by wheat. After vegetables the land lies fallow for a year, and then wheat is sown. After tobacco comes a six months' fallow, and then wheat. Onions are sometimes followed by turnips; and after wheat jawar is sometimes sown for fodder. But most of the land on the wells produces wheat year after year, and in the whole district the area which produces a kharif and rabi crop in the same agricultural year (dofasli) is quite insignificant.

Agricultural implements and operations. kind of animal that is to draw it, from the large plough drawn by oxen to break up hard land for the rabi crops to the small plough (haleri) drawn by donkeys or women in the sandy soil of the uplands; but the plough generally consists of a wooden boot (hur or chan) in front of which the iron

plough-share (phálá) is fastened, while to an upright shaft let into the boot a small peg is attached as a handle; another shaft attached to the boot stretches forward between the bullocks and is tied to the yoke (panjálí) resting ou their necks. The donkey-plough is yoked in the same way, but is much smaller than that drawn by bullocks and is only rarely used by the Kumhárs. Sometimes the light sandy soil of the Rohi is ploughed by a small plough drawn by a single bullock which pulls it by two poles attached to the plough as shafts, and passing one on each side of the bullock to a collar on its neck. The camel-plough, which is very commonly used by the Bágrís in sowing kharif crops in light soil, is smaller than the ordinary bullockplough and is drawn by one camel by means of a rope attached to the shafts of the plough and to a cross-beam from which a rope passes along each side of the camel to a small saddle fastened round his hump. The ploughman directs the camel by a pair of very long reins fastened to the rings in the camel's nose. The ploughing in the Hitar is all done with bullocks, a pair of which can plough about a third of an acre of the hard soil in a day, the plough usually penetrating about six inches into the ground. An ordinary two-bullock plough can turn up in a day about three-fourths of an acre of the loam of the uplands when penetrating six inches into the ground in preparing it for rabí cultivation, and 12 acre when penetrating only three inches to prepare it for the kharif. A camel can plough in a day about two acres for the kharif, the plough penetrating only three inches into the ground; and the light plough drawn by donkeys simply scratches the ground. In sowing with the drill, a hollow bamboo (nálí or por) is attached to the upright shaft of the plough with its lower end in the ground just behind the boot, and with its wide mouth at the plough-handle so that the ploughman can feed it with the hand with which he holds the plough. That hand he keeps supplied with seed from a bag slung behind him, so that he is kept fully occupied holding the plough, guiding the bullocks, giving their tails a twist or their backs a stroke now and then, and at the same time keeping the drill regularly supplied with seed. As the soil has generally been prepared beforehand unless it is sandy and light, a plough can sow more land than it can break up in preparation for sowing, Thus in the Hitar a pair of bullocks can sow about half an acre for the rabi in a day; and in the Rohi about two acres for the kharif or one acre for the rabi, the furrows being much closer in sowing rabi . than for kharif crops. A camel can sow about two acres in a day, and a single bullock about three-fourths of an acre.

The soldod or clod-crusher is simply a log or beam which is drawn over the field by bullocks while the driver stands on it to increase its weight. It pulverises the clods, levels the ground and gathers together some of the weeds. It is not ordinarily used in kharif cultivation in the uplands, but is employed in preparing the land for a rabi crop and in levelling the ground after the seed has been sown. The karáhi is a board with a handle attached to it at right angles, held by one man while another holds two ropes attached to its ends; it is used for making the boundaries of fields or of plots for irrigation

(kyárí) one man standing on one side and pushing up the earth while the other man on the opposite side helps him by pulling the ropes. The dandráli is a sort of wooden rake used chiefly for raking manure together. In reaping (wadhan) the dátri or datti is used, a sickle with a curved blade like the hook sometimes used by reapers in Scotland, but with a saw-edge. The reaper squats on the ground and cuts the crop in handfuls at a time, tying it in sheaves as he moves along. This is the usual way of cutting the rabi crops, but in reaping jawar and bajra it is usual to ent off the ears only (dunggan) leaving the stalks standing to be cut afterwards. A man can reap about a fourth of an acre of rabi crop in a day, and sometimes when there is s great demand for labour, wandering reapers get as much as 3 annas each and their food for a day's work. The sheaves or ears seldom require to be dried, but are carried at once to the threshing-floor (pir) which is simply a piece of bare hard ground swept clean for the purpose. Here they are heaped round an upright stake round which bullocks are driven so as to tread out the grain and separate it from the straw; while a man with a pitch-fork (tringal or tangli) heaps it up under their feet. Sometimes as many as 15 bullocks are driven round the stake together. When the grain and straw have been separated, the straw is removed with the pitch-fork and the grain drawn together with the sabarkatta or wooden scraper, and if necessary again trodden out. The grain and chaff are separated by winnowing with the chhaji, a sort of tray made of the thin part of the stalk of sarr grass. One man fills the chhajj from the heap and hands it to another who stands up over a bare spot of the threshing-floor and shakes out the stuff gradually so that, as it falls to the ground, the wind may blow the chaff some distance while the heavier grain falls straight down. The chaff and stalks are then swept to one side and the grain to another. For winnowing a dry day and a fairly strong wind are required, and these are rarely wanting in Sirsá at harvest-time; too strong a wind blows part of the chaff beyond the threshing-floor so that it is lost. Winnowing is the special duty of the Chuhra, where he helps in harvest operations, and the chhajj is generally made by him. When the grain is winnowed it is put up into a heap, which is sometimes protected from evil spirits by having a circle of ashes traced round it, or by having a slit stick stuck into it with a piece of paper put in the slit on which a verse of the Qurán has been written as a charm. If the owner takes rent in kind and is not ready to divide it, little bits of wet mud are stuck all over the heap and a Chuhra (the thapi) stamps them with a small wooden stamp (thappa) that no one may take away the grain before it is divided. When all are ready, the division is made in presence of the parties interested, and each takes away his share. In some villages it is the duty of the Kumhar to bring the grain in from the field to the village, but often it is brought in by the peasant himself in his cart or on his camel. It is then stored in the 'earthen receptacles (bharola or palla) until wanted for consumption or to be taken to market. Barley and bajra keep longest, sometimes as long as 15 years, but have to be protected from the attacks of white ants by being kept above the ground in a raised bin (bharola) or granary (bukhárá

or burj). Barley is also liable to be attacked by weevil (súsrí). Jawár only keeps for two years or at the most four, and is also attacked by the súsrí. Gram keeps for four or five years and is attacked by a different weevil called dhorá.

The straw of jawar and bajra is put up in large stacks, sometimes near the homestead, but generally in the field. The chaff and broken straw of wheat and barley, and the leaves of gram, moth, mung and pala are generally brought to be stored in the homestead, but often left in a heap on the field with a covering of thorns and surrounded by a ditch to protect them from the cattle, or put up in a stack thatched with bajra straw to keep out the rain. Jawar and bajra straw are chopped

into little pieces for fodder with the chopper (toká).

Other implements are the kulhárí or hatchet used for cutting wood; the pháorá or wooden scraper used for scraping up manure, &c.; the gandása which consists of a thin broad blade fastened at an oblique angle to a long handle, and is used for cutting thorns and low brushwood by a sidelong blow with the right hand while the left hand catches and tosses them up with a pitch-fork; the khurpa, a small trowel used for digging up grass and weeds; the kahí or kassi, a shovel-mattock, the broad blade of which is placed at an acute angle to the short handle; this does the work of a spade, but instead of being pushed outwards and upwards like a spade, it is struck downwards and pulled inwards like a hoe. Carts are little used in the district except by the Sikhs and the Ráíns, who have fine large four-bullock carts and sometimes go long distances with them for hire. According to the enumeration made in 1880 there were then 2,013 carts in the district, of which 1,172 were for two bullocks and 841 for four bullocks.

169. The use of manure is almost unknown in the Sirsa district; indeed, the report furnished in 1878 to the Famine Use of manure and Commission gave the percentage of area annually deterioration of soil. manured as only '014 of the irrigated area, which is itself only 6 per cent. of the cultivated area; and this is much less than the proportion in any other district of the Province. Manure is not used at all in the Dry Tracts, Chaks Bagar, Rohi and Utar. It is only rarely applied to lands irrigated from wells and to rice-The best rice-cultivators think it enough to give their lands a cart-load of manure per bigha before the rains every second year, i.e., about two tons to the acre, and they double this quantity if a layer of soil has been taken off the field to make it low enough for irrigation. For rice the best manure is the dung of sheep and goats scraped up at long intervals from the folds where they are shut in at night. In the Ghaggar valley land irrigated from wells often gets manure every year, one or two cart-loads per bigha of rubbish taken from the general heap of ashes, sweepings and waste outside the village exposed to sun and rain. In the Satlaj valley some lands irrigated from wells get manure once in two years, and especially when sown with tobacco or sugarcane. Maize, turnips and wheat are said to get 50 tons of manure per acre; pepper, tobacco and onions twice as much; and sugarcane thrice as much; but probably the land is excep-

tionally lucky that gets this quantity of manure. On the lands flooded by the Ghaggar and Satlaj the silt annually deposited by the river-floods to some extent refreshes the soil and renders but a considerable area of land on manure less necessary, the Ghaggar which formerly produced a wheat crop annually will now produce nothing better than gram. In the Dry Tract so much of the cultivated land has been only lately brought under the plough that of a great portion it may be said, in the words of a former Deputy Commissioner, that it is a virgin soil requiring only to be tickled to make it smile with blooming harvests. The people however with some show of reason say that after five years or so of cultivation the land deteriorates and requires more labour to make it produce crops like what it did at first, and that after twenty years' cropping the kharif produce is so poor that it becomes necessary to cultivate the land more carefully and grow vabi crops. The often-recurring enforced fallows due to the failures of rain must help to restore the fertility of such land. In 1838, Major Thoresby estimated the average produce of the Dry Tract at from 3\frac{1}{2} to 5\frac{1}{2} maunds per acre, which is not far from the present average; and there cannot have been any very great deterioration in the productive power of the soil, though no doubt there has been some, the effect of which has been to make the better soils first brought under the plough more nearly equal to the soils which being somewhat inferior were broken up at a later date. In the Dry Tract the fields are comparatively free from weeds, which there cause no trouble; but in the irrigated lands of the Ghaggar and Satlaj the camel-thorn (janwása), the wild onion (pivázi), the thistle-like kateli and leh, and other weeds, are very common and often do great harm by exhausting the soil and choking the crops; indeed many fields seem to produce more weeds than grain. Yet, except in the rice-fields, the people seldom make any attempt to eradicate them, and as weeding hardly finds a place in the agricultural operations of the Sirsa peasant, it is probably true that weeds are annually becoming more numerous and troublesome.

170. When a crop has been sown, the outturn depends chiefly on the quantity and seasonableness of the rainfall it Causes reducing the gets when growing and ripening, but there are outturn of crops. other things which affect the result. The rabi crops, and especially gram and sarson, are apt to suffer severely from late frosts which prevent the fruit from forming and sometimes kill the whole plant. In 1868 Mr. Oliver estimated that severe frosts had reduced the outturn of the rabi by one-third, and a severe frost in February 1882 greatly reduced the outturn of the rabí harvest of that year. Late jawar and moth are also sometimes injured by early frosts. Both kharif and rabi crops often suffer from dry hot west winds blowing while the grain is forming, and shrivelling up the grain so as to make the outturn much lighter. For instance, in 1876 a hot wind in September reduced the produce of the kharif, and in 1880 a hot west wind in March reduced the rabi outturn throughout the district, the loss being in some places estimated at one-fourth. Sometimes, but rarely,

a hail-storm in March or April passes over a tract of country, laying low the ripening grain and quite destroving the crop. instance, in 1876 remissions were granted to five villages on account of damage done by hail. Such storms, however, affect only a few villages at a time, and they seem to be very rare in this district, not visiting any village oftener on the average than once in 20 years. The crops are also very liable to attacks of animals. Antelope, which are very common in some parts of the district, do great damage to the crops and much trouble is taken to preserve the growing grain from their ravages, especially in and near the Bishnoi villages and along the Bikaner border. Field-rats are always numerous enough to do some damage to the crops, but sometimes they suddenly multiply enormously, and devour almost the whole produce of the fields; for instance, in 1876 they swarmed in several parts of the district and even filled the houses. On such occasions the villagers are practically helpless and do little more than look on and wait until the rats disappear, which they do as suddenly and as inexplicably as they increased. The kharif crop especially is liable to the attacks of birds of many sorts, parrots, crows, doves, starlings, sparrows, goliyas, sarágás and other small birds in flocks. To protect the crops, the peasants erect platforms from which a watchman with his sling or rattle or rope to be cracked like a whip and his shouts and cries frightens off the depredators. Flights of locusts (tidi) often come up from the west in the rainy season and disappear towards the east where they are supposed to die from eating saline earth (reh). Such swarms do great harm to the crops where they settle for the night by devouring everything eatable; and even if the crop recovers somewhat and grows again, both grain and straw are bad and The greatest damage is done where they lay their eggs, for after 21 days the young locust (phakka) appears above ground and prepares for his devastating career. It is while his wings are gathering strength that he does so much damage to the neighbouring crops, but at this stage he is easily killed. A shallow trench is dug and the young locusts driven into it, beaten down and buried alive with earth. If he escapes this treatment, he flies about a month after appearing above ground. The villagers near the Bikaner frontier complain that they suffer from the carelessness and want of system of the Bikaner people in killing the young locusts; but if the crawling army sometimes extends, as it is said to do, 20 or 30 miles without a break, the people are to be excused for letting some of them escape. The worst flight of locusts in late years was in 1872, but there is hardly a year in which they do not visit the district, and on the average each village endures a flight of locusts about once in 20 years. In dry weather the roots of all crops are liable to be attacked by the white ant (syonk or dimak), especially wheat and gram in hard dry ground. A kind of grasshopper (toká) eats sprouting bajra and jawar, and sometimes wheat and barley. A small insect (teliya) attacks the leaves of autumn crops which are also sometimes eaten by a caterpillar (khtra) and other insects, most of which are said to be produced by excessive rain and east wind and to be killed

by a dry west wind. One of them, the karwa, a flying insect which injures the bájra blossom, is got rid of by a man taking his sister's son on his shoulder and feeding him with rice-and-milk while he repeats the following charm:—"Máme charhke bhánjá áyá—nikal karwa khet paráyá," i.e., "The nephew has got on his uncle's shoulder—go away karwa to somebody else's field." A white maggot (gindár or baggá kíra) eats the heart of the jawár stem where it joins the roots and injures the plant so that it produces no grain. Rice is injured by a bluish insect (kara) which sucks the juice of the plant. Other insects also attack the crops and they are liable to diseases, such as rust (kúngí) which destroys or diminishes the produce of wheat and barley; it is caused by continuous damp weather and is cured by sunshine or strong wind.

Mode of observation of harvests.

Since the commencement of Settlement operations in 1879 a careful field-to-field inspection has been made of each harvest, and an estimate framed of the

outturn and its value at the prices of the day. The areas returned by the patwaris as cultivated with each crop were carefully checked by the supervising staff, and I believe them to be very approximately correct. In order to estimate the average outturn. a number of fields were selected by the Superintendent or tahsíldár such as he considered to represent the average outturn of the assessment circle for that harvest, and their actual produce was weighed and reported; and after comparison of the averages given by the different fields so chosen and enquiry from the Superintendents and the peasants checked by my own observation of the crops, I formed a rough estimate of the average outturn of that harvest for each crop in each assessment circle. In a tract where the produce of the fields varies so immensely as it does in Sirsa, it would be very difficult even for a skilled farmer after an inspection of the whole tract to say what fields represented the average of the harvest, and although the fields were of considerable size (often more than 10 acres in area) and were chosen with care after a good deal of inspection and enquiry, the estimate of average outturn so framed must be only a very rough one, and I took care always to err on the safe side by assuming the average outturn as something less than the observations would seem to warrant. Having ascertained the gross produce of the harvest by multiplying the area sown with each crop by the estimated average outturn of the crop, I made an estimate of its gross value by ascertaining the prevailing prices of the time in the several towns and large villages, and after making full allowance for cost of carriage, applying those prices to the estimated gross produce. I submitted for each harvest a detailed report giving these calculations and discussing the condition and prospects of the district, the export of grain, the state of the cattle, and the realisation of the land-revenue of the year; and as these are the most trustworthy statistics available for an estimate of the average produce of the district, I give below a short account of each of those harvests.

Rabi harvest 1880. 1879-80 was as follows (in inches):—

	Month.		Sirsá.	Dabwálí.	Fázílká.
April	•••				
May		•			
June	•		1.2	2.9	2.5
July			1.7	2.	1.1
August			5-9	4.4	3.6
September	•••		2.	1.0	.8
October				-1	
November					•••••
December			.2	1.4	.4
January					•••••
February			•5	.7	.9
March			•		
	Total	w.	11.5	12.5	9-3

This is below the average everywhere. The rainfall in June and July was much lighter than usual, and the kharif crops did not get a good start. Very little rain fell until near the end of August and by that time the kharif crop had almost wholly dried up. Except in a few villages in the south of Fázilká tahsíl, which had a fair crop, the kharif was everywhere very poor, and in most villages a total failure. The Ghaggar did not come down in good flood and the rice-crop was poor. But in the end of August and in September there came good rain, which moistened the soil well for the rabi sowings, and a larger area was sown for the rabí than had ever been sown before. Some villages sowed rabí for the first time, and much land on which the kharíf had failed was ploughed up and sown for the rabi. The crop sprouted well. and a most opportune fall of rain in the end of December and again in February brought it near maturity with great promise. Unfortunately in March and April, just as the crop was ripening, a strong hot west wind blew for several days and dried up the grain, which thus lost much of its weight; and the crop which had promised to be a bumper everywhere in the Dry Tracts was brought down nearly to the average. It suffered from no other evil however-no trouble of hail or blight or insects-and in the three Dry Chaks both the area and the outturn were above the average and the crop was an unusually good one, especially

about Dabwálí where the rain in December was heavier than elsewhere. As always happens, however, a dozen or twenty villages, this time to the west of Rori were an exception to the general good fortune and had a poor rabí. The Ghaggar floods irrigated much less than usual, and the area of flooded lands under crop for the rabí was much below the average, but the outturn of wheat was fair and of gram good. The area irrigated by the Satlaj floods was also much below the average and the outturn too was poor, so that the rabí crop in the Hitar was a very poor one. But the gross produce of the rabí for the district as a whole was decidedly above the average.

The areas returned were as follows (in acres) :-

Areas sown for Rabi 1880. .

Assessment Ci	rcle.	Soil.		Wheat.	Barley, gram, oil- seeds, &c.	Total.	Total of Assess- ment Circle.
Bágar		Báráni Cháhi		206	12,432 274	12,432 460	12,432
Náli	}	Rez		6,360	3,967	10,357	
Robí	1	Báráni Báráni		1,057	19,114 2,07,611	20,171	30,978
		Cháhi		4,993	2,01,011	2,12,604	2,12,604
Utár	{	Báráni		880	10,180	11,060	11,125
Hitár	{	Cháhi		4,136	630	4,766	
	1	Rez		2,793	e1,605	4,398	9,164
Total of district	1	Cháhi Rez	- 1	4,398 9,153	913 5,572	5,311 14,725	•••
10tal of district)	Báráni		6,930	2,49,337	2,56,267	•••
Grand total	***	Grand total		20,481	2,55,822	2,76,303	2,76,303

The actual produce of 44 fields chosen by the Superintendents was observed, and after comparison of all available data and enquiry from the people, I estimated the average outturn of the different crops for that harvest as follows (in maunds per acre):—

		GR	AIN.	STRAW.		
Assessment Cir-	Soil.	Wheat.	Barley, Gram and Oilseeds.	Wheat.	Barley, Gram and Oilseeds.	
Bágar	Bárání Cháhi	.;	4	**	10	
Nálí .	Rez	6	10 8 5 5	6 5	6	
	Báráni	6 3 3 3 9	5	5 5 5	5	
Robí	Barání	3	5	5	5	
Utár	Báráni	3	4		1	
Hithr }	Cháhi	9	10	9	10	
	Rez	6	8	6	8	

This gives the gross grain produce of rabí 1880 approximately as follows:—

			GROSS PRODUCE OF GRAIN IN MAUNDS.						
Asses	SMENT CIRCI	E.	Wheat.	Barley, Gram and Oilseeds.	Total				
Bágar Nálí Rohí Utár Hitár	•		40,000 14,000 3,000 51,000	48,000 1,22,000 10,00,000 38,000 14,000	48,000 1,62,000 10,14,000 41,000 65,000				
	Total		1,08,000	12,22.000	13,30,000				

In order to estimate the value of this grain the actual market prices of the 1st June 1880 of the eight chief towns and villages throughout the district were ascertained, and the lowest prices on that date were 20 sers per rupee for wheat and 40 sers per rupee for barley and gram. Allowing for the cost of carriage I estimated the average harvest prices as follows:—

Alexander of the same	AVERAGE PRICE IN I	RABI 1880 (SERS PER RUPRE.)
Assessment Circle.	Wheat.	Barley, Gram and Oilseeds,
Bágar		42 •
Nálí	. 20	40
Rohí	. 20	40
Utár	. 20	36
Hitár	2.2	32

These rates applied to the estimated gross produce gave the following as the approximate money value of the grain produce of the rabi crop of 1880. I add an estimate of the value of the straw similarly calculated, the rates taken being from five to seven maunds per Rupee for wheat straw, and from four to six maunds per rupee for barley straw.

Assessment Ci	rele.	Value of Grain.	Value of Straw.	Total.
		Rs.	Rs.	Rs
Bágar	***	46,000	12,000	58,000
Nali	•••	2,02,000	38,000	2,40,000
Rohi		10,38,000	1,70,000	12,08,000
Utár		48,000	10,000	58,000
Hitár	•	1.10,000	11,000	1,21,000
Total Rs		14,44,000	2,41,100	16,85,000

This estimate gives the value of the total produce of the cultivated land in rabí 1880 at the market prices of the day as Rs. 16,85,000, which is about nine times the former assessment of the whole year, and six times the new assessment of the whole year. There was little difficulty in realising the revenue, and on 31st October 1880 the balances of land-revenue were only Rs. 730 and of Takkávi Rs. 122.

Kharif harvest 1880.

173. The rain-fall of the year 1880-81 was as follows:—

Month.		Sirsá.	Dabwáli.	Fázilká.
April			•••	
May		-1		-5
June		3.9	4.1	4.3
July		3.9	-1.6	2
August			26	-3
September		-6	.6	
October	•••			
November				
December		.7	.9	. 5
anuary		.1	-1	1
ebruary			1.	-8
March		2.8	1 .	.7
Total		8.9	11.9	9.2

In the end of June and the beginning of July there was an excellent fall of rain over the whole district. Every plough was set to work and as much land as possible was sown while the soil remained moist. The total area sown was, notwithstanding the large area sown for the previous rabi, somewhat above the average, especially in the Robi and Utar circles; but in the Nali circle owing to excessive floods of the Ghaggar in July many of the rice-kunds were overtopped and drowned and the area sown with rice was very small; on the other hand an unusually good flood from Fathabad irrigated a large area of Sotar land east of Sirsá in the Choya valley, and enabled it to produce good jawar crops. In July the prospects of the kharif harvest were excellent everywhere in the Dry Tracts, but with the exception of slight and partial showers no rain fell in August, September or October, and the promising crops dried up. In many villages the failure was complete and not a straw grew worth the gathering; in many there was some straw but little grain; and in only a few did the partial rain-showers bring the crop to maturity, and produce a fair harvest of grain. According to the patwaris' inspection, which was checked on the field by the supervising establishment, of the 7,27,200 acres sown for the kharif only 3,64,746, or about half, produced any grain, and in Chak Nálí less than a fourth of the area sown produced a crop; indeed, in great part of that Chak every blade of crop sown on unirrigated land dried up, and came to nothing. In more than a third of the villages in the district less than a fourth

of the area sown produced a crop, and in 140 villages the failure was so complete that nine-tenths of the area sown produced no grain. The Bágar Chak suffered least, as its sandy soil produces a crop with very little rain, and a few opportune showers in September were enough to save something of the harvest. In the Rohi Chak some villages southeast of Chautála had a fair crop, but elsewhere, with the exception of a few fields here and there, very little grain was produced. Even where the crop came to something the outturn was as a rule very poor. Sixty-two fields averaging 12 acres each were selected by the Superintendents as representing the average outturn of the harvest, their actual produce was weighed, and from the results of these experiments and my own observations and enquiries, I framed a rough estimate of the average outturn. The statistics and estimates are as follows:—

		AR	EA REPORT	RD.	AS HAVING	PRODUCED	A CROP (IN ACI	RES.)	
Assessment Circle. Total area sown for Kharif 1880.		Total area		Jawar (alone and with pul- ses.)	Bajra (alone and with pul- ses.)	Moth, Mung, Mash, and Gwar	Til	Rice.	Miscellaneous.	
Bagar	1,18,954	80,911	Barani	•••	2,493	76,643	1,734			1
Nali	1,29,420	29,611 {	Rez		96 3,936 2,052	2,317 15,580	2,518 1,609	275 110	994 34	86 1 1
Rohi	4,35,607	2,26,363	Barani		• 30,846	1,61,988	27 156	6,328		45
Utar	37,898	24,160	Barani		2,473	19,906	1,342	430		9
Hitar	5,321	3,7.1	Rez		1,361 197 138	36 197 759	15 205 57	98 24 36	33	444 101
Total of district	7,27,200	3,64,746		_	43,582	2,77,426	34,688	7,301	1,061	688

My estimate of the average outturn and the resulting approximate gross produce were as follows:—

	ESTIMATED AVE	PER ACER.		ESTIMATED GROSS PRODUCE IN MAUNDS,			
Assessment Circle,	Soil.	Jawar, &c	Bajra, &c.	Jawar, &c.	Bajra &c , including pulses.	Miscellaneous	Total.
Bagar { Nali { Rohi } Utar {	Barani Barani Barani Chahi Barani	2 t 1 t 1 t 1 t 1 t 1 t 1 t 1 t 1 t 1 t	2 }	1,000 11,000 46,000 4,000 8,000	39,000 17,000 2,53,000 51,000	12,000 10,000 1,000 4,000	40,000 40,000 3,09,000 36,000 13,000
Total of the district		-		70,000	3,41,000	27,000	4,38,000

The war in Afghanistan had created a brisk demand on the frontier, and over six lakhs of maunds, chiefly the produce of the previous good rabí harvest, had been exported through Fázilká during the previous year; but although owing to the two bad harvests the stocks of the agriculturists had become low, there was still a considerable quantity of grain in store in the towns. Prices remained at a high figure from August 1880 to June 1881, when the lowest market-prices quoted were for wheat 20 sers and for barley 29 sers per rupee. The prices adopted and the resulting estimate of the value of the produce of rabi 1881 are given below together with an estimate of the value of the straw. The produce of straw on unirrigated land I estimated at two maunds per acre for wheat and 21 for barley, &c., and on irrigated land about six maunds per acre for wheat and ten maunds for barley. Owing to the failure of the previous kharif there was a fodder famine in the district, and fodder was selling in the towns at prices at which a few years before grain itself could be got; barley and gram straw sold in Sirsá and Fázilká at a maund and a half per rupee. The prices I adopted for valuation were three maunds per rupee for wheat straw, and two maunds per rupee for barley straw. The resulting values are as follows :-

Assessment Circle.		Harvest (sers per	Prices Rupee.)	Value of Grain pro-	Value of Straw.	Total.	
			Wheat,	Barley &	duce.	•	
€ Bágar			20	30	Rs,	Rs.	Rs.
Nálí			20	30	1,45,000	21,000	1,66,000
Rohi	***		18	28	4,96,000	1,44,000	6,40,000
Utár	***	***	20	25	21,000	5,000	26,000
Hitár	***		20	25	1,32,000	20,000	1,52,000
Total of	district				7,94,000	1,90,000	9,84,000

This estimate makes the value of the rabi produce of 1881 Rs. 9,84,000, which is five times the former total land-revenue of the year, and more than three times the new total assessment. As two successive harvests had been more or less failures, I considered separately the condition of each village in the district with regard to its ability to pay the rabi instalment, but only eleven villages seemed to require a suspension of the instalment; and on 30th November 1881 the total balances were Rs. 5,001 suspended from kharif 1880 and Rs. 1,782 suspended from rabi 1881—total Rs. 6,783. Thus notwithstanding the failure of three harvests out of four, the lightness of the demand made it possible to realise almost the whole of the revenue without hardship to the people.

In June 1881 the district was in a somewhat critical condition, for the kharif and rabi harvests had both been poor and the stocks of

grain and fodder were low; and had the rains again failed there might have been a scarcity and a serious loss of cattle. There would probably not have been a famine, for the area which had suffered was small and scarcity prices would have caused a large importation. The people as a body were still well enough off to purchase grain from without, and the poorer classes are so ready to migrate that so long as work and food are within easy reach they readily go to it, and in June 1881 half the population of many villages in the Fázilká Rohí had wandered westwards; but such migrations are not necessarily a sign of distress. When the rains fail the cultivator has nothing to do at home, so he takes his wife and children and drives off his cattle and sheep towards the river where at the time of the rabi harvest he and his family find plenty of work and live in luxury on wheaten bread, while his cattle are allowed to graze free of charge. Many such families crossed the Satlaj into Montgomery to help in reaping the unusually good harvest there and returned to their homes as soon as The cattle held out wonderfully notwithstanding the drought. In August 1880 they had been at their best with plenty of water and grass everywhere; but after that time for want of rain the grass gradually dried up and the cattle grew thinner and thinner. There was no grass within reach in Bikaner, and though some cattle were driven to the banks of the Ghaggar and Satlaj, few left the district. Partial showers here and there produced a little grass, and the rain of March and April was a great boon to man and beast by filling the village-ponds and giving them a supply of water, which is generally got with great difficulty in the hot weather. In many villages there were stacks of bajra straw of several years back, and these were the means of saving many cattle. However, few cattle died though they were getting much reduced and the stocks of fodder were almost exhausted, and had the rains of 1881 held off much longer, many thousands of cattle must have died as in former seasons of drought.

Kharif harvest 1881. 175. The rain-fall of 1881-82 was as follows :-

	ala ya Mareta Ya	Month	.			Sirsá.	Dabwáli.	Fázilká
April		enter an in Ma	Mark 1 1911			.7	100 L	•6
May			Charles on the State		•••	-3	•3	•8
fune		13 - L. 14 14 5 14	***		Constitution of the Consti	.3		1:4
uly	•••					12.8	9.6	9.6
Lugust					anagar Mar	9.4	7	5.
leptember	il.		•••		•••			-9
Detober	•••		•••	***	***	***		
November	•••		***	•••	***	•••	***	•••
December		10 May 10 10 10 10 10 10 10 10 10 10 10 10 10	***		***	***	10000000000000000000000000000000000000	***
	***			1 7 TO 10 IN	Seers Mile		2.1	***
lanuary	***	•••	***	***	***	2.5		2 3
February	***	•••	***	•••	***	.3	-5	·KE MA
March	**		•••		•••		•••	***
	Tale of	• Total	NE CONTRACTOR	Sala Artis		26.3	20.5	20.6

This was an exceptionally heavy rainfall, almost as much as in the previous two years put together. No such heavy rainfall has been recorded at Sirsá since the mutiny, and only once during that period has there been so heavy a fall at Dabwálí or Fázilká. Some little rain fell in April, May and June, but up to the 9th of July there had been no fall of rain sufficient for the kharif sowings, and things were getting very critical. During the following two months, however, there was such a heavy and continuous fall of rain as has not been seen in this district for many years and the general complaint was that there was too much instead of too little. In the lighter sandy soil of the Bagar and Rohi the seed was washed out of the ground by the violence of the downpour and in many fields had to be resown twice or even thrice before the young plants developed sufficiently to retain their hold on the soil; and in the stronger soils a heavy shower of rain falling within two or three days after sowing made the soil cake so firmly that the young shoots could not force their way through. The people did not fail to make the most of their opportunities to sow as large an area as possible, and the area sown (7,72,858 acres) was larger than had been cultivated for the kharif during the previous six years. Besides the large area which suffered from too much rain at seed-time, a considerable area of crop which started well suffered from want of rain later on to bring it to maturity, as practically no rain at all fell after the end of August. The moisture which the soil had retained sufficed to develope the straw to an unusual degree, and a little rain towards the end of September would have enabled the ear to form well, and we should have had such a bumper crop as Sirsá never saw before; but the premature cessation of the rainfall made the outturn of grain much lighter than might have been expected from the favourable rain at seed-time, except in a few favoured tracts which got some partial showers. In those villages the harvest was really a bumper, even as compared with those of districts generally much more fertile than Sirsá; but as a rule throughout the district, while the outturn of fodder and straw was far above the average, the grain produce was not so very much better than in an ordinary kharif harvest. Still the contrast between this kharif and that of the previous year was a great one. In 1880 the rains had been almost equally favourable for sowing, but they commenced about a fortnight earlier than in 1881, and ceased about a month earlier, so that throughout the greater part of the district the crops dried up entirely and produced neither grain nor straw; while in 1881 hardly an acre sown failed to produce some little grain and a fair amount of straw. Even in this year of general prosperity, however, some tracts were denied the good rainfall granted to their neighbours, and 42 villages were returned as having had a poor crop, chiefly in two small tracts, one in the north corner of the Dabwali tahsil and the other in the west of Fazilka tahsil. The floods of the Ghaggar were on the whole favourable for the kharif, and as the rice-cultivators had taken care to repair the embankments of their kunds which had been breached by the heavy floods of the previous year, they were able to sow a large

area of rice most of which produced a crop. In the Hitar also the area sown was unusually large; the jawar crop was at first very promising, but the failure of the latter rains prevented the ear from forming, and although there was plenty of straw everywhere, there was comparatively little grain.

The area cultivated for the kharif of 1881 was as follows :-

	No. 1780		ARBA CULTIVATED (IN ACRES.)								
A-cossment Circle.	Soil.		Jawar (alone and with paises.)	Bajra (alone and with pulses)	Moth, Mung, Mash and Gwar.	Til.	Rice	Miscellaneous.	Total.	Total of Assess- ment Circle.	
Bagar Nali Rohi Utar Hitar	Barani Chahi		10,018 74 652 14,996 1,02,418 7,143 2,653 895 962	3,466 1,06,755 2,84,357 29,230 121 386	4,023 7 152 5,581 55,286 3,652 38 46 178	40 900 20,502 1,913 216 320 237	6,740	31 3 8 38 31 313 84	1,21,738 119 11,053 1,28,248 4,62,601 41,949 3,341 1,731 2,078	1,39,420	
To	otal of the distr	iet	1 39,811	5,32,717	68,963	24,130	6,748	489	7,72,858	7,72,856	

My estimate of the average outturn and resulting average gross produce were as follows:—

				PER ACRE		ESTIMATED GROSS PRODUCE (IN MAUNDS.			
Assussi	STATE OF THE PARTY	• Soil.	•	Jawár, &c.	Bájra, &c	Jawar (alone and with pulses,)	Bájra and pulses.	Miscellaneous.	Total,
Bagar	***	Barani	***	1	1	10,000	1,12,000		1,22,000
Nali		(Rez) Barani	***	4	24	63,000	2,98,000	71,000	4,32,000
Rohi Utar	•••	Barani Barani		4	1	4,10,000 29,000	12,48,000 1,28,000	51,000 6,000	1,63,000
Hitar	2007	(Chahi Rez	***	3 3	2}	20,000	3,000	2,000	25,000
Total of th	e district					5,32.000	17,89,000	1,30,000	24,51,000

The high prices which had ruled from August 1880 when the failure of the rains showed that the kharif of 1880 would be a poor harvest continued pretty steady until the end of May when they rose still higher; and in June and July when the rain seemed to hold off prices rose rapidly until on 31st July, in Sirsá market, barley was selling at only 23 sers per rupee, jawár at 22, bájra at 20, and gram, the cheapest grain, at 25; these were, for Sirsá, scarcity prices. The rains of August caused a rapid fall of prices, and they continued to fall as the harvest became assured, and more especially after it became

certain that the rabi would be a good crop, and in six months the prices of barley and jawar fell by half, i.e., barley, which in July sold at 23 sers per rupee, sold in January 1882 at 50 sers, and the price of bajra fell from 22 sers per rupee to 46. On the 1st December 1881 the lowest prices quoted in the chief villages were for jawar 53 sers per rupee, for bajra 40 and for moth 49. I estimated the average outturn of jawar straw at about 8 maunds per acre, bajra and gwar 3 and moth 2 maunds per acre, and the average prices at about 12 maunds per rupee for jawar straw, 20 for bajra straw and 5 per moth and pala. These give the following as the approximate value of the kharif crops of 1881;—

Asser	sment Circ	le.	Harvest p	rices (sers pe	r Rupee)	Value of grain,	Value of straw.	Total.
	100 Y		Jawar.	Bejra.	Moth.	Rs.	Rs.	Rs,
Bagar	***		40	36	40	1,34,000	21,000	1,55,000
Nali			40	36	40	5,50,000	41,000	5,91,000
Robi	***		50	40	45	16,63,000	1,35,000	17,98,000
Utar			45	35	35	1,83,000	18,000	2,01,000
Hiter	***		45	85	35	26,000	10,000	36,000
Total of	the district					25,56,000	2.25.000	27,81,000

The value of the total produce of kharif 1881 is thus estimated at Rs. 27,81,000, or enough to pay the new assessment of the whole year nine times over. The new assessment came into force in the three Dry circles with effect from this harvest, and although the enhancements were 31, 81 and 47 per cent. respectively, the land revenue in those tracts was fully realised without difficulty.

176. Although almost no rain fell for four months after the end of August, the rainfall of July and August Rabi harvest 1882. had been so heavy that the moisture remained in the soil and enabled the people to sow a large area for the rabi. The crop was beginning to dry up when an exceptionally heavy fall of over two inches in January almost all over the district refreshed it and gave promise of a bumper harvest. Towards the end of February, however, a severe frost injured the crop in may parts of the district and greatly reduced the outturn. Had it not been for that frost, the rabi harvest would have given an amount of produce larger than was ever before given by a rabí harvest in this district, and even as it was, the outturn was excellent and much above the average. Almost every acre sown produced something, and no village which depends much on the rabi crop failed to reap some produce, though in a few scattered villages the harvest was much inferior to what it was elsewhere. In the Sotar valley east of Sirsá the wheat crop was exceptionally good both in area and in outturn, as the unusually heavy rain shad thoroughly soaked the hard soil and made it possible to sow an exceptionally large area of wheat. There were good floods on the Ghaggar in

August, but the late floods failed, and the villages below the Annakai Chhamb were unable to sow much land for the rabí; the Dhanúr and Annakai Chhambs, however, and the low land towards Ellenábád dried up early and left an unusually large area of moist land fit for the rabí sowings. This was benefited like the Sotar crop by the heavy rain of January, while it escaped being drowned by winter floods of the Ghaggar. The Satlaj, chiefly owing to the neglect of the inundation canals, did not flood much land, and the area irrigated on wells showed a slight falling off, but the heavy rainfall permitted the sowing of a large area without irrigation, and the outturn of many fields was very good. The areas sown were as follows (in acres):—

Assessment Circle.	Soil.	Wheat.	Barley, Gram, Oilseeds, &c	Total.	Total of Assessment Circle,
Bagar	Barani	8	2,041**	2,049	2,049
Nali	Chahi Rez	68	285 5,895	353 15,129	
	(Barani	4,780	10,143	14,923	30,405
Rohi	Barani Barani	5,683	189,453 10,791	195,136 12,061	195,136
Ular	Chahi	3,764	795	4,559	12,001
Hitar	Rez	3,807	877	4 684	
Company of the second	(Barani	641	594	1,235	10.478
Total of the district.		29,255	220,874	250,129	250,129

In order to ascertain the average outturn, 62 fields aggregating 558 acres were selected by the Superintendents as representing the average of the harvest and their produce weighed. The estimated average outturn and the resulting estimate of the gross outturn are as follows:—

Assessment Circle,	Soil.	OUTTUR	DAVERAGE N(MAUNDS ACRE).		OF GROSS OUTTURN (IN MAUNDS).			
CONTRACTOR		Wheat.	Barley, &c	Wheat.	Barley, &c.	Total.		
Bagar	Barani (Chahi		4		8,000	8,000		
Nali	Rez Barani	8 7 8	5 7	103,000	102,000	205,000		
Rohi	Barani	• 6	7	34,000	1,326,000	1,360,000		
Utar	Barani (Chahi	6 71 6	10)	8,000	75,000	83,060		
Hitar	Rez Barani	6	65	55,000	13,000	68,000		
Total of the district.		***	Table 100 and	200,000	1,524,000	1,724,000		

Owing to the failure of the kharif crop of 1880 and the rabi crop of 1881 and to the cessation of the war in Afghanistan export

had greatly fallen off, but the good kharif crop of 1881 replenished the stocks of the peasants and left them free to sell a large portion of The imports into the their excellent rabí crop in 1882. towns for storage or export greatly increased, and an especially large quantity of grain found its way into Sirsa town. There was a large demand to the east, where the rabi crop had not been so good, and many traders from Bhiwani or Dehli bought largely; and it was estimated that two lakks of maunds of grain were in a few months stored in Sirsá town alone. At the beginning of March prices were low, and barley and jawar were selling in Sirsa market, at 47 sers per rupee, but the demand from the east then sent prices steadily but slowly up, and at the end of July the cheapest grain' at Sirsá was jawar at a maund the rupee. At the beginning of June the lowest market prices were, for wheat 27 sers per rupee, barley 50, gram 45. The prices adopted and the resulting estimate of values are given below. The weight of the straw was taken, according to the roughly correct estimate of the peasants, as about equal to that of the grain, and the prices at about eight maunds per rupee for wheat straw and sia maunds per rupee for barley straw. The resulting values are as follows:-

* Assus	SMENT CIRCL	B.		RICES (SERS	GRAIN PRODUCE.	VALUE OF STRAW.	TOTAL VALUE.
	District Control		Wheat.	Barley, &c	Rso	Rs.	Rs.
Bágar	***	***	24	40	8,000	1,000	9,000
Náli			24	40	274,000	30,000	304,000
Rohi			28	50	1,110,000	225,000	1,335,000
Utar			25	45	79,000	16,000	- 95,000
Hitár			25	45	1,00,000	11,000	111,000
Total	of the district	***			1,571,000	283,000	1,854,000

This estimate makes the value of the produce of rabi 1882 Rs. 18, 54,000, or nearly seven times the total assessment of the year. The enhanced assessment was realised without any difficulty. In July 1882 after the first rains of 1882 had fallen, the district was perhaps in as prosperous a condition as it had ever before experienced. The cattle had had a year of plenty and were in excellent condition, and there was plenty of grass and water everywhere. Large stores both of grain and fodder had been laid by from the last two harvests, and many of the peasants had a considerable amount of cash in hand after their recent large sales of grain and cattle. The contrast between the state of things then and that of only a year before was marvellous.

Kharif Harvest 1882. 1882. The rainfall of 1882-83 was as

Month.	13	Sirsá.	Dabwálí.	Fázilká.
April			200	·2 ·7
May				
June		1.2	.6	•2
July		5.4	6.1	1.8
August		.2	1.	2.7
September		3.8	2.9	7.6
October	[******		••
November				
December				
January		1.5	2.3	1.9
Feburary			•4	
March		1.4	.5	.5
Total		13.5	13.8	15.6

The total rainfall was not far from the average, being slightly below it at Sirsá and slightly above it at Fázilká. Almost no rain fell until the 10th of July, and very little land had up to that date been sown for the kharif. During the next fortnight there was very little rain, and it was not until the last week in July that really good rain fell. In that week almost the whole of the district, with the exception of a small tract below the Danda near Fázilká, had two or three inches of rain. It was not too late to sow for the kharif, and the area cultivated, though considerably less than the area-sown in the previous year, was not much below the average kharif area. Indeed nearly 50,000 acres of new prairie-land were for the first time brought under the plough. During August very little rain fell, except a partial shower in Fázilká and its immediate neighbourhood, and for more than a month no rain fell in the greater part of the district, and in many places the crops dried up irretrievably. Had the rain held off a fortnight longer, very little grain would have been produced except in a few villages which had been fortunate enough to get partial showers; but in the second week of September abundant rain fell most opportunely over almost the whole of the district and saved the greater part of the kharif crop. The village ponds were filled and the grass revived, and the supply of fodder was assured. A considerable area had dried up so badly that this rain was too late to revive it and it produced no grain, and in a large area the outturn was poor, but most fields produced something, and the harvest was about three-fourths of the average. The floods of the Ghaggar came down at a good time for the rice sowings, and again in the beginning of September in good time to save it from drying up, and the rice-crop was unusually good, especially below the Annakai Chhamb; the villages dependent on the Dhanur Jhil however did not get good rice owing to the erosion of the Ghaggar channel which has lowered the level of that lake. Satlaj floods were low, and owing to this and the almost total failure of the early rains in the Hitar, very little land was there sown for the kharif. The areas cultivated and the reported quality of the produce were as follows :---

					AR	ba Sown bo	в Кнави	r 1882 (1	n Acres).			QUAI	ATY OF PROI	DUCB.	
LSSHSBMENT CI	RCLR.	Soil.		Jawar. (alone and with pulses.)	Bajra (alone and with pulses.)	Moth, Mung, Mash and Gwar.	Til.	Rice.	Miscella- neous.	Total.	Total of Assessment Circle.	None.	Very poor	Fair.	
Bagar		Barani	***	8,134	105,216	2,393	5		1	115,749	115,749	955	13,361	101,433	
		Chahi	***	69	3			12	36	120		14	19	87	
Tali		Rez	***	350	3,528	204	·	6,163		10,245		458	1,321	8,466	
		Barani	***	16,641	101,801	3,233	412	31	6	122,124	132,489	13,083	57,353	51,688	
ohi		Barani	***	87,581	307,038	13,812	14,668	***	15	426,114	426,114	136,023	223,542	66,549	
tar		Barani		5,642	28,457	1,810	695	***	4	36,608	36,608	3,880	8,605	24,123	
		Chahi	***	2,000					98	2,098		279	886	933	
litar 🚉		Rez		200						200		200			
		Barani		155	871	148	311	•••	443	1,928	4,226	276	682	970	
Total of the	district			120,772	546,914	24,600	16,091	6,206	603	715,186	715,186	155,168	805,769	254,249	

Thus, according to the patwaris' returns as checked by the supervising staff, of the 715,186 acres sown for the kharif only 254,249 acres produced a fair crop, and 155,168 acres produced no grain at all. In order to estimate the average outturn of the harvest the tashildars selected representative fields and had their produce weighed and also consulted the Zaildars. The average outturn and the resulting estimate of gross produce are approximately as follows:—

Assessment Circle.	OUTTURN	TED AVES (IN MAUN CRE.)		ESTIMATED	Тотаь.		
	Soil.	Jawár, &c	Bájra, &c.	Jawár, &c	Bájra &c.	Miscella- neous.	
Bágar	Báráni	1	2	8,000	210,000	•••	218,000
Náli	Rez Báráni	2	11/2	30,000	142,000	100,000	272,000
Rohi Utár	Bárani Barani	13/4	11 2	130,000 16,000	330,000 55,000	16,000 2,000	476,000 73,000
Hitar	{Chahi Barani	4	~i}	7,000	1,000	8,000	18,000
Total of the district			-	191,000	738,000	126,000	1,055,000

The very low prices of June 1882 gave way to the failure of rain, but did not rise very rapidly as the stocks of grain in the district were large and there was no great demand for export; and in the beginning of September barley, grain, jawar and moth were selling at 35 sers per rupee in Sirsá market. The favourable rainfall and the prospect of fair kharíf and rabi harvests sent prices down again, and on 1st January 1883, the lowest prices quoted at the large villages throughout the district were 70 sers per rupee for jawar and 50 for bajra. The average outturn of straw may be taken as six maunds per acre for jawar, and three maunds for the other crops, and the average selling prices of straw at ten maunds per rupee for jawar, 20 for bajra, and five for moth and pala. These estimates give the value of the kharif produce of 1882 approximately as follows:—

	A STATE OF THE PARTY OF THE PAR	VEST P		VALUE OF	VALUE OF	
Assessment Circle	Jawár,	BELLEVICE BELLEVICE OF THE STATE OF THE STAT		GRAIN PRODUCE.	STRAW.	Rs. 2,50,000 3,40,000 5,25,000 1,00,000 20,000
				Rs.	Rs.	EEP.5.10.55925.571674785708857259824880038
Bágar	50	40	50	2,20,000	* = 30,000	E. S. N. S. P. (1977) MIDELLING THE RESIDENCE OF THE PROPERTY
Nálí Rohí	70	40 50	60	3,00,000 3,75,000	1,50,000	16-2 (A) 图 (安治) [2] 图 (B)
Utac	60	45	50	70,000	30,000	COUNTY STATES OF THE RESIDENCE OF THE PARTY
Hitar	60	45	50	15,000	5,000	20,000
Total of the district				9,80,000	2,55,000	12,35,000

This estimate makes the value of the produce of kharif 1882 Rs. 12,35, 000 or about $4\frac{1}{2}$ times the new assessment of the whole year. Notwithstanding the partial failure of the kharif in some parts of the district, the previous harvests had been so good that the enhanced land revenue was realised with ease, and on 31st March 1883 the arrears amounted only to Rs. 159.

The general and abundant rain of September 1882 was very opportune for the rabi sowings, and a much Rabi Harvest, 1883. larger area was sown for the rabi than had' ever been before. The rain in January came in good time to save the crop from drying up, and the result was a bumper rabi crop throughout the whole of the Dry Tracts. The Ghaggar floods were not very favourable, and a smaller area than usual was sown with wheat; the gram crop in the low-lying parts of the channel was drowned by a flood in March. The Satlaj floods were low and the inundation canals had not been put in order, and the area flooded in the Hitar was small; but the abundant rain in September allowed of the sowing of a larger area of unirrigated land than usual in the Hitar also. The area sown on wells was about up to the average and the crop good. Throughout the district little of the land sown failed to produce a crop and on a large proportion of the area the outturn, both of grain and straw, was unusually good. The areas cultivated were as follows:-

		ARI	BA SOWN I	FOR RABI,	1883.	QUAL	ITY OF PE	ODUCE.
Assessment Circle.	Soil.	Wheat.	Barley, Gram & Oilseeds.	Total.	Total of Assess- ment Circle.	None.	Very poor.	Fair.
Bágar	Báráni Cháhi	09	4,676 273	4,679	4,679	988	1,112	2,579 283
N411 }	Rez Sotar Báráni	7,571	6,610 1,454 13.185	14.181 2,660 13,408	30,545	1,594 583 849	2,337 625 1.238	10,250 1,452 11,321
Rohí	Báráni	15,970	241,745	257.715	257,715	3,150	38,831	215,734
Utár	Báráni Cháhi	4 000	15.550 779	19,100	19,100	C200	100	14,559 4,479
Hitar {	Rez Báráni	3,000	753 1,146	3,753 2,675	11,107	400 321	100 277	3,253 1,977
Total of the	district	. 36,975	286,171	323,146	323,146	8,145	49,114	265,887

According to the results of observations made by the tahsildars on representative fields and enquiries made from the Zaildars, the average outturn and the gross produce may be estimated as follows:—

Assessment Circle.	Soil	OUT	D AVERAGE TURN, per acre.)	Hier Today	IMATE OF GROSS OUTTUEN. In Maunds.)			
		Wheat,	Barley, &c.	Wheat.	Barley, &c.	Total.		
	# 1 (20) (1 (42) (42)							
Bágar	Báráni Cháhi	61	3 71)		12,000	12,000		
náli }	Rez Báráni	4 41	6 5	30,000	120,000	150,000		
Rohi	Bárani	5	7	80,000	1,600,000	1,680,000		
Utár	Barani Chahi	5 8	7 8)	17,000	100,000	117,000		
Hitár {	Rez Barani	6 5	3	50,000	20,000	70,000		
Total of the distric	t			177 000	1,852,000	2,029,000		

There was a considerable export trade, and a large quantity of grain was purchased and stored in Sirsá, but there was no unusual demand in any direction, and a large quantity of grain remained in the hands of the peasants, whose stocks were fully replenished by two such good rabí harvests. Prices continued very low, and in Malaut in June 1883 wheat was said to be selling at 30 sers per rupee, and barley at 70 sers. The produce of straw was estimated as usual as about equal in weight to that of grain, and the average telling price of straw may be taken at ten maunds per rupee for wheat straw, and six maunds per rupee for barley straw. The resulting values are as follows:—

Assessment Circle.	HARVEST (Sers per	PRICES,	VALUE OF GRAIN	VALUE OF STRAW.	TOTAL VALUE.
	Wheat.	Barley, &c.	PRODUCE.		
Bagar Nali Rohi Utar Hitar	25 30 28 28 28	50 50 80 55 55	Rs 10,000 1,50,000 12,00,000 1,00,000 1,00,000	Rs. 2,000 23,000 2,75,000 18,000 10,000	Rs 12,000 1,73,000 14,75,000 1,18,000 1,10,000
Total of the district			15,60,000	8,28,000	18 88,000

According to this estimate, the value of the gross produce of the cultivated land in rabí 1883 was Rs. 18,88,000, or nearly seven times the total assessment of the year. The rabí instalment was realised with ease, and at the end of September 1883 the arrears of land revenue, including suspensions, were only Rs. 243. In July 1883 the district was in an unusually prosperous condition. The cattle had not suffered from scarcity of fodder for two years, and there was plenty of water and grass everywhere, with considerable stores of fodder from the past good harvests. The peasants had had two good rabí harvests,

and their private stocks of grain had been replenished; there were no arrears of revenue against them, and most of them had considerable savings in hand. In short the district, as a whole, was in a more prosperous condition than probably it ever was in before, and the accumulated capital was larger than at any previous period in its history.

Summary of observations of seven harvests.

Statement, a rough estimate of the kharif harvest of 1879, which was not actually observed, being added to complete the four years. The statistics are given in even thousands, the last three figures being omitted:—

AGRICUL-		sown. In acre		ARRA WHICH PRO- DUCED A CROP. (In acres.)			Total PRODUCE OF			VALUE OF GRAIN AND STRAW.		
YBAR.	Kharif.	Rabi.	Total.	Kharif.	Rabi.	Total.	Khafif.	Rabi.	Total.	Kharif.	Rabi.	Total,
1879-80 1880-81 1881-82 1882-83	7,00 7,27 7,73 7,15	2.76 1.53 2.50 3,23	9,76 8,80 10,23 10,38	5,50 3,65 7,50 5,60	2,65 1,50 2,40 3,15	8,15 5,15 9,90 8,75	6,00 4,38 24,51 10,55	13,30 5,13 17,24 20,29	19,30 9,51 41,75 30,84	Rs. 10,00 10,12 27,81 12,35	Rs. 16.85 9.84 18.54 18,88	Rs. 26.85 19.96 46,35 31,23
Average	7,29	2,50	9,79	5,56	2,43	7,99	11,36	13,99	25,35	15,07	16,03	31,10

Thus, according to the average of the last four years, 9\frac{3}{4} lakhs of acres were sown, but only eight lakhs produced a crop of grain; the gross annual produce of grain was 25 lakhs of maunds, or rather more than three maunds per acre; and the value of the grain and straw was 31 lakhs of rupees, or about Rs. 4 per acre producing a crop, a sum equivalent to eleven times the total new assessment. And although the area sown for the rabi has been little more than a fourth of the whole, the grain produce of the rabi, and the money value of its crops, have exceeded those of the kharif. The total grain produce of the year has varied from 9\frac{1}{2} lakhs of maunds to 42 lakhs, or more than four times as much.

the harvest prices vary so much as they do in Estimate of area sown the Sirsá district, a true average could only be and area producing a crop obtained by comparing the statistics of more than four years, especially where the crops observed have been so exceptional as in the past four years. The last two rabi crops have been exceptionally good in the Dry Tracts, and three of the four kharif harvests have been poor, yet there has been no such very poor year as in 1877-78 or in any of the years of real scarcity, and some allowance must be made for them. The floods of

the Ghaggar have not been favourable, and the area irrigated by the Satlaj floods has been much less than it used to be and than it might fairly be estimated at. I propose then to correct the estimate of the average by making allowance for these considerations, and to state my estimates of what may be considered an average harvest in the Sirsá district as it was in 1882, without allowing for further extensions of cultivation or irrigation or development of new canals not already in existence. They are estimates only, founded upon the observations of the past four years and the available statistics. It does not seem necessary to give in detail in every case the grounds on which the estimate is framed, especially as the detailed calculations have in many cases been already reported in submitting proposals for assessment and reports on the different harvests. The best way of checking and correcting the estimates will be by comparing them with the actual results of future harvests.

The area actually sown on an average of years and the area which

produces a grain crop I estimate as follows (in acres) :-

				m - m m - m - m - m - m - m - m - m - m	ED AVERAG			D AVERAGI	
Assessment CIRCLE.	Soil.		Total Cultirated area returned at Settlement.	Total,	Khárif.	Rabi.	Total.	Kharif.	Rabi.
Bagar Nali {	Barani Chahi Rez Barani		125.607 893 39.915 145,0°6	25,000 127,000	120,000 10,000 115.000	4,000 15,000 12,000	103,000 22,000 101,000	9,000 91,000	3,000 13,000 10,000
	Total	•••	185,813	152,000	125,000	27,000	123,000	100,000	23,000
Rohi Utar {	Barani Báráni Chahi Rez Báráni	:::::::::::::::::::::::::::::::::::::::	651,348 53,818 9,389 13,868 1,890	600,000 45,000 6,500 9,000 3,500	425,0 0 35,000 2,000 1,000 2,000	175,000 10,000 * 4,500 8,009 1,500	480,000 38,000 6,0 0 7,5 0 2,500	320,000 30,000 2,000 500 1,500	160,000 8,000 4,0 0 7,0 0 1,000
	Total	***	25,147	19,000	5,000	14,000	16,000	4,000	12,000
Total of the district		•••	1,041,733	910,000	710,000	230,000	760,003	554,000	206,000

That is, of the total area under the plough, which may be stated roughly as 10½ lakhs of acres, on an average only 940,000 are cultivated, and of this only 760,000 acres produce a grain crop, the remaining 180,000 acres sown being estimated as producing practically no grain at all.

The 760,000 estimated as producing a crop on an average of years may be taken as sown with the following crops:—It will be seen that about a lakh of acres are estimated as producing jawar, 4 lakhs bajra, half a lakh autumn pulses and til, and nearly two lakhs barley, gram and oilseeds, while wheat is produced in only 26,000 acres and rice in 5,000.

						1	KHARIF	CROP.			1	ABI CROP	
Аванавивит Ство	CLB,	Soil,	*	Total area producing a Crop	Total Kharif,	Jawar (slone - nd with pulses)	Bajra (alone and with pulses.)	Moth Mung, Mash Gwar	Til.	Rice,	Total Rabi	Wheat	Barley, Gram, Oilseeds, &c.
Bágár		Bárani		103,000	100,000	6,000	92,000	2,000			3,000		3,000
	1	Rez		22,000	9,000	1,000	2,000	1,000		5,000	13,000	9,000	4,000
iáli	}	Báráni		101,000	91,000	12,000	75,000	3,500	500		10,000	1,000	9.000
		Total		123,000	100,006	13.000	77.000	4,500	500	5,000	23,000	10,000	13.000
lohi		Báráni		480,000	320,000	70,000	2,10,000	30,000	10,000		160,000	5,000	155,000
iár		Báráni		38,000	30,000	4,000	23,000	2,000	1,000		8,000	1,000	7,000
	1	Cháhi		6,000	2,000	2,000					4,000	3,500	500
		Rez		7,500	500	2	500				7,000	6,000	1,000
litár	"]	Báráni	1	2,500	1,500	500	1,000	m	°		1,000	500	500
	,	Total	# P	16,000	4,000	2,500	1,500				12,000	10,000	2,000
Total of district				760,000	554,000	95,500	403,500	38,500	11,500	5,000	206,000	26,000	180,000

Estimate of average outturn and average gross produce of grain and straw.

181. The average outturn of grain on the area producing a crop I estimate as follows (in maunds per acre.)

Assessment Circle.	Soil.	Jawar and pulses	Bajra and pulses.	Moth, Mung. Mash, Gwar.	Til.	Rice.	Wheat.	Barley, Gram, Oil- seed, &c.
Bagar	Barani	11	11	11				2 6
Nali §	Rez Barani	3	3 2	2	2	12	6	6
Rohi	Barani	21	3	2	2		4	5
Utar	Barani	21	2	2	2	***	3	4
(Chahi	4	***				8	9
Hitar }	Rez		2			***	6	5
1	Barani	2	2				5	4

These estimates of average outturn applied to the areas producing a crop give the following as the estimate of average gross produce of grain (in thousands of maunds):—

Assessment Circle,	Jawar and pulses.	Bajra and pulses.	Moth, Mung, Mash and Gwar.	Tal.	Rice.	Total Kharif Produce.	Wheat,	Barley, Gram and Oiiseed.	Total Rabi Pro- duce.	Total produce of the year.
Bagar	9	138	3			150		6	6	150
Nali Rohi	40 175	156 630	60	20	60	268 885	59	775	119 795	387 1,680
Utar	10	46	4	2		62	3	28	31	93
Hitar	9	3				12	67	11	78	90
Total of the District	243	973	78	23	60	1,377	1.49	880	1,029	24,06

According to this estimate the average gross grain produce of the district is 24 lakhs of maunds. The average of the last four years only would have given over 25 lakhs of maunds, but instead of 11 lakhs in the kharif and 14 in the rabi as the four years' average gives, I estimate that on the average of a longer series of years 14 lakhs are produced in the kharif and 10 in the rabi. Taking all crops together the average outturn per acre is 2½ maunds in the kharif and 5 maunds in the rabi, or 3½ maunds per acre for both harvests taken together.

The average outturn of straw available as fedder for cattle may be estimated as follows in maunds per acre:

Assessmen	at Circle.	Soil.		Jawár.	Bájra.	Moth,	Rice.	Wheat.	Barley, Gram, &c.
Bágar		Báráni		2	2 4	2 3	8	6	9 6
Náli	}	Rez Báráni	***	10	3	2		5	4
Rohi		Báráni	***	6	3	3		4	5
Utár		Báráni Cháhi	***	15		2		8	9
*Hitár	}	Rés Barani	•••	10	3	2		6 5	9 5 4

This gives only the weight of that part of the straw which cattle will eat, apart from the hard woody stalk which is hardly eatable. Our observations tend to corroborate the assertions of the peasants that in the rabi harvest the weight of straw is approximately the same as that of the grain. Ordinarily a much larger area produces straw than the area producing grain, for the rainfall is often sufficient to develope the stalk and leaves to some extent although not sufficient to develope the ear and grain. Allowing for this and for the leaves of the jhárberi (pála) which are a valuable fodder and in many fields, especially in the south-east of the district, average a maund per acre, the following estimate of the average amount of fodder produced by the cultivated land may be framed:—

		AVERAG	E Proi	DUCE O	F FODD	ER (IN TO	OUSAN	Ds of	MAUND	•)
As&sement Circle.	Jawar, &c	Bajra, &c.	Moth.	Rice.	Pala.	Total Kharif.	Wheat.	Barley, &c	Total Rabi.	Total of year.
Bagar Nali R-hi Utar Hitar	0.0	184 233 630 69 4	4 10 90 4	40 	50 50 5	250 373 1,190 98 37	59 20 3 66	6 60 775 28 12	6 119 795 31 78	256 492 1,985 129 116
Total of District	555	1,120	108	40	125	1,918	148	881	1,029	2,977

According to this estimate the cultivated land produces 20 lakhs of maunds of fodder in the kharif and 10 lakhs of maunds in the rabi.

Account of the different take up different constituents of the soil, and it is found that the produce of the two when sown together is often somewhat larger than if they were sown separately in the same area, and partly in order that the cultivator may have something of each kind of produce or that, should the season prove unfavourable to one kind, there may be a chance of getting something of the other kinds of produce. Sometimes they are reaped separately, as in the case of javoar and moth when sown together, but sometimes the produce re-

mains mixed, as in the case of barley and gram, which, once sown together, are rarely afterwards separated. This makes it difficult to frame an estimate of the produce of each kind of grain separately.

Rice (dhán) is always sown by itself in the kunds specially prepared for its cultivation, which have been already described. Mr. Oliver is said to have distributed some rice seed got from the Peshawar valley, but it was not very successful, and only a very few fields are now sown with it; it is recognised as a finer and more delicate rice than the ordinary variety, but the produce is not so great, and it is not found remunerative. Several varieties are grown in small areas as luxuries, such as the ram-gawayan, sunkharcha, and bansmatti varieties, but practically the only two kinds of rice cultivated here are the munji and the kharsu varieties. The latter, which is also called seora, is much coarser and less valuable than the munji and of a darker reddish colour but can be sown later, and is ordinarily only sown when the kunds have been unfit for rice-sowing until September, when it is too late to sow munit, but not too late for kharsu. Ordinarily very little kharsu is grown (in 1882 only 117 acres), and it may be said that practically the only variety sown is the munji, which is sown broadcast, about 25 sers or 30 sers of seed going to the acre. The mode of cultivation of rice has already been described. The sowing commences as soon as possible after the end of June and continues until September; and the reaping commences in the end of November and continues until the end of December. The ground is ordinarily quite dry by that time, and the rice is bound in sheaves and stacked to be threshed in the field by bullocks. The unhusked rice (dhán) is sometimes sold just as it comes from the threshing-floor, but before using it has to be husked in a mortar, and many of the grains are broken in the process, which reduces greatly the value of the cleaned rice (chanval) unless it is sifted and only the full grains sold. Husking (kutái) costs about four annas a maund. Of the 8,927 acres of land set aside for rice-cultivation about 6,000 acres are sown on the average, and of this 5,000 acres are estimated as producing a crop averaging 12 maunds per acre. When rice dries up, no grain is formed in the ear, and it is then called marayan. Such dried up rice is good fodder, but the ordinary rice-straw (purali) is not of much use as fodder.

Jawar or great millet is sown about July and reaped about November or December, and is said to require at least 90 days to ripen. It is not often sown for fodder only (chari), even on the wells of the Hitar. In that tract it is usually sown alone broadcast, about 10 sers to the acre, the land being watered, ploughed and levelled first. If kept well irrigated it produces a good crop, but if irrigation and rains fail little grain is produced, especially if the white maggot (gindár) attacks it. The average area sown on the wells of the Hitar is about 2,000 acres, and the average produce may be taken as 4 maunds per acre. On the sotar lands of the Ghaggar valley also jawar is sown unmixed with other crops; the land is rich, and if the rainfall is good it sometimes produces excellent crops as in 1881, but more often the crop is very poor for want of sufficient moisture. In the Dry Tracts jawar is

sown in good soil, sometimes alone, but more often mixed with moth, mung, gwar and other seeds in varying proportions, and it is difficult to estimate how much of each kind of grain is produced. In these Dry Tracts the amount of jawar seed is about 8 or 10 sers per acre, and the average outturn of the mixed crop is estimated at about 1½ maunds per acre in the Bagar and 2½ maunds elsewhere. In 1881, however, some exceptionally favoured fields in the Rohi produced more than 12 maunds of jawar per acre. Jawar grain is not thought so nutritious as bajra, and always sells much cheaper, but the straw makes good fodder and in times of scarcity commands a good price.

Bájra is the staple kharif crop of the Dry Tracts, and is sown in about 5 lakhs of acres annually, but on the average only some 4 lakhs of acres produce a crop. It grows on the lightest soil and is almost always sown mixed with moth, mung, gwar and other crops in different proportions. Whenever favourable rain falls, at any time from April to August, the mixed seed, about 2 to 4 sers per acre, is drilled into the ground often without any preliminary ploughing. The crop often produces no grain, only stunted straw (bukwána), but if it comes to anything it is reaped in October and November, the ears being generally cut off first and thrown in a heap on the threshing-floor to be trodden out by bullocks, while the stalks are left to be cut and stacked at leisure. Sometimes the ears do not all come to maturity at once, as the stalks go on earing until the frost kills them. Bajra straw is very poor fodder and used to be left to rot on the ground, but recent fodder famines have taught the people its value, and it is now generally carefully cut and stacked against a drought, and stacks of several years back may sometimes be seen standing about the fields. Bajra grain is the staple food of the inhabitants of the Dry Tracts for half the year, and of most of the Bágri population all the year round; it is considered good and nourishing and generally commands a high price as compared with jawar, barley and gram. The outturn varies very much, but may be estimated as averaging 11 maunds per acre in the Bagar and 3 maunds in the Roli.

Moth is sometimes sown alone, but much more commonly with bájra or jawár. It is almost always grown on light upland soil dependent on the local rainfall, and when sown alone takes from two to four sers per acre of seed and produces on the average about 2 maunds per acre. The area returned as under moth alone was 24,000 acres in 1880 and 37,000 acres in 1881, but a large proportion of the area returned as under bájra and pulses also produces moth. The whole plant is collected, separately from the bájra, and its leaves are preserved as carefully as the grain, for they form a much-prized fodder which always commands a good price, as it is very nourishing and greedily eaten by cattle. The grain is sometimes given to cattle, but more commonly ground up

into flour with bajra, or split and eaten as pulse (dál).

Múng is not nearly so much grown as moth, and the area under múng alone was only 424 acres in 1880 and 3,306 acres in 1881, but like moth, it is often sown with bájra and jawár and its leaves and pulse are similarly used. Másh, another pulse similar to moth and múng, is still less common and is grown chiefly on the flooded lands

of the Hitar; but only 242 acres were returned as under mash in 1880 and 101 acres in 1882.

Gwar is more common, and though sometimes sown with bajra and other crops on the dry uplands, is often sown alone from 3 to 10 sers to the acre, and produces about two maunds per acre. The areas under gwar alone in the last three years have been 10, 29 and 17 thousand acres respectively. It is sown and reaped at the same seasons as the other kharif crops, and as the grain is coarse

and inferior it is ordinarily given to cattle as well as the fodder.

Til is a crop of some importance in the Dry Tracts, where it is ordinarily sown in July and reaped in November. It is sometimes sown with other crops, but more often sown alone broadcast, about three ser to the acre, and the average produce is about two maunds per acre. The seed is pressed in the oil-mill (kolhú), by a Telí or oil-maker, and the resulting sweet oil (míthá tel) is used with food instead of ghí, or for anointing the body, and more rarely for burning in lamps. The number of oil-mills in the district is returned as 229.

Water-melons (matira) are grown largely, where one would hardly expect to find them, in the fields of the dry uplands. A little seed is mixed with the bajra sown, and in September the creeping stems of the plant produce melons larger than a man's head containing a large quantity of pure cool juice, which is much prized in that dry country where the drinking water is often so hot and filthy. These melons are very common in the season all over the Dry Tract, but they do not keep long unless very carefully packed so as not to touch each other and covered up from the action of the air. They are sometimes cut up, dried in the sun and preserved for use in the cold weather. Another kind of small melon grown in the uplands is the kakri, which is similarly preserved; so is a sort of cucumber called kachri. A few sweet melons (kharbúza) are grown in the low lands near Sirsá flooded by the Ghaggar.

Cotton is grown only on a few wells on the Satlaj, and the area sown is insignificant, being only 148 acres in 1880 and 257 in 1881. About 300 acres of maize and less than a hundred acres each of sngarcane (paunda), red pepper (mirch) and hemp (san) and a few acres of china are also grown in the Satlaj Hitár. Sugarcane is said to have been introduced only a few years ago. No sugar is extracted from it; the cane (paunda) is sold in the Fázilká market at a paisa per stalk and cut into pieces which are chewed and sucked raw. The produce of a kanál of land sells for about Rs. 6 or Rs. 8,

say Rs. 60 per acre.

Account of the different rabi crops.

Account of the different rabic crops.

Account of the differ

maunds per acre on the land irrigated from wells and 6 maunds per acre on the flooded lands of both rivers. (In England the average outturn of wheat is 25 bushels or about 18 maunds per acre.) In favourable years the hard clay of the Sotar where thoroughly moistened is sown with wheat and sometimes produces excellent crops, as in 1882; and on the Satlaj, when the floods fail, patches of lowlying land where the rainwater has collected are often sown with wheat. In both cases the average outturn may be estimated at 6 maunds per acre. The estimated average area under wheat in the Dry Tracts is only 7,000 acres, but its cultivation seems to be spreading. It requires more careful cultivation than barley, and the seed is more expensive, but the produce is much more valuable, though the average outturn is rather less, being about four maunds per acre. Wheat is much preferred for food to any other grain except rice; it is the ordinary food of the Musalmans of the Satlaj Hitar, but is a luxury to the inhabitants of the rest of the district. The straw (túri) of wheat is used for fodder, but is not nearly so nutritions as that of barley and commands a lower price. For wheat cultivation the ground is prepared during the rains by ploughing and pulverising it twice or thrice, and the seed is sown in October or November, about 20 sers to the acre on unirrigated land, 30 sers on flooded land, and 40 sers on land irrigated from wells. (In England the ordinary quantity of seed is 21 bushels or about 65 sers per acre.) On flooded and unirrigated lands the crop is left to itself and not even weeded, but on wells, especially in dry seasons, it is irrigated as often as the capacity of the well allows. Some of the wheat-crops on wells and even on flooded lands in the Hitar are excellent, almost as good as can be seen anywhere in India, but these are the exceptions, and often the straw is stunted and the outturn very poor. The grain produced in the Hitar is often large and of excellent quality, but that of the uplands is generally small, dry and shrunken. The crop is reaped in April and threshed by means of bullocks treading it out on the bare ground. Considerable care is taken to clean the wheat and to keep the seed free from other grain, especially from barley. Sometimes wheat is sown mixed with barley and gram (gojiyá or berra): if mixed with barley alone it is called jawali wheat, and if there is no barley in the mixture it is called nijawi; but the area sown with these mixtures is small (only 500 acres being so returned in 1881 and 1,700 acres in 1882) and wheat is ordinarily sown alone. In the Hitar three varieties of wheat are grown. (1) The ordinary red wheat (ratti or lál kanak), also called kálchingárí because the beard (chingár) gets of a dark colour when ripe. This is the wheat ordinarily grown in the uplands also. Its ears (sitta) are about three inches long and the grains run in two lines, which make the ear look two-sided; the grain itself is short and thick. (2) Pamman, with a dark-green beard and a general dark colour in the sheaf, has distinctly two-sided and heavy ears, from 21 to 3 inches long, and a long thick and soft grain. (3) Daudí, which again is divided into three subvarieties (a)

goni or beardless (literally "hornless") with short light-coloured ears from two to three inches long but sometimes (changhundi) having as many as four lines of flowers (ghundi); its grain (dána) is white, thick and soft and makes very white bread (b) nárí dáudí or bearded (literally "horned") having whitish ears from 3 to 4 inches long, and a white beard; its grain also is white, thick and soft (c) paháran,so called because its seed (bih=bij) was brought from the hills (pahár) some ten years, ago by the Banya who owns part of Rána—with long pointed beardless ears from 4 to 5 inches long, and a reddish appearance; the grain is thin and reddish, and although the ear is longer than that of gon't daud't it is thinner and has only three lines of flowers (trighund) and so produces less than the indigenous gont. Except the paháran variety these are all said to be old wheats well-known in this part of the country; and it says much for the care with which the seed is grown that they should be so well distinguished. They seem to be sown on all kinds of irrigated soil, but pamman and ratti kanak are said to require more moisture (tarawat) than goni Goní dáúdi is considered the best variety because of its whiteness, and if ordinary red wheat is selling at Rs. 2 per maund, paháran will fetch about Rs. 2-1 and goní, nári and pamman about Rs. 2-4 per maund. (These prices may be compared with the average price of wheat in England, say 40s. a quarter, or about Rs. 4 per maund.) Sometimes a grain or two of oats finds its way into a wheat field, but the people do not know it as a separate crop, and say that it is wheat which has deteriorated or become diseased (dhanak).

Barley (jan) and gram (chholá or channa) are the staple rabí crops in the uplands. They are often sown separately but very often together (bejhar), when they are reaped, threshed and sometimes even made into flour together. The areas sown were returned as follows for the last four years:—

Year	Barley alone.	Gram alone.	Barley and gram
	(acres)	(acres)	(acres)
1880	1,12,000	41,000	88,000
1881	55,000	33,000	40,000
1882	78,000	27,000	1,10,000
1883	88,000	40,000	1,45,000

Barley is not sown much on irrigated land, and when sown separately it is generally grown on the light sandy soil of the uplands. It is sown in October, about 20 sers to the acre. Sometimes if the early seed-time has not been favourable, it is sown as late as December, but such late crops (kanadji) seldom produce much grain. The average outturn of barley on the uplands is about 5 maunds per acre. It is reaped in March or April and trodden out by bullocks. The straw (turi)

is considered good fodder and commands a fair price. The husks are separated from the grain by pounding it in a mortar, and the flour of barley and gram is made into scones or porridge, which form the ordinary food of great part of the population of the

uplands for half the year.

When gram is sown alone on the unirrigated uplands it is ordinarily sown on the stronger soils about 20 sers to the acre. It is sown in the early part of October and reaped in April, the average produce being from 4 to 6 maunds per acre. A good deal of gram is sown unmixed with other seed on the low lands irrigated by the Ghaggar, about 25 sers to the acre, and the average ontturn is about 6 maunds per acre; but the crop is specially liable to be drowned by the winter floods of the Ghaggar. The young leaves of gram are plucked and used as greens (ság) with scones and porridge, and the flour of gram mixed with that of barley is much used as food. The dry leaves are used as fodder but are not

very nutritious.

Sarhon or sarson is rarely sown by itself, and the area sown with sarson alone was less than two thousand acres in 1881 and less than a thousand in 1882. When sown alone in the uplands it is sown in October and reaped in April, the seed is from 3 to 6 sers per acre and the average outturn about 3 maunds per acre. But it is very common in the uplands to sow sarhon in lines in the barley and gram fields, and a large quantity of sarhon is grown in this way; it is reaped and threshed separately from the other crops. The young leaves of sarhon are largely used as greens (say) and from its seed bitter oil (karwa tel) is expressed in the oil-press (kolhú) a maund of seed being pressed in a day and giving about 12 sers of oil, which is used for burning, &c., and more rarely consumed with food. The refuse or oil-cake (khal) is given to cattle. A considerable quantity of sarson seed is exported, chiefly towards Bombay. The similar plant, tara or tára míra, is grown in the same way, but is not unite so largely grown as sarson, because although less liable to suffer from frost, it is coarser and its products are less palatable.

Churál, masar and methra are low plants of the pea-kind which are sown in the Satlaj Hitár, chiefly on new land recently thrown up by the river, but the area sown is seldom so much as a thousand acres. They are generally sown broadcast with one ploughing, and the crop is very precarious. The produce is generally used for fodder, but the peas are sometimes used for human food. The people say that a constant diet of churál is apt to cause a sort of paralysis, and point to several invalids as instances of its evil effects. There are several other plants of the pea-kind which grow half-wild on the cultivated

lands of the Hitar and are used as fodder.

Turnips (ghonglu), both white and red, carrots (gájar), radishes (múlí) and other vegetables are grown chiefly on lands irrigated from wells or by the floods of the Ghaggar and Satlaj, but the areas returned as under these crops in 1881 and 1882 were only 734 and 1,467 acres respectively. They are grown partly for human food

and partly for cattle. Tobacco is grown on well-lands in the rabi, but the areas sown in 1881 and 1882 respectively were on the Ghaggar only 149 and 191 acres, and on the Satlaj 93 and 60 acres only: the flower-buds (kaliyán) of tobacco are broken off in order to make the leaves ripen with a bitter (karwa) flavour, and the leaves are buried in the ground for some time with the same object. A few acres of other fancy crops, such as dhaniya, haliya, and kasumbh are also sown on irrigated lands.

Estimated average gross crops together there is no means of making an produce of each crop. exact estimate of the proportion borne to each other by the various grains whose seed and produce are thus mixed. It is possible however to make a rough estimate approximating to the truth, and my estimate of the average gross annual produce of the district is as follows (in thousands of maunds):—

Кнав	F PRODUCE.	RABI PE	CODUCE.
Crop.	Average Gross Produce.	Crop.	Average Gross Produce.
Jawár Bájra Moth Múog	2,00 8,00 1,80 47	Wheat Barley Gram Sarson	1,49 5.00 3,00 40
∃wár Fil	60	Tara Churál, Masar Methra	20
	60	Vegetables	10,29

According to this estimate, of the 24 lakhs of maunds annually produced in the district, about 22 lakhs are grain ordinarily used for human food, one lakh is oilseed (til, sarson and tara) and one lakh vegetables and grains sometimes used for human food, but ordinarily given to cattle.

According to the estimate furnished to the Famine Commissioners, a typical family of five persons, on an average for the whole Province, consumes 3.2 sers of grain per day. The Sirsá population is almost wholly agricultural and is unusually well-to-do, so that most of them have large appetites and the means to satisfy them, and the average consumption of a typical Sirsá family of five may be taken as 4 sers of grain (8lbs.) per day or say 36 maunds per annum, which gives for the whole population of 2,53,275 persons an average annual consumption of grain alone of about 18½ lákhs of maunds, besides vegetables, greens, berries, spices, oil, milk, butter and the butcher-meat

consumed by the Musalmans. This leaves available for export an average surplus produce of $3\frac{1}{2}$ lakhs of maunds of edible grain and nearly a lakh of maunds of oilseed. According to the Municipal returns the average imports of grain during the last eight years into Sirsá and Fázilka have been about seven lakhs of maunds annually. Allowing a lakh and a half of maunds for the consumption of their population of 19,143, there remain $5\frac{1}{2}$ lakhs to be re-exported out of the towns. Some of this is consumed in the district and some comes from Pattiála, Bikáner and other places, and there is nothing inconsistent between this estimate and the estimate of $3\frac{1}{2}$ lakhs of maunds of grain as the average surplus produce of the district.

185. In order to ascertain the average prices at which the harvest produce should be valued, enquiry was made in History of prices. three different ways. (1) The prices reported every fortnight by the Deputy Commissioner as those prevailing in the Sirsá market were extracted from the Panjáb Gazette and an average struck for each year since the Mutiny, and a separate statement was drawn up showing for each year the prices reported as prevailing about harvest-time. (2) The books of the chief traders of Sirsa and Fázilká towns were examined and the prices of wholesale transactions extracted, and an average of these taken for each year since the Mutiny as representing the average trade prices of these two principal markets. (3) The books of the grain-dealers of all the principal towns and villages in the district, viz., Sirsá, Ránia, Ellenábád, Rori, Guda, Sohuwála Chautála, Dabwálí, Abulkharána, Malaut, Abohar and Fázilká, were examined and the harvest rates for each village were obtained by comparing a number of transactions about harvest-time between Banyas and peasants, and taking the average of the rates given by them. It is these last prices that concern us, for they show at what price the peasant could at harvest-time convert his grain into cash in the nearest market. The Gazette prices and the trade prices are chiefly important as corroborating the harvest prices obtained by the peasants, and as the general results of the very numerous independent enquiries on the whole corroborate each other, they may be considered quite trustworthy. order to obtain an average for each assessment circle the average of the prices prevailing at each harvest in the different villages in or near the circle was taken, and in order to get an idea of the general changes in prices, an average of the harvest-prices was calculated for five-year periods. This is of course a rough method of ascertaining average prices, for it leaves out of account the quantity of grain which changed hands at each particular price; but it is the only method practicable and gives a fair idea of the rise and fall of prices. We were not able to get much evidence as to the prices prevailing before 1850, and the prices of the earlier years were calculated on fewer transactions and are therefore less trustworthy than those of later date. The following statements give the average prices so calculated for quinquennial periods from 1850 to 1883 (the last period however comprises only four years' rabí prices and three years' kharíf prices.) The prices are everywhere stated in standard sers per rupee.

Chaks Nálí and Bágar.

(Average prices of Sirsá, Rániá and Ellenábád.)

	RABI I	PRODUC	E (ABO	UT 1sT	JUNE.)	KHARIF PRODUCE (ABOUT THE END OF DECEMBER.)								
Five-year period.	Wheat.	Barley.	Gram.	Barley and Grem.	Sarson.	Jawár.	Bájra.	Moth.	Máng.	Til.	Rice (un- husked.)	Gwår.		
1850 to 1854	4.1		49				80	* King			38			
1855 to 1859	45	111	75				54	105	***	4	39	120		
1860 to 1864	24		37				37	42		22	29	57		
1865 to 1869	20	33	• 32	27	17	44	26	33	27	16	22	46		
1870 to 1874	22	36	34	32	23	48	30	33	***	13	27	41		
875 to 1879	21	44	39	35	20	56	31	39	25	18	28	48		
880 to 1883	20	39	34	38	18	38	30	36	28	17	27	49		

Chaks Hitár and Utár.

(Average prices of Fázilká.)

1850 to 1854	44				40						
1855 to 1859	48			*	36	78	52			34	
1860 to 1864	29	24	51		23	33	35	34	19	16	 ***
1865 to 1869	23	38	33		20	26	22	20	16	13	
1870 to 1874	24	35	29		16	37	29	31	26	14	 55
1875 to 1879	24	40	36 .		17	30	30	35	18	11	 38
1880 to 1883	22	38	31	.38	16	38	31	33	25	13	 52

Chak Rohi. (Average of eight villages.)

1850 to 1854	36	68	48	79	48	73	65	72	86	30		
855 to 1859	39	114	93		51	102	71	107	93	33		
860 to 1864	24	71	48	63	30	67	50	55	37	19		53
865 to 1869	21	52	42	46	36	34	27	27	25	15		37
870 to 1874	22	43	33	41	24	43	32	35	28	14	1 2 NA	6
875 to-1879	21	45	40	41	21	43	33	34	24	13	***	4
880 to 1883	21	43	35	39	19	43	35	40	29	15	***	5

The available information may be summed up as follows. Upon the whole the prices of the different food-grains of both harvests have gone up or down together according to the nature of the harvest of the year and latterly with some reference to the demand for export. Barley has almost always been somewhat cheaper than gram, both being much cheaper than wheat. Similarly, jawar has always been somewhat cheaper than bajra, both being much cheaper than rice. Upon the whole the cheapest grains have been jawar and barley, sometimes the one being cheaper and sometimes the other; then gram, then bajra and lastly wheat. In years of scarcity the difference of price between the different grains becomes very small. In 1838, just after the famine of 1837, when grain was probably dearer than usual,

Captain Thoresby valued the ordinary produce at more than two maunds per rupee. In 1844-45 gram and barley sold at 44 maunds and bajra at 3 maunds per rupee. From 1850 to 1855 the cheaper grains sold at about 13 maunds per rupee; and during the next five years prices were very low, the average rate of barley and jawar in the Rohi being over 21 maunds per rupee, the cheapest grain sometimes selling at over 3 maunds. But the great era in the history of prices in the Sirsá district was the famine of 1860-61 o in that year, owing to the widespread scarcity, prices rose higher than they had ever been before and trade was greatly stimulated. The peasants and grain-dealers of the district learned how profitable it was to watch the market and carry grain from where it was cheap to where it was dear; and they never forgot the lesson. Prices never again fell so low as they had been before the famine, and they did not again fluctuate nearly so much from year to year and from village to village. The high prices of the famine year 1861 did not hold and prices were pretty low in 1862 and 1863, were high from 1864 to 1867, but suddenly rose still higher in the famine year 1868-69, when they were higher than they have ever been before or since. They then fell gradually until the beginning of 1877, when they were lower than they had been for eleven years and than they have since been. The failure of the rains of 1877 and the demand caused by the famine in the South then sent them suddenly up, and notwithstanding the good harvests of 1878-79, the demand on the frontier caused by the Afghan war and the scanty rainfall of 1880 kept them at a high figure until the good rains of 1881 brought them down, and owing to the good rabi harvests of 1882 and 1883 prices continued to fall and in 1882-83 were lower than they had been since 1877. The periods of low and high prices since 1861 have been (1) 1862 to 1867, six years of low prices; (2) 1868 to 1871, four years of high prices: (3) 1872 to 1876, five years of low prices; (4) 1877 to 1880, four years of high prices; and (5) 1881 to 1883, three years of low prices. Prices have however on the whole been much higher since 1860 than they were before, and the five-year periods, which contain both good and bad years, give approximately the same average prices since 1860. There is little reason to expect that the average prices-will fall lower than these, for communications have improved and are continuing to improve, and no doubt the opening of the Rewarf-Firozpur Railway will cause prices to rise greatly throughout the tract and will keep them steadily high. The peasants even of out-of-the-way villages keep themselves informed of the state of the market and bring their grain long distances to sell for eash at the market rates in Sirsá and Fázilká. The ordinary rate of carriage is one anna per maund per stage of 10 or 12 miles, and the differences in price in different villages are never for any length of time greater than are to be accounted for by the expense of carriage. When there is a demand towards Sind the peasants within reach take their grain to sell at Fázilká; and if the demand is towards Delhi, grain goes to Sirsa. Yet the distances are so great and the produce of the district so varying in quantity and so uncertain from harvest to harvest,

that the prices of grain still fluctuate greatly, though not so much as before 1860. Thus in 1868-69 prices were double what they were the year before, and in 1877-78 nearly treble those of the previous year; and in the Sirsá market the harvest price of barley has fluctuated between 66 and 12 sers per rupee, and that of bájra between 50 and 10. Where the fluctuations are so great averages are apt to be misleading; but if the average prices of the five years ending 1879 be compared with the average of the six years ending 1854 as given for the Sirsá market in the annual Revenue Report as follows:—

Average of years.	Wheat.	Barley.	Gram.	Jawár.	Bájra.
1848 to 1854 (sers per rupee) 1874 to 1879 (sers per rupee)	32 20	50 37	44 35	51 26	43 32
Rise per cent	60	35	26	42	34

and if the prices prevailing in the villages of the Rohi from 1850 to 1860, which may be taken as those of last Settlement, be compared with those of the years since 1870, it may safely be asserted that the prices at which the peasants of the district as a whole can dispose of their harvest produce are more than 50 per cent. higher now than they were at last Settlement.

Estimated average prices of grain and fodder.

Estimated average prices of grain and fodder.

Last few harvests which have been carefully observed, but without taking into account the Rewari-Firozpur Railway and future developments of communications, I estimate the average prices for the different assessment circles as follows (in sers per rupee):—

	R	BI PR	DUCE.		KHARIF PRODUCE.							
Assussment Circle.	Wheat.	Barley.	Gram.	Sarson.	Jamár.	Bájra.	Moth.	Mung.	71.	Rice unbusked.	Gwar.	
Bagar and Nali Rohi Utar and Hitar	24 25 25	48 55 50	42 50 45	• 24 24 20	45 50 40	35 40 35	45 50 40	30 30 25	20 18 15	80	80 60 60	

It is more difficult to estimate the average prices of fodder, for owing to its bulk the cost of carriage is greater than that of grain, and owing to its being less carefully preserved and to the precariousness of the annual growth of grass on which the need of artificial fodder greatly depends, the variations in its price are even greater than in

that of grain. For instance, in the Sirsá market in June 1881 barley straw was selling at 1½ maunds per rupee, and wheat straw at 1½ maunds, and in June 1882 their prices were 3 maunds per rupee for barley straw and 4 maunds for wheat straw, and in the villages the differences were even greater. Bájra straw sometimes fetches almost nothing in the market, but in February 1881 it was selling in the Sirsá market at 3 maunds per rupee. I estimate the average prices of straw as follows (in maunds per rupee.)

Assessment Cir		CIR-	KHARIP STRAW.						Rabi [®] Straw		
*	CLB	1100	Jawar.	Bajra.	Moth and Mung,	Rice.	Gwar.	Pala.	Wheat.	Barley,	Gram
Bagar Robi Utar	and	Nali Hitar	8 10 6	15 20 12	4 5 4	12	10 12 10	. 6 6 6	6 10 7	4 7 5	4 8 4

187. These estimates, combined with the estimate of gross produce, give the following as the estimate of the average value of gross produce of the cultivated land (in thousands of rupees):—

Assessment	VALUE OF KHARIF PRO- DUCE.			VALUE	VALUE OF RABI PRODUCE.			TOTAL OF YEAR.		
CIRCLE.	Grain.	Straw.	Total.	Grain.	Straw.	Total.	Grain.	Straw.	Total	
Bagar Nati Rohi Utar Hitar	70	23 35 1.00 11 5	1,83 3,35 9,20 81 20	5 1,60 1,50 30 1,20	2 25 1,20 6 12	7 1,85 7,70 36 1,32	1,65 4,60 14,70 1,00 1,85	25 60 220 17 17	1,90 5,20 16,90 1,17 1,69	
Total of district.	13,65	174	15,39	905	165	11,30	23,30	339	26,69	

The average of the observations of the last four years gives a gross value of 31 lákhs of rupees, of which 15 lákhs worth are produced in the kharíf and 16 lákhs in the rabí, but according to my estimate the average annual value is 263 lákhs of rupees, of which 153 lákhs worth are produced in the kharíf and 114 lákhs worth in the rabi. This estimate is certainly a low one, especially in the Robí and Utár circles, but according to it the average annual value of the produce of the cultivated land alone equals nine times the total new assessment of the district.

188. At last Settlement the area of land still uncultivated was re turned as 13,21,618 acres or 66 per cent. of the The uncultivated area. total area, and now according to the measurements of the present Settlement the uncultivated area is only 8,56,622 acres or 45 per cent. of the total area. Of this only 71,657 acres, or less than 4 per cent. of the total area, is returned as occupied by villages, roads or ponds, or otherwise incapable of cultivation. There are no hills or stony places in the district and the only classes of soil which are not fit for the plough are the hillocks of pure sand, which are of no great extent, the beds of pure river-sand on the Satlaj, and the kallar soil rendered barren by salts (reh or shor). Such soil is found in patches here and there all over the district and is easily distinguished by its being totally devoid of vegetation. Besides the patches so impregnated with salts as to produce nothing, there are many fields which owing to the presence of salts produce very little in places. Some soil also owes its barrenness to the presence of a bed of kankar too near the surface. The culturable area not yet brought under the plough is as follows:-

Assessment Circle.			Culturable waste (in acres.)		Per cent. on Total		tal
Bágar Nálí Rohí Utár Hitár			38,302 1,31,661 • 5,33,095 58,716 23,191			22 39 43 51 38	4
Tot	al of district		7,84,965		14 - 15 - 14 - 14 - 14 - 14 - 14 - 14 -	41	

Some of this land is set apart round the village-ponds to collect rainwater, and no one is allowed to cultivate such reserved portions; and in many villages in the Rohi, and still more in the Sotar valley where the soil is hard and requires to be well soaked before being ploughed, large areas of land are kept uncultivated, and the rain which falls on them is guided by shallow surface drains (súá or ágam), sometimes as much as half a mile long, on to the adjacent cultivated fields. Practically the whole of this area, say eight lákhs of acres, is available for cultivation and little inferior in productive capacity to much of the land already cultivated. Meanwhile it annually produces abundance of grasses of all kinds in the rains and affords food to great numbers of cattle.

The cattle of the district.

The cattle of the district.

The cattle of the district.

The cattle of the disdrove about from place to place for grass and
water; but as cultivation spread and the produce
of the cultivated land increased in value, they became less dependent
on the produce of their herds and accumulated other forms of capital.

There seem to be some signs that the rapid change from the pastoral to the agricultural mode of life which this district has witnessed, and the breaking up of the prairie, may have caused a diminution in the numbers and perhaps a deterioration in the quality of the cattle of the tract, but their food-supply must be less precarious now than it was formerly when they were entirely dependent on the grass produced annually in the rainy season, which was rarely cut and stored, and must, have died in immense numbers in seasons of drought. Yet even now, notwithstanding the care with which straw and even grass are preserved, and the high prices which they sometimes command, a serious drought deprives the cattle of their usual supply of food and the stocks soon becoming exhausted vast numbers of them die of simple starvation, especially when, in an emaciated condition, they are exposed to severe cold such as often follows rain in the cold weather. Mr. Oliver reported in 1863-64 and 1864-65 that the cattle were fast diminishing in numbers and deteriorating in value, a large number had died of murrain and starvation, herds driven towards Karnál in search of pasture had returned diminished by two-thirds of their number, and a great many cattle had been sold and taken down country. The breed had greatly deteriorated within Mr. Oliver's experience of 21 years, and none of those fine bullocks for which the country was once famous were to be met with. This deterioration he ascribed to three causes;-(1) the diminution of good pasture, as all the best of the land was brought under the plough; (2) the frequency of murrain; and (3) breedingin, due to the smaller herds and their more limited range. In 1866-67 the eattle were still suffering from want of pasture, and in 1867-68 it was reported that more than half the cattle in the district had perished within the preceding two years from scarcity of fodder and the severe frost of March 1868, and the survivors were so tottering and emaciated that they could hardly be driven out to graze. Cattle in such a condition were ill-prepared to meet the drought of 1868-69, and it was estimated that in that year of 2,02,327 horned cattle 1,48,590 died and little more than a fourth were left. The Bagris turned their cattle loose, and the Musalmans killed and ate theirs, but the Sikhs spared no trouble and expense to obtain fodder for their bullocks. So few were left that in the following March women were to be seen drawing the plough. In 1874-75 cattle-disease was prevalent and was ascribed to the drought; of 13,000 cattle affected 7,000 died. Again in 1877 the rains failed, and 53,532 head of cattle or nearly half the entire number in the district were estimated to have died during the year. According to the enumeration made in 1879 there were then about 80,000 horned cattle in the district, only two-thirds of the number estimated in 1875-76, and only two-fifths of the number said to have existed before the famine of 1868-69. That this estimate was very inaccurate, however, was shown by the more complete enumeration made in the following year.

Enumeration of livestock in 1880. In August and September 1880 we made an enumeration of the cattle of all sorts in the district. The opportunity was a good one, for the abundant and general fall of rain in the end of June and

beginning of July had given a plentiful supply of grass and water everywhere, and for the time each village had enough for its own cattle, so that few herds had either entered or left the district and at the time of the enumeration almost all the cattle were in their own villages. As it is usual for all the cattle to be driven inside the village enclosure every evening, all the patwari had to do was to take the village headman with him in the morning, shut all the gates of the village but one, count all the cattle that went out to work or graze, and then go from house to house and count those left in the village. I believe that the enumeration was fairly accurate and complete and that its results approximately represent the actual number of cattle owned by the residents of the district in the rainy season of 1880, being however probably somewhat under the truth. The following statement gives the number of live-stock, carts, &c., as then enumerated, with the number returned in 1875-76 (in which year the number given is the largest since the drought of 1868-69) and the number returned in 1882-83.

Year.	Horned cattle.	Horses.	Ponies.	Mules and donkeys.	Sheep and goats.	Camels.	Carts.	Ploughs	Boats.
1875-76 1880 · 1882-83	1,18,030 1,77,152 1,80,472	1,930		8.564	1.28,886 1.44,435 1,89,788		425 2.013 1,86	33,01(34,286 34,672	30 29 31

It is evident that the previous enumerations were incomplete and that the effects of the drought of 1877-78 had been exaggerated. I am inclined to believe that the losses of previous bad years were also exaggerated, but there can be no doubt that in 1868-69 a very large proportion of the cattle in the district died. At all events the number of horned cattle now in the district is nearly up to the two lakhs estimated as the number existing previous to the drought of that year. After August 1880 the rainfall was very scanty, the grass dried up, and little fodder was produced, and in June 1881 the cattle were in a most critical condition. The grass had all been completely burnt up, the fodder of the two previous harvests and the stacks of bajra straw preserved from former seasons had been almost entirely consumed, and fodder was hardly to be got at the ordinary price of grain. rain held off much longer many thousands of cattle must have died of starvation, but the plentiful rains of 1881 came just in time and very few cattle were actually lost. Three of the four following harvests gave a plentiful supply of fodder, and there was little cattle-disease, and in 1883 the cattle were perhaps more numerous and in better condition than they had been since 1868.

Bullocks and cows. 191. According to the enumeration of 1830, the number of ordinary cattle in each assessment circle was as follows:—

Asse	Assessment Circle,		Bulloeks.	Cows.	Bull-calves.	Cow-calves,	Total.
Bágar Náli Rohí Utár Hitár			1,964 8,819 38,240 2,615 3,741	4,439 7,360 38,383 2,778 3,153	2,355 3,553 13,314 1,176 1,097	2,835 3,442 15,317 1,381 1,290	11,593 23,174 1,05,254 7,950 9,281
Total of	the district		55,379	56,113	21,495	•24,265	1,57,252

The number of bullocks and cows is about the same except in the Bágar Chak where the Bágrí peasants do much of their ploughing by camels and sell their bull-calves, keeping the cow-calves, to supply them with milk. In the Nálí and Hitár the smaller number of cows than of bullocks is probably due to the large number of milch-buffaloes kept in those tracts. Far more bullocks than cows are sold out of the district, but on the other hand cows are not so carefully tended as the more valuable bullocks and their death-rate must be higher, so that the numbers of cows and of bullocks in the district probably remain about equal.

192. A great cattle fair is held annually at Sirsá during the month of Bhádúa (August-September), and at this fair a large number of cattle bred in the district are sold for export. A number of cows, buffaloes, camels, &c. are sometimes sold, but it is essentially a bullock fair; for instance in 1880 of the 19,149 animals sold, 18,541 were bullocks. The number of bullocks brought to the fair every year with the number sold and the average price have been as follows:—

	NUMBER OF I	BULLOCKS.	TOTAL PRICE.	AVERAGE PRICE
YEAR.	Brought to the Fair.	Sold.	Rs.	Rs.
1863		11,971	2,07,647	17
1864		26,188	4.83,439	18
1865	1	10,066	2,13,174	21
1866	24.953	21,953	5,22,403	24
1867	13,000	10,769	3,06,419	28
1868	15.275	11,775	2,80.758	24
1869	7.600	5,576	1,58,054	28
1870	17 000	13,854	8,90,362	28
1871	7,430	5,426		
1872	6,400	4,885	1. 16.70	Late Manager
1873	12,436	11,051		
1874	23,408	10,787	2,09,807	19
1875	14,222	5,869	1,61.703	28
1876	22,970	8,093	1,95,482	24
1877	27,625	14.031	2,89,474	21
1878	22,095	11.398	2,98.371	26
1879	28,028	22,839	6,29,522	28
1880	33,031	18,541	4,97,027	27
1881	17,491	8,901	2,84,593	30
1882	31,246	19,210	4,41,717	. 23
erage	19,071	12,659	3,26,468	25

The animals are counted as they are brought into the enclosure where the fair is held, and it is possible that some are omitted or counted twice over, but the number of animals sold is trustworthy, for each purchaser is careful to have the sale recorded by the clerks employed for the purpose, to state the price paid and to get a certificate of the sale, for which he pays a fee of ½ anna per rupee on the price. The fair is now held under direct management and the income from fees was Rs. 4,213 in 1881 and Rs. 7,114 in 1882. Notwithstanding the large concourse of men and cattle and the rude arrangements made to prevent confusion, there is wonderfully little disorder, no one attempts to evade the regulations, and although many of the dealers carry large sums in cash there are seldom any offences committed—so quiet and law-abiding are the people.

The bullocks sold at the fair are almost all young animals bred in the neighbourhood, many of them untrained, and they are bought principally by dealers who take them away to the Upper Panjáb and across the Jamna, sometimes as far as Cawnpore. The animals sold during the last seven years came from the following districts and States:—

Year.	Sirsa.	Other British districts, chiefly Hissar and Rohtak.	Bikaner.	Pattiala.	Other States.	Total.
1876	3,615	4,028	1,080		1	8,724
1877	5,324	5,909	3,099	417	9	14,758
1878	6,096	2,954	1,666	658	253	11,627
1879	11,174	8,258	2,212	1,255	294	23,193
1880	8,372	7.842	1,194	1,543	198	19,149
1881	4,395	2,371	1,215	1,135	74	9,190
1882	6,851	5,852	3,414	2,455	1,731	20,303
Average	6,547	5,316	1,983	1,068	366	15,278
Per cent	43	34	14	7	2	100

The number of cattle brought to the fair and sold and the average price depend chiefly on the nature of the seasons. The prospect of a drought and a scarcity of fodder in the neighbourhood bring a large number of cattle to the fair, as their owners having difficulty in providing for them are anxious to sell; thus in the years 1877 and 1880 the number of cattle brought to the fair was large. On the other hand, when the supply of fodder is abundant and the prospects of rabi cultivation good, owners are not anxious to sell and few cattle are brought to the fair, as in 1875 and 1881. Again, a drought in the east reduces the demand from that quarter and few cattle are sold, as in 1877 and 1880 when few buyers came from the North-West Provinces. The prices of course depend on the relation of supply and demand and vary accordingly, being ordinarily lowest in years of drought such as 1877, when many are anxious to sell and few to buy, and highest in years of plenty such as 1881, when their owners can

easily keep them at home and when many have means to buy and folder to support more cattle. The effect of the bad years culminating in 1868-69 in diminishing the number of cattle in the district may be seen in the rapid rise of price and the smaller number of cattle sold for years afterwards; but now, although the price has not again fallen so low, the numbers have recovered. On an average, of about 20,000 bullocks brought to the fair, nearly 13,000 have been sold for 34 laklis of rupees at an average price of Rs. 25 per bullock. During the last seven years nearly half the animals sold at the fair have come from the Sirsa district, and almost all of these are sold out of the district; so that on an average the Sirsá cattle-breeders sell at Sirsá fair alone a surplus stock of 6,000 young bullocks for about a lakh and a half of rupees in hard cash, or more than half the total new assessment of the district, and in some years their realisations at this fair amount to two lakhs of rupees. There is no other cattle-fair of any importance in the district, but many cattle are taken from the district to other fairs in the neighbourhood, such as those at Hissar, at Guga in Bikaner and at Jalálábád in Firozpur, and many cattle are sold to dealers in the villages, so that it may safely be estimated that on an average of years the Sirsá peasants sell out of the district a surplus of 10,000 bullocks for more than two lakhs of rupees.

193. Few bull-calves are allowed to grow up as bulls. The lucky animal selected to succeed his sire as the lord of the village-herd has an easy time of it. is given no work to do and is often allowed to help himself to whatever takes his fancy in the fields. The peasants are careless about the quality of the bull, though they do try to select a promising animal and appreciate the value of the bulls which have been obtained in recent years from the Hissar Cattle Farm and distributed among the villages with the view of improving the breed. A bullock is expected to do full work from the age of four to nine, and sometimes survives work and droughts to the age of 15. His work is constant but seldom very severe, and he is carefully tended by his master, especially if he is a valuable animal and his master a Sikh. A young bullock grazing. in the prairie is sometimes given 8 sers of chopped straw in the house daily, and a bullock in full work may get 21 sers of grain and 15 sers of fodder every day. In the Dry Tract the bullocks are brought home to the village in the evening and tied in the owner's courtyard or sometimes in a covered shed, and are there fed in the evening and again in the morning before they go out to work. They are given a rest and a feed in the course of the day and if there is grass within reach are allowed to graze for an hour. When not at work they are often, if not very valuable, driven out to graze with the cows of the village. In the Hitar the bullocks are often kept in temporary shelters at the well both night and day and fed there in troughs with chopped straw and green fodder. The bullocks used on wells by the Musalmans of the Hitar are small and feeble as compared with those in the dry uplands, where the Sikhs especially have excellent bullocks. The Aráins on the Ghaggar also usually have very fine animals. The average price of a

young bullock at the Siršá fair has varied from Rs. 17 to 30, and may be taken as Rs. 25. An ordinary bullock in the Hitár costs from Rs. 20 to 30, but an ordinary animal of the kind used by the Sikhs costs from Rs. 40 to 50, and a good bullock fetches Rs. 60 or more. In 1876, a hundred standard artillery bullocks were bought at the Sirsá fair at an average of Rs. 50 each.

194. Few cows are sold out of the district; they are kept for breeding purposes and for mik. At the Sirsa Cows. fair only about 100 cows are sold annually and their average price there is from Rs. 8 to Rs. 9; but these are poor specimens. The price of a fair cow in the Hitar may be taken at from Rs. 15 to Rs. 25, and in the Rohi from Rs. 25 to Rs. 35, but sometimes as much as Rs. 45 orRs. 50 is paid for a good cow giving 8 sers of milk daily. A cow calves after nine months of pregnancy, generally between January and June. She drops her first calf when four years old, and often gives six calves or even seven before she ceases bearing. A cow's life is stated as about 12 years. Cows are milked twice a day, morning and evening; for the first fortnight the calf is allowed to take all the milk, and for three months it gets half the milk and then a quarter of it. A Sirsa cow will not allow herself to be milked unless her calf is present, and the milker ties the calf to her leg before commencing operations. She can hardly believe that it is the calf that is milking her, but this is a form with which she will not dispense. An ordinary cow gives about 4 sers of milk a day from which 2 or 3 chhatánks of ghí can be extracted. sells in Sirsá at about 16 sers per rupee, but in the villages it is much cheaper. The cows of the village are ordinarily driven out every day to graze in the common, and if a cow is in milk she sometimes get 2 sers of grain and 10 sers of fodder daily; but ordinarily she has to be content with some chopped straw.

Buffaloes.

195. The number of buffaloes in the district in 1880 was as follows:—

Assessment Circle.		Male Buffaloes.	Female Buffaloes.	Total.	
Bágar Nálí Rohí Utár Hitár		361 2,120 1,819 101 237	1,076 3,621 8,187 517 1 861	1,437 5,741 10,006 618 2,098	
Total of district		4,638	15,262	19,900	

Buffaloes cannot stand heat well and are seldom made to work in this district, but sometimes a buffalo may be seen in a cart or plough or working at a well, yoked along with a bullock. Male buffaloes are ordinarily sold to dealers from the districts farther, north where they are used in cultivation, while buffalo-cows are kept for milk and breeding purposes. Thus three-fourths of the buffaloes in the district are cows, and almost all the buffaloes sold at the Sirsá fair, where

rometimes nearly a thousand change hands, are males (jhotá). The average price of a male buffalo at Sirsá fair is about Rs. 12, but sometimes Rs. 15 or Rs. 20 can be got. Buffalo-cows are highly valued for their milk. A buffalo calves when five years of age, in the eleventh month of pregnancy, usually about July; she ordinarily gives six or seven calves at intervals of two years. The calf is allowed to take all the milk for the first month, and then half the milk for three months and a quarter of it for three months more. A buffalo gives milk for about a year. She is milked only once a day in the evening, and an ordinary buffalo gives about 6 sers of milk a day, from which sometimes as much as half a ser of ghi can be extracted. The buffaloes of the Ghaggar and Satlaj river-sides are much superior to those of the dry uplands. On the Satlaj a good buffalo-cow giving 10 sers of milk a day fetches as much as Rs. 100, and the ordinary price may be taken as about Rs. 60 or Rs. 70 in the Hitar, and Rs. 40 to Rs. 50 in the Rohi. A buffalo is generally considered to consume nearly twice as much as an ordinary cow, and the grazing and watering fees for buffaloes are usually double those for cows. A good buffalo-cow is well looked after and when giving milk sometimes gets 31 sers of grain and 15 sers of fodder daily; but an ordinary buffalo-cow is driven out with the village cattle to graze in the common, and gets a little chopped straw at home. and perhaps a feed of grain.

The Sirsá district has for many years produced a large quan-Price of tity of ghi over and above its own requirements Produce for export northwards to Firozpur and eastwards towards Delhi. With the improvement of communications and the increase of population the price of ghi has gradually risen, and as it is valuable in comparison with its bulk its price has not been subject to such violent fluctuations as that of the heavier food-grains, although it has varied considerably with the number of cattle and the supply of fodder. Like the other produce of the district, ghi made its first decided start upward in price after the drought of 1860. From 1850 to 1859 its price remained pretty steady between Rs. 10 and Rs. 15 per maund; in 1860, when a large number of cattle died in the scarcity. it rose to Rs. 18; and remained between Rs. 15 and Rs. 18 until 1864. In that year a large number of cattle died from disease and starvation, and it was probably in consequence of this that in 1865 the price rose to over Rs. 20 per maund. The next three years were bad for the cattle and the price went on steadily rising until, after the drought of 1868 which killed a large proportion of the cattle, the price of ghi was from Rs. 30 to Rs. 34 per maund. It fell during the next two years to about Rs. 20, and remained thereabouts until the drought of 1877 which again carried off a large number of the cattle. It then steadily rose to Rs. 25 or Rs. 30 per maund, or about double the price of thirty years ago. Ghi is not much consumed by the poorer peasants except on festive occasions, when the consumption is sometimes enormous. .It is sold to the richer classes, or stored for sale and export. The average imports of ghi into the municipalities during the last eight years have been as follows :-

			2,300 maunds.
•••			2,600 ,,
•••	••••	•••	300 ,,
	ale I to	•••	125 ,,
•••		• • • •	100 "
	Total		5,425 maunds.
	· · · · · · · · · · · · · · · · · · ·		

A good deal of this comes from Bikaner, but on the other hand a good deal of ght is exported from the district without passing through the towns, and it will be safe to estimate the average annual surplus produce of ghi in the district at 4,000 maunds, which at Rs. 20 per maund represents an income of Rs. 80,000 to the cattle-owners of Sirsá from this source alone.

197. Cattle-disease of some kind is always present in the district, but is rarely very wide-spread or fatal. Accord-Diseases of cattle. ing to the reports annually made by the patwaris, about 4,500 cattle are attacked on an average every year, and of these The worst year of late was 1874-75 when 7,000 about 2,000 die. cattle died. The most fatal disease is sitlá or cow-pox which occurs at all seasons of the year, and from which the animal attacked seldom recovers. The sick animal is put in a closed stable and protected from the cold, and is sometimes given balls of pounded bhang. mouth disease (munh-khur) is common but seldom fatal. Sores form in the bullock's mouth and on its feet and it loses its appetite and gets very thin and miserable. By way of remedy a pound of molasses (gur) is put in its mouth, which is tied for some hours so as to keep it shut. Viláya or til seems to be a kind of rheumatism and is rarely. fatal; the animal affected gets stiff and unable to walk. It is said to be caused by eating a small black insect covered with a spittle-like secretion which appears in the rains, and the treatment is to give the bullock half a pound of onions and to tie a wisp of dry grass in its mouth. Golf ki sat is fatal and there is no remedy for it; it seems to be anthrax fever, and the swellings which appear on the animal's body are ascribed to congulation of the blood. When juan or maggots appear in the skin, the part affected is rubbed with a solution of tobacco. Buffaloes are subject to dhá or tákú which seems to be rheumatism, as the legs stiffen and the animal is unable to walk and loses its appetite. It is ascribed to wallowing in water heated too much by the sun, and is treated by shutting the sick buffalo in a warm stable, bleeding it at the ear and giving it dried dates as medicine. Galghuta or malignant sorethroat is often fatal; the neck swells and the animal has difficulty in breathing. One remedy is to scorch the neck by applying burning grass to it, and another is to get a holy man to exorcise the disease by making mesmeric passes (jhárná) over the part affected. Little care is taken to guard against contagion by segregating diseased animals, and the wonder is that cattle-disease does not spread more rapidly than it does. It is generally thought sufficient to tie a charm over the village gateway so that the cattle may pass under it on their way to and from the pasture-ground.

198. Almost every village in the district has still a considerable area of pasture-land, and it is usual to send out Cattle-breeding in Sirsa. all the unemployed cattle of the village daily under the charge of a cowherd to graze in the village common, and when there is no crop on the ground, in the cultivated fields also. bullocks and more valuable milch-animals are sometimes, but rarely, kept by their owners apart from the rest of the herd, but usually all the cows and buffaloes and all the calves with the village bull are driven ' out into the common in the morning after milking-time, and brought back again in the evening. There is sometimes great difficulty in supplying them with water. So long as there is water in the villagepond they are allowed to drink and wade in it, but when the pond dries up, water has to be drawn for them from the well every morning and evening with great trouble; and if, as is the case in many villages, the water of the well is too salt to drink, the cattle have to be driven daily sometimes as much as five miles to some neighbouring village to drink from the pond or well there; and a considerable sum has often to be paid for this privilage (pilet). In many villages a customary grazing-fee (bhunga) is paid by the cattle of the village, and usually a higher fee is charged on any cattle belonging to other villages which may come to graze on the village pasture-land. There are still vast tracts of unenclosed prairie in Bikaner, and there the same custom still prevails that was prevalent in Sirsá before the spread of cultivation. In the rainy season, after the annual growth of grass has sprung up, large herds of cattle are driven south into the prairie from Sirsá. Pattiála and even from as far north as Ludhiána, and are kept grazing there until the grass dries up, when they are driven northwards home again. Considerable fees are paid to the Bikaner State and local authorities for permission to pasture and water these herds of cattle (gol): sometimes as much as Rs. 2 per bullock for grazing and Re. I for watering; but commoner rates are 8 annas for grazing and 4 annas for watering. The growth of grass on the Bikaner prairies, like that on the Sirsa uplands, is wholly dependent on the local rainfall, and when, as in 1880, rain fails generally in that neighbourhood, no grass is produced and no cattle are driven southwards. In such years of drought little or no grass grows on the village commons and the cattle are left in dependence on the produce of the cultivated land. Mr. Oliver was of opinion that the prairie was being brought under cultivation too rapidly, and urged that in the interests of the cattle steps should be taken to check the spread of cultivation in the more advanced parts of the district, and to reserve a large area in each village as a grazing-ground for the cattle, and similar proposals have recently again been made. But however desirable such arrangements may be to keep up a supply of fresh green fodder in ordinary times; they would have little effect in protecting the cattle from starvation in years of drought; for in such years the uncultivated land produces almost nothing, and

indeed up to a certain limit not yet nearly reached in Sirsa, cultivation actually increases the produce of fodder, and especially of storeable fodder, and thus renders the district better able to support its cattle in years of drought. The average produce of straw even on unirrigated fields may be estimated at four maunds per acre, which is much more than the storeable produce of grass on the same land when uncultivated; and even in a year of drought like 1880, when the grass-wastes produced almost nothing, the cultivated land was estimated to have produced from two to three maunds of fodder per acre. Pála too, which is considered one of the best fodders, grows more plentifully on cultivated than on uncultivated land. Considerable efforts are made by the people to store up fodder for their cattle. Some crops, such as turnips, gwar and moth, are cultivated chiefly for the fodder they give, and care is taken to gather and store not only the produce of these crops, but the straw of barley, wheat, jawar and even bajra, stacks of which may be seen in the fields and about the homestead of almost every village. Pála and some of the better grasses are also cut and stored. It seems that greater care is now taken than formerly to store fodder in these ways and preserve it for seasons of drought, and that the experience of the last two or three scarcities has taught the people to use every available means of storing fodder against such seasons. But as a rule all that they can do is to store up a quantity sufficient to support their cattle during the dry months of the hot weather, when there is no green food available and the cattle have to be supported wholly on the stored fodder. This gets exhausted towards the end of June, and if the rains then fail and no grass springs up, the cattle are left without afood, and numbers of them die. -Yet, except perhaps for a short time after one of these recurring scarcities. the district has always more cattle than it requires for agriculture, and can afford to export a large number of young bullocks. If the peasants of the district were to reduce their stock of cattle and breed fewer, the supply of fodder would be sufficient to support a larger proportion of them through a drought, and the loss of cattle from starvation in a season of scarcity would be smaller, but then their surplus stock and their profits from the sale of young cattle in ordinary seasons would be less. Cattle-breeding in such a country is a very speculative business, and the peasants seem to find it more profitable in the long-run to allow their cattle to multiply up to the limit of subsistence, that is, up to the number which can be supported by the year's fodder until the usual season for a new growth of grass, and to take their chance of the rains failing. If the rains come as usual, the speculation is a success, and the cattle are safe for another year; if they fail, the speculator loses his profits and some portion of his capital, but one or two good seasons soon make it up to him again. It is not improbable that improvements in the methods of storing fodder would be utilised by the Sirsá peasant not so much in guarding against the consequences of drought as in multiplying his stock still further, and taking his chance of the rains as before; and this is perhaps in the circumstances the most profitable way of conducting his trade as a cattle-breeder.

It is probable that, whatever improvements in the way of storing fodder and increasing its supply be made available in Sirsa, there will always be great mortality of cattle in seasons of drought. Still every increase in the supply and improvement in the method of storing must tend to some extent to render the food-supply of the cattle less precarious and to diminish the mortality in droughts, while the increase in the value of cattle makes it better worth the owner's while to take precautions to protect them from starvation. There cannot be a doubt that the district is much better off in this respect than it was. The annual produce of fodder is increasing, instead of diminishing, with the spread of cultivation, and if the number of cattle in the district is now fewer than it used to be (which I doubt), the mortality in a season of drought is much less, as they are less dependent on the precarious produce of the prairie and have larger stores of fodder to supplement it. It is probable that the drought of 1880 which caused hardly any deaths among the cattle would twenty years ago have killed a considerable proportion of the total number.

I am also inclined to doubt whether, as Mr. Oliver reported, the breed has really deteriorated in quality. At the time he wrote the cattle seem to have been suffering from a series of bad years and to have been greatly subject to starvation and disease. For a number of years now disease has not been seriously prevalent and the fodder-supply has not been so precarious as it was. The herds are certainly not yet so small as to lead to any deterioration from breeding-in. It is possible that the natural grasses developed a better breed than the straw and leaves of cultivated crops, but nearly half the area of the district is still virgin prairie, and there are boundless stretches of grassland within easy reach in Bikaner. Perhaps the good quality of the breed of this tract is partly due to the frequent droughts which killed off the weaker animals and thus exercised a sort of natural selection leading to the survival of the fittest; and this principle is still at work. It is true that the Musalmans of the Satlaj have an inferior breed of cattle and that those owned by the Bagris and Musalmans of the uplands are only of fair quality, but the breeds owned by the Sikhs and the Arains are excellent, and finer bullocks than they have are seldom to be seen anywhere. The demand for Sirsá bullocks and the prices they fetch show no signs of falling off, but the contrary, and this would seem to show that the breed is at least as good as it used to be. Some attempt has been made to encourage artificial selection for breeding purposes by offering prizes at the Sirsá Cattle Fair for the best animals and by introducing good bulls from the Hissar Cattle Farm, and although the progress made in this direction has as yet been little, there are signs that the peasants are learning to appreciate the advantages of artificial selection sufficiently at least to prevent the breed from deteriorating.

199. The number of camels in the district in 1880 was as follows:—

Assessment Circle	ADULT CAMELS.		Young	TOTAL,	
	Male.	Female.	Male.	Female.	
Bågar Nálí Rohí Utár Hitár	823 •712 2,242 234 24	1,208 993 4,931 705 38	486 455 1,998 184 10	342 • 281 1,201 135 10	2,859 2,441 10,422 1,358 82
Total of district,	4,135	7,925	3,133	1,969	17,162

This is a much larger number than had ever been returned before. Camels are kept chiefly in the Dry Tract by the Bágris who use them not only as beasts of burden, but do their ploughing with them also. Taking young and old together, there are 7,268 male camels to 9,894 female, the reason for the difference being that male camels are more readily sold than the females which are kept for breeding purposes. In 1878-79, during the Afghan War, orders were issued to impress camels and send them to the front, and camel-owners to escape impressment promptly sent their camels into Bikaner, so that the number returned as in the district was only 1,099. The number required was therefore distributed over the villages of the district in proportion to their assessment, and each was required to produce so many whether it had them or not. The peasants of most of the villages had to buy the camels, which they took to the tah-il and made over to the tabsildar. Few of them ever returned and most of those that disappeared were never paid for. If the 2,792 camels thus impressed be taken at Rs. 70 each, it seems that the first Kabul campaign cost the Sirsa peasants in camels alone nearly two lakhs of rupees -a large contribution to the expenses of the war. For the second campaign 915 camels were bought by Government in 1879, and 975 in 1880, making a total drain of 4,682 in three years. These camels were all supplied through the Deputy Commissioner of Sirsá, and though many of them came from Bikaner, the number of camels in the district must have been much reduced. Yet it was after this drain that we enumerated 17,162 in the district. The average price paid by Government was Rs. 77 in 1879 and Rs. 97 in 1880, and the total price paid from the Sirsá Treasury for camels during the two years 1879 and 1880 was Rs. 1,65,000. The circumstances were exceptional, and ordinarily the number sold out of the district and the price paid are much less. few camels are sold at the Sirsa Fair, and the average price is only from Rs. 50 to Rs. 60. But it may be estimated that on an average of years about 1,000 camels are annually sold out of the district at an average price of Rs. 60, giving a total value of surples camels of Rs. 60,000 per annum. The Bagris have often very good riding

camels which can go eight miles an hour for several hours at a time, or travel 70 miles at a stretch. Such a camel fetches Rs. 150 or more, but an ordinary male camel fetches from Rs. 70 to Rs. 90, and an ordinary female from Rs. 50 to Rs. 80. A camel begins to work at four years of age, and a female gives her first young in her fifth year, after thirteen months of gestation, and bears five or six times at intervals of two years. A camel's lifetime is considered to be 15 or 20 years. An ordinary camel-load is 6 maunds. Camel's milk is often consumed as food, and the hair (jat) is shorn and made into ropes and sacks. When a camel is in full work, he sometimes gets 2 sers of grain and 15 sers of moth or pála leaves daily, but ordinarily he is turned out into the prairie to pick up what he can find. Although camels are subject to many diseases, they eat almost any kind of plant and can live on food that does not support horned cattle, and are not like them liable to die off in great numbers in times of scarcity, so that in this district they are not so precarious a property.

Among the numerous ailments to which camels are subject the following may be noticed :- In báo-bája the eyes water badly and sometimes the animal cannot raise his head or move his legs. The part affected is branded, or he is given 4 sers of sugar in 4 sers of sweet oil, or is bled below the eyes. In hubbi the neck swells and the month waters and the animal ceases to wag his tail. The diagnosis is made by applying fire to the animal's tail; if that does not cause him pain he has certainly got hubbi. The proper medicine is hedgehog soup, made by boiling a hedgehog in water with a pound of red pepper; another medicine is a solution of roasted salt. Kupálí is supposed to be due to a growth on the brain which causes the camel to keep his head constantly raised in the air. The cure is branding the head, which causes the growth to burst and the matter to come out through the ears. In ras the camel's hindlegs get quite stiff and he becomes lame; the legs are poulticed and oil is given as a medicine. In pátlí the camel's nose gets filled with blood so that he cannot breathe properly: it is cured by clearing out the nostril with the fingers or by drawing two lines over the nose with a heated

Horses and donkeys.

200. The number of horses and donkeys was returned in 1880 as follows:

Assessment Circle.	Horses.	Mares.	Ponies.	Mules & Donkeys.	Total.
Bágar Nálí Robí Utár Hitár	52 146 428 61 60	166 285 468 11 253	52 621 135 14 88	425 4 687 2,527 92 833	695 5,739 3,558 178 1,234
Total of district	747	1,183	910	8,564	11,404

The distinction between horses and ponies is not trustworthy. The horsestesh of the district generally consists of wretched little ponies, some of