

ON
BENGAL DYSENTERY

AND
ITS STATISTICS,

WITH
A NOTICE OF THE USE OF LARGE ENEMATA IN
THAT DISEASE,

AND OF
QUININE IN REMITTENT FEVER.

By
JOHN MACPHERSON, M.D.,
1ST ASSISTANT PRESIDENCY GENERAL HOSPITAL.

CALCUTTA:
W. THACKER AND CO., ST. ANDREW'S LIBRARY.

1850.

CONTENTS.

	PAGE.
INTRODUCTION.	
Treatment,	1
GENERAL STATISTICS.	
Mortality from Dysentery,	3
Prevalence of the Disease and Mortality from it, according to season,	7
PATHOLOGICAL FACTS.	
Preliminary Remarks on Tables,	12
Table of Acute Dysentery,	13
Summary of Table of Acute Dysentery,	34
Table of Chronic Dysentery,	35
Summary of Table of Chronic Dysentery,	44
On Chusan Dysentery,	45
Comparison of Acute and Chronic Dysentery,	46
True description of the changes produced by Bengal Dysen- tery,	47
Frequency of complication with disease of Liver,	48
On certain appearances mentioned by authors,	50
Conclusion,	51
NOTES.	
On Treatment,	52
On Dysentery in Children,	53
APPENDIX.	
On the use of large Enemata in Dysentery,	54
On the use of Quinine in Remittents,	57

ON
BENGAL DYSENTERY

AND
ITS STATISTICS,

WITH
A NOTICE OF THE USE OF LARGE ENEMATA IN
THAT DISEASE,

AND OF
QUININE IN REMITTENT FEVER.

By
JOHN MACPHERSON, M.D.,
1ST ASSISTANT PRESIDENCY GENERAL HOSPITAL.

CALCUTTA:
W. THACKER AND CO., ST. ANDREW'S LIBRARY.

1850.

CALCUTTA:
PRINTED AT BISHOP'S COLLEGE PRESS.

CONTENTS.

	PAGE.
INTRODUCTION.	
Treatment,	1
GENERAL STATISTICS.	
Mortality from Dysentery,	3
Prevalence of the Disease and Mortality from it, according to season,	7
PATHOLOGICAL FACTS.	
Preliminary Remarks on Tables,	12
Table of Acute Dysentery,	13
Summary of Table of Acute Dysentery,	34
Table of Chronic Dysentery,	35
Summary of Table of Chronic Dysentery,	44
On Chusan Dysentery,	45
Comparison of Acute and Chronic Dysentery,	46
True description of the changes produced by Bengal Dysen- tery,	47
Frequency of complication with disease of Liver,	48
On certain appearances mentioned by authors,	50
Conclusion,	51
NOTES.	
On Treatment,	52
On Dysentery in Children,	53
APPENDIX.	
On the use of large Enemata in Dysentery,	54
On the use of Quinine in Remittents,	57

XVII.E 12

To

SIMON NICOLSON, ESQ., F.R.C.S.,

Surgeon of the Presidency General Hospital.

MY DEAR SIR,—To whom could I with greater propriety present these gleanings from the records of the Institution, with which your name has been so honorably associated for more than a quarter of a century, than to you, even did I not lie under a weight of personal obligation to yourself, which I am proud to acknowledge?

Had the Medical and Physical Society been in existence, or were there now any local Medical Journal, these pages would probably not have appeared in a separate form. Bengal Dysentery has been admirably described by former officers of this Hospital. The present is a slight attempt, but one of the first of its kind, to apply the numerical method to the subject. It is not an exposition of opinions, but a statement of facts, (perhaps too copious in detail,) from which the reader may draw his own conclusions. I have not thought it desirable to expand these materials, as they might easily have been expanded, into a regular treatise.

In the numerical statements it has been found impossible to obtain absolute accuracy, but I have had the advantage of comparing them with some carefully constructed tables of Mr. Hare's, and they are accurate enough for all practical purposes. You

will be surprised at the high rate of mortality that has been found to prevail, but if a rigid analysis of cases were made, and if those which were evidently the terminations of other maladies, and those which died within twenty-four hours after admission, were excluded, the mortality would be considerably less.

I should hope, that the short historical notice of the use of enemata in dysentery, and of quinine in remittents, may, at the present time, be of some interest to you and to the profession in this country.

Believe me,

My dear Sir,

Your's very sincerely,

JOHN MACPHERSON.

GENERAL HOSPITAL:

May 20th, 1850.

ON
BENGAL DYSENTERY.

INTRODUCTION—TREATMENT.

I propose—1st, to give some returns of the mortality from dysentery, and of the diffusion of the disease throughout the year, from the books of the General Hospital, and to make a few comparisons with what is found to occur elsewhere.

2nd, to give an abstract of the pathological appearances presented in 160 cases of acute and in 55 of chronic dysentery, which have occurred during the last nine years.

The symptoms and history of Bengal dysentery have been so fully described by Twining and Raleigh, that it is unnecessary on the present occasion to add any thing to what they have said on those heads: nor do I intend to say more than a few words on the management of the disease. Practitioners seem to have been gradually losing faith in the mercurial treatment, which dates back from the latter half of the last century, and has often been exclusively followed. The virtues of ipecacuanha, also a very old remedy, though they were brought prominently forward by Twining, are not so much trusted to now as formerly. Opium again, which since the days of the liquid laudanum of Syden-

ham down to the present time, has been more or less used, but which had latterly got the character of masking the disease, seems of late to have regained its rights. In Bengal, Drs. Mackinnon and Goodeve and others have suggested the propriety of returning to a more free use of this drug, and that most sound observer, Dr. Morehead of Bombay, has borne testimony to its value. My own belief is, that the return to a milder and more soothing treatment has been attended with great advantage—though it may be difficult to prove this by reference to any Hospital records within reach. One obvious difficulty in arriving at safe conclusions is, that the disease itself varies so much in intensity in different years, as well as in different periods of the same year.

Thus, H. M. 55th at Secunderabad, lost in 1837. . 1 in $4\frac{1}{2}$
 1838. . 1 in $7\frac{1}{2}$
 1839. . 1 in 10

yet the treatment was the same, and by the same Medical Officer throughout.

The only data bearing on this point in my possession are afforded by the records of the Seaman's Hospital. Such as they are, they are subjoined, though they must be taken with great reservation—1st, because the number of facts is small—2nd, because it is possible to support by statistical returns almost any pre-conceived notion.

Treatment by bleeding and calomel	} 227 cases, 48 deaths, or 21 per cent
for five years,	
Treatment by leeching, opium and	} 80 „ 10 „ or 12 per cent.
astringents for three years, . .	

Still I would not be understood as advocating any exclusive treatment. In an account of the latest Irish epidemic, a most intelligent physician, Dr. Mayne, states—“opium most certainly aggravated the disease.” “Mercury must be considered the principal remedy.” As the character of dysentery varies, so no doubt should our practice.

I may allude here, as being quite a curiosity in its way, to the most singularly successful mode of treatment that I have seen on record. Mr. Marbot tells us, that in a French vessel on the coast of Zanzibar, he has treated 300 cases of dysentery with aconite and ipecacuanha without a single casualty. Two things seem very plain: first, that Mr. Marbot deceived himself: second, that the disease he treated is something very different from Bengal dysentery.

GENERAL STATISTICS.

MORTALITY FROM DYSENTERY.

1. *Out of Bengal.* It is difficult to procure any accurate statistical data regarding the proportion of deaths to cases treated in Northern Europe, and dysentery does not occur in Major Tulloch's tables of the diseases of troops in Great Britain. Dr. Williams of St. Thomas', however, states, that more than 25 per cent. of cases of chronic dysentery die in the London Hospitals, and says, "in candour it must be allowed, that no class of diseases in them offers so few chances of recovery." In Dublin, Dr. Mayne lost 32 per cent., or omitting cases in old men after the age of 60, as much as 22 per cent.; children under 10 years died at the rate of 65 per cent. At Gemünden, in Southern Germany, where dysentery is often epidemic at the fall of the year, the mortality during the great epidemic of 1834 was 11 per cent. In the Peninsular war, scarcely more than 2 out of 3 recovered. Further South in Malta, which can scarcely be said to have an European climate, the mortality among H. M.'s troops is about 8 per cent.

The accompanying table, compiled from Major Tulloch's reports, shows the percentage of deaths to cases among H. M.'s troops in various parts of the world :

Bermuda,	3.2	Mauritius,	6.0
Nova Scotia,	7.4	Jamaica,	4.3
Canada,	5.0	Windward and Lee-	} 7.1
Mediterranean,	4.0	ward Islands, ..	
St. Helena,	10.6	Ceylon,	13.0
Sierra Leone,	18.0	Tenasserim,	8.5
Cape of Good Hope, ..	4.7	Three Presidencies, ..	8.3

2. *In Bengal* Major Tulloch gives the mortality of H. M.'s troops at 8 per cent., and Dr. Mackinnon finds, that among the European troops for 7 years at Cawnpore, it was 6.16. In Calcutta and its neighbourhood, the percentage among European troops appears to be rather more than 9 per cent.; but it varies much, for some regiments have suffered much more heavily. Thus, H. M.'s 21st Fusiliers lost in Fort William and Chinsurah, at the rate of 17 per cent.; and H. M.'s 70th, in crowded barracks at Dum-Dum, suffered at the rate of 35 per cent. during the first six months after its arrival from England.

3. In an Institution like the General Hospital, to which patients are often sent in the last stage of disease, and which receives all sick soldiers left behind by their regiments, as well as all sick invalids on their way home, one would naturally expect to find a high rate of mortality. Raleigh gives it as from 10 to 14 per cent., and Martin says, it is 10.27.

But the following table of admissions and deaths during the last 20 years will shew that they have greatly underrated it. The mortality from all "bowel complaints" is about 19 per cent., much higher than the rate assigned by them, but not equal to the reality. Of course the mortality in returns can be made to vary much according as cases are classed under the heads of diarrhœa or dysentery, a point often requiring nice discrimination.

This will be abundantly evident if we contrast the proportion of deaths from bowel complaints, with that from dysentery during the last 16 years, in periods of four years.*

	1834—38.	1838—42.	1842—46.	1846—50.
Bowel complaints, ..	68.6	71.8	70.8	69.
Dysentery, ..	24.3	21.2	25.6	26.3

Thus, while the mortality from dysentery has, on the whole, been increasing, that from all bowel complaints has varied to but a trifling extent.

The following have been the admissions and deaths from dysentery in the General Hospital, from 1830 to 1850 :

Year.	Admission.	Deaths.	Year.	Admission.	Deaths.
1830	144	22	1840	68	11
31	128	20	41	172	42
32	124	20	42	147	41
33	128	19	43	88	19
34	147	39	44	141	28
35	71	20	45	91	31
36	55	10	46	87	27
37	54	15	47	78	17
38	52	13	48	87	25
39	78	15	49	94	24

Total admissions 2044, total deaths 457, or 22.3 per cent.

The extremes of mortality have been 14.8 in 1833, and 34. in 1845.

The average is higher than that of the Bombay General Hospital, which, for a period of 5 years, was 18.3, and lower than that of the Madras one, in which, for a period of 10 years, it was 30 among Civilians, though only 5.3 for the Military. The average mortality in a series of years appears to have been almost the same in the Calcutta General and Medical College Hospitals, and the extremes in the latter have been 14.1 and 33.

The mortality in the year 1840 in the General Hospital was 16, in 1849, 25.5, and in the Medical College Hospital at the same periods 25, and 27.

* I have taken a period of 16 years, because in the year 1834 there was a sudden increase of mortality in bowel complaints, which has ever since continued. This was the year of the great Continental epidemics of dysentery, and what is perhaps more to the point, the port of Calcutta began to be crowded with ships, many of an inferior class.

In the table of admissions and deaths, acute and chronic dysentery are classed together: indeed, they cannot be separated with advantage, and their severance in Major Tulloch's tables gives rise to some very strange results. In them chronic dysentery (which is returned as infinitely more fatal than acute) is made to kill in Malta 1 in 4, while in the Mauritius it kills only 1 in $14\frac{1}{2}$, and deaths by acute dysentery are made to vary from 1 in 57 in the Bermudas, to 1 in $2\frac{1}{2}$ in Sierra Leone. There is manifestly some error in such statements.

On analysing the classes, among whom the mortality occurred in the General Hospital during the only two years, 1847 and 1849, in which that analysis can be easily made, we find that the mortality of the Military in those years was 10 and 21 per cent., giving a mean of $15\frac{1}{2}$ per cent., which is considerably less than that of the non-military.

It might be expected, that some statement should be given of the average period after the first invasion of the disease, or after admission into Hospital in which death occurs, and it may be stated, that death within a week from the first attack is extremely rare. But patients are so seldom brought to the General Hospital at the commencement of the attack, and the attack itself varies so much according to the prevailing character of the disease, that no such statement would be satisfactory. In like manner, almost all the dysenteric patients being soldiers and sailors, are between the ages of 18 and 40, and the chief mortality is of course between those two periods of life. Most admissions also occur between the ages of 18 and 30, and consequently most deaths.

The number of fatal cases among women is extremely small: for in the following tables of the appearances in 215 dissections, the names of only 5 females are found. I have imagined, that Bengal acute dysentery is peculiarly fatal in boys from 14 to 20 and in middle aged men: boys suffer less from the chronic form. I have been repeatedly astonished to discover after death an immense extent of structural

change in boys whose illness could not be ascertained to have exceeded 8 or 10 days.

It may be finally remarked, that about 1-10th of all admissions into the General Hospital are cases of dysentery, and that while 10 to 11.5 per cent. is the average mortality on all admissions, that on dysentery is about 22, cholera and hepatitis alone proving more fatal.

PREVALENCE OF THE DISEASE AND MORTALITY FROM IT, ACCORDING TO SEASON.

I. 1. *Out of India.*—The following are a few notices of the prevalence of the disease, according to season, in Europe. In the Infirmary for children in London, Dr. West found the rate of prevalence of dysentery to be, spring 8.3: summer 13: autumn 24.4: winter 7.2. In the last Dublin epidemic, in the year 1847, the admissions in the Work-house were, 1st quarter 136: 2nd quarter 159: 3rd quarter 206: 4th quarter 157:—and at Grätz, where dysentery is an autumnal disease, the distribution was—

Spring 8 cases: Summer 67: Autumn 163: Winter 11.

2. *In India*, Annesley found on the large scale that out of 13,900 cases which occurred in 5 years in Bengal, 2,400 were in the cold, 4,500 in the hot and dry, and 7,000 in the hot and moist season. This accords with the results in Ceylon among Queen's troops, who have most cases of dysentery in the second and third quarters of the year. In like manner Dr. Mackinnon found at Cawnpore, in a series of 7 years, that dysentery among Europeans was most frequent in the rains, next in the hot weather, least frequent in the cold season, and far most frequent in the months of August and September. The results in the General, the Seaman's, and the Medical College Hospitals, will be found generally similar, although the disease is in Calcutta somewhat more prevalent

in the cold than in the hot weather. These points will be illustrated by the following tables. The prevalence of the disease, according to season, is modified by the climate of the particular place, and we thus find, that in the General Hospital at Bombay, it is said to be most common in the cold season.

Table of Admissions and Deaths from Dysentery in the General Hospital, for 10 years.

Months.	1840.		1841.		1842.		1843.		1844.		1845.		1846.		1847.		1848.		1849.		Total Admission.	Total Deaths.	Percentage of Deaths.
	Admitted.	Died.	Admitted.	Died.	Admitted.	Died.	Admitted.	Died.	Admitted.	Died.	Admitted.	Died.	Admitted.	Died.	Admitted.	Died.	Admitted.	Died.	Admitted.	Died.			
Jan. . .	6	1	14	2	7	1	8	3	10	0	18	3	10	2	6	1	21	4	7	1	107	18	16
Feb. . .	5	2	7	2	6	2	19	0	5	2	5	0	3	2	6	2	5	1	4	1	61	14	23
Mar. . .	6	1	13	3	7	2	8	2	5	2	1	1	1	2	4	2	3	0	4	1	52	17	32.6
April, .	6	0	4	1	4	1	6	1	6	2	2	1	7	1	7	0	7	2	7	1	56	10	14.2
May, . .	4	0	13	3	8	3	2	2	5	1	4	0	9	3	8	3	7	1	8	5	66	21	31.8
June, . .	1	2	6	3	10	2	6	1	10	7	9	3	3	1	3	1	6	2	5	0	59	22	37
July, . .	4	1	21	2	15	3	7	3	13	7	10	5	11	3	8	3	5	2	12	3	106	32	30
Aug. . .	6	0	17	7	18	7	6	3	10	1	11	5	7	2	5	0	9	2	10	1	98	28	28
Sept. . .	5	1	19	5	25	8	6	0	3	2	4	2	7	3	5	1	9	3	7	4	96	29	30
Oct. . .	11	0	21	7	10	7	5	0	5	2	5	1	3	1	7	3	8	4	12	3	87	28	32.1
Nov. . .	7	1	4	4	25	2	8	2	21	1	8	2	16	2	10	1	4	3	9	2	112	20	17.8
Dec. . .	7	2	33	3	12	3	7	2	48	1	14	7	10	5	9	0	3	1	9	2	152	26	17
Total,	68	11	172	42	147	41	83	19	141	28	91	29	87	27	78	17	87	25	94	24			
Percentage of Deaths.	} 16		24.4		27		21.5		20		31.8		30		21.8		28.7		25.5				

The average prevalence of the disease in the different months is fairly enough represented in the foregoing table: if we make a few corrections to allow for the increase of cases caused by the arrival of the invalids of the season in

the end of November, and in December and January, the number of cases would stand thus—

January, 75	July, 106
February, 60	August, 98
March, 52	September, 96
April, 56	October, 87
May, 66	November, 82
June, 59	December, 80
1st half of the year, 368	2nd half of the year, 549

or, according to season, in the following proportions: four cold months 74, three hot ones 68, five of rains 88.

The results are nearly the same in the College Hospital.

3. Regarding the degree of prevalence of the disease in different years, nothing of distinct value can be gathered from the Hospital records, because the number of admissions is in great measure dependant on the number and strength of detachments of troops arriving at the Presidency and on the number of shipping lying in the river. Thus the year 1842, in which there were most admissions from bowel complaints, was that of the return of the invalids from Chusan.

II. Regarding the rate of mortality according to season, out of India I possess no data, but if we proceed to investigate the rate of mortality in the General Hospital, we find it to be in this proportion, almost equal in the first and second halves of the year; but according to season, cold weather 18.4: hot weather 26.2: rains 31.4.

The results in the College Hospital are nearly the same. The most fatal months in the General Hospital have been March, May, June, and October, and in the College Hospital May, June, August, September and December. In Ceylon the disease was most fatal in April, May, and June, while at Cawnpore, October and November were the worst months. This comes nearest to Bombay, where the cold season is said to produce most deaths, and the monsoon, or July, August and September, next most.

The information on this point is not very full or satisfactory, but probably for India generally, August, September and October will be found to be the most fatal months, as the fall of the year is in Europe, though perhaps from different causes.

The unusually high mortality in the General Hospital in the month of March, when the prevalence of the disease is smallest, cannot be very readily explained.

Nor can any satisfactory information be given as to the rate of mortality in different years. It was remarked above, that the average mortality in a *series of years* was the same in the General and in the Medical College Hospitals, yet in *individual* years the mortality in the two institutions differed most widely.

Through the kindness of Dr. Mouat, I am enabled to give the following tables of admission and deaths among Europeans and Natives in the Medical College Hospital, during the last 10 years.

Table of Admissions and Deaths of Europeans from Dysentery, in Medical College Hospital.

Months.	1840.		1841.		1842.		1843.		1844.		1845.		1846.		1847.		1848.		1849.		Total Admission.	Total Deaths.	Percentage of Deaths.
	Admitted.	Died.	Admitted.	Died.	Admitted.	Died.	Admitted.	Died.	Admitted.	Died.	Admitted.	Died.	Admitted.	Died.	Admitted.	Died.	Admitted.	Died.	Admitted.	Died.			
January, ...	8	0	2	0	10	1	3	0	8	0	9	2	9	1	13	4	8	0	12	2	82	12	14.6
February, ...	4	1	1	0	12	2	4	0	10	2	8	1	13	1	11	1	10	2	8	2	81	13	16
March, ...	4	1	5	0	8	2	7	2	10	2	10	0	15	1	10	1	13	0	11	2	93	11	11.8
April, ...	5	0	3	0	9	5	8	1	7	0	14	3	6	2	11	1	16	3	11	1	90	16	17.7
May, ...	4	2	7	2	12	5	9	2	1	1	11	4	9	2	10	1	15	4	13	7	91	30	33
June, ...	4	1	10	2	6	2	11	3	0	0	9	4	6	1	10	0	16	1	13	7	85	24	28
July, ...	4	0	8	3	7	3	13	3	2	0	21	3	18	4	15	2	18	6	16	3	124	27	21.7
August, ...	4	2	9	5	13	4	16	4	5	2	17	4	18	3	12	4	23	4	12	2	129	34	26.2
September, ...	4	2	17	7	6	2	17	4	10	2	16	3	18	2	10	2	14	6	13	3	125	33	26.4
October, ...	7	2	12	3	10	2	14	2	9	1	12	1	16	3	13	3	10	2	11	4	116	23	19.7
November, ...	5	2	7	2	7	3	13	3	11	3	18	2	14	0	11	3	19	1	10	2	127	21	16.7
December, ...	2	1	6	1	12	6	12	7	16	4	24	5	14	2	10	1	17	3	4	1	117	31	26.3
Total, ...	55	14	87	25	112	37	127	31	89	17	169	32	156	22	136	23	179	32	134	36			
Percentage of Deaths ..	25.2		28.7		33.0		24.4		19		19		14.1		17		17.8		27				

Table of Admissions and Deaths of Natives from Dysentery, in the Medical College Hospital.

Months.	1840.		1841.		1842.		1843.		1844.		1845.		1846.		1847.		1848.		1849.		Total Admission.	Total Deaths.	Percentage of Deaths.
	Admitted.	Died.	Admitted.	Died.	Admitted.	Died.	Admitted.	Died.	Admitted.	Died.	Admitted.	Died.	Admitted.	Died.	Admitted.	Died.	Admitted.	Died.	Admitted.	Died.			
Jany.	5	1	8	1	5	2	5	2	6	1	8	2	8	1	11	1	7	0	6	0	68	11	16
Feby.	5	0	3	2	4	2	3	0	6	1	7	0	6	1	7	1	8	1	9	2	58	10	17
March	3	3	4	0	3	1	5	1	6	2	9	0	8	0	7	1	6	1	5	0	56	9	16
April,	4	1	3	0	2	1	6	2	2	0	8	3	7	1	6	0	7	1	7	1	52	10	19
May,	8	0	5	0	3	1	9	4	4	1	9	0	8	0	7	0	7	0	9	1	69	7	10.1
June,	5	4	6	1	6	0	5	1	5	0	11	0	2	0	4	0	5	0	7	2	56	8	14
July,	6	2	3	1	5	1	3	1	6	2	7	2	7	2	8	0	7	0	7	0	59	11	18.6
Augt.	7	2	6	0	6	1	8	2	10	1	8	1	4	2	12	4	6	1	6	2	73	16	21.9
Sept.	5	2	4	1	5	1	4	2	7	2	8	0	6	0	4	0	11	0	9	0	63	8	12.6
Octr.	4	0	3	0	3	1	6	1	9	1	5	1	9	2	9	2	7	0	5	0	60	8	13.3
Novr.	4	3	4	1	5	1	7	2	7	1	8	1	4	0	10	0	8	2	3	1	60	12	20
Decr.	4	2	2	0	3	0	9	0	10	2	8	2	8	1	7	1	8	2	2	0	61	10	16.2
Total,	60	20	51	7	50	12	70	18	78	14	96	12	77	10	92	10	87	8	75	9			
Percentage of Deaths.	33		13.7		24		25.6		18		12.4		13		10.8		9.2		12				

Though it is foreign to the object of these pages to treat of dysentery among Natives, it is worthy of remark, how uniform the rate of admission among them seems to be throughout the year. Though they do not vary much, the admissions are most numerous in August and in May, while the mortality is highest in August, and next highest in November. The general results are much the same as with Europeans.

The average mortality among Natives has been 16.9, that among Europeans 22.5. This accords with general experience, which has shown the disease in Natives to be more amenable to treatment than in Europeans.

PATHOLOGICAL FACTS.

PRELIMINARY REMARKS ON TABLES.

The following tables exhibit a true representation of the structural changes commonly effected by fatal Bengal dysentery. Cases complicated with pthisis, syphilis, or any other constitutional taint, have been, as far as possible, excluded,—(and here I cannot help remarking, how strange it seems that Rokitansky should have asserted the antagonism of phthisis and dysentery, which in this place so commonly occur together.) An abstract is given of the pathological appearances in all cases which are at all fully recorded in the Hospital books, so that the reader may draw his own conclusions from them. A few results only have been noted, but every opinion of a theoretical nature is studiously avoided. On going over the records of these cases it is impossible not to be struck with the fact, that no two observers saw with the same eyes. Thus one gentleman has invariably found the mesenteric glands enlarged, while another describes the liver as dry in one half of the cases in which he mentions its condition. Where there has been no notice of the state of a particular organ, the space is left blank. In such cases its condition may be presumed not to have differed widely from the normal one. In the column of remarks any striking variation from the usual symptoms is noted.

ACUTE DYSENTERY.

No.	State of large Intestines.	State of small Intestines.	State of Liver and Gall-bladder.	State of Mesentery, Mesenteric glands or other organs.	REMARKS.
1.	Cæcum and colon in a sloughing state throughout.	Cicatrix of ulcer on lower surface of right lobe; bands of adhesion.	Seaman—died in 3 weeks.
2.	Cæcum ulcerated and sloughing, firmly tied to omentum: ulcers in transverse arch and sigmoid flexure of colon.	Stomach natural.	Liver natural.	Seaman, ætat 42—died in 20 days.
3.	Cæcum slightly injected, one or two small points of ulceration: these increased in the transverse arch and sigmoid flexure. Rectum one mass of ulceration.	Small intestine healthy.	Liver healthy. Gall-bladder with healthy bile.	Soldier caught dysentery from sleeping on deck—died in 20 days.
6.	Spotted ulceration of large intestine, increasing as it went downwards: internal ulceration, not perceptible outside.	Small intestine distended.	Liver large, pale and soft. Gall-bladder flaccid: a little yellow pale fluid.	Effusion into mesentery.	Engineer—died in 10 days.
7.	Slight partial ulceration of cæcum. General ulceration of lower portion of large intestine.	Healthy.	Liver large and pale. Gall-bladder distended.	Townsmen—died in 15 days after 1st attack.
8.	Ulceration of large intestines.	A hepatic abscess.	European Seaman—whole illness 12 days.

No.	State of large Intestines.	State of small Intestines.	State of Liver and Gall-bladder.	State of Mesentery, Mesenteric glands or other organs.	REMARKS.
9.	Much ulceration of large intestines.	Liver enormously enlarged.	European from house of correction—died in 31 days.
10.	Cæcum and transverse colon healthy: mucous coat of the descending down to rectum, ulcerated and sloughing.	Seaman—died in 21 days.
11.	Much ulceration at cæcum and sigmoid flexure, coats thickened and easily torn.	Liver much enlarged: abscess in right lobe. Gall-bladder half full of greenish thin bile, ducts pervious.	Seaman—died in 13 days.
12.	Cæcum in state of mortification, transverse coat and sigmoid flexure covered with minute ulcers.	Liver large. Gall-bladder full of dark bile.	Seaman—died in 22 days.
13.	Large intestine congested, mucous membrane sloughing: large coagulum in colon.	Liver large and gorged.	Mesenteric glands enlarged and indurated.	Seaman—ill a few days: doing well—suddenly passed a quantity of blood and died.
14.	Mucous surface of whole large intestines sloughing.	Liver large and mottled. Gall-bladder full.	Boy of 14—ill three weeks.
15.	Cæcum in sloughy state, distended with large coagulum of blood: colon inflamed and ulcerated.	Liver enlarged, pale—abscess in posterior portion of right lobe.	Seaman from house of correction—ill say a fortnight—died suddenly after passing a quantity of blood.

<i>No.</i>	<i>State of large Intestines.</i>	<i>State of small Intestines.</i>	<i>State of Liver and Gall-bladder.</i>	<i>State of Mesentery, Mesenteric glands or other organs.</i>	REMARKS.
16.	Extensive ulceration of cœcum and throughout colon.	Distended.	Seaman—died in a fortnight.
17.	Ditto.	Small abscess, size of a walnut, on lower surface of liver.	Seaman—died in 3 weeks.
18.	Ditto, and cœcum perforated.	Seaman, intemperate—died in 8 days.
19.	General ulceration of cœcum and colon.	Sailor—ill nearly 6 weeks.
20.	Old disease of cœcum—ulceration of whole colon and rectum.	Healthy.	Healthy. ..	Healthy. ..	Old Seaman : only 14 days' illness.
21.	Whole large intestine, especially cœcum, in sloughing state.	Cook—ill some weeks.
22.	Do. do., with adhesions to peritoneum.	African Seaman—died in 18 days.
23.	Large intestine inflamed and ulcerated throughout. Cœcum perforated in various places, containing pus.	Healthy.	Liver much enlarged.	Sailor, got dysentery in hospital—died in a fortnight.
24.	Sloughing of valve of cœcum. Colon thickened, ulcerated and cartilaginous.	Inflammatory blush on internal surface of stomach : small intestines little affected.	Seaman—had recovered from cholera—ill 10 days.

No.	State of large Intestines.	State of small Intestines.	State of Liver and Gall-bladder.	State of Mesentery, Mesenteric glands or other organs.	REMARKS.
25.	Ulceration of cœcum and transverse colon.	Small abscess.	Old Seaman—10 days in hospital.
26.	Cœcum and colon ulcerated throughout.	Small intestines here and there ulcerated.	Abscess, size of walnut, in right lobe.	Seaman—27 days ill.
27.	Colon inflamed and ulcerated throughout: in some places perforated.	Healthy.	Liver rather pale.	Had been treated for same complaint in previous month—died in 9 days: passed large quantity of blood.
28.	Whole intestine in right iliac region matted together. Cœcum ruptured.	Liver much enlarged: 3 abscesses, one in right lobe containing 3½j of pus.	African—ill 5 weeks.
29.	Cœcum thickened and sloughing: lower intestines ulcerated but less so.	Healthy.	Healthy. Gall-bladder distended.	Inflammation of peritoneum and tying down of omentum.	Seaman—died in 17 days.
30.	Intestines dark outside; internally covered with dark grumous blood: when wiped off, mucous membrane pale and bloodless. Cœcum thickened and ulcerated.	White and bloodless.	Liver enlarged and pale.	Seaman—died in 28 days: stools bloody.
31.	Cœcum disorganized and ulceration of colon.	Seaman—died in 20 days.

No.	State of large Intestines.	State of small Intestines.	State of Liver and Gall-bladder.	State of Mesentery, Mesenteric glands or other organs.	REMARKS.
32.	Large intestine ulcerated and sloughy throughout.	Small intestines in several places inflamed.	Liver somewhat large.	Seaman—died in 22 days: had diarrhoea before.
33.	Ulceration of large intestines, especially towards rectum.	Congested.	Seaman—died in 10 days.
34.	Cæcum semi-cartilaginous, perforated and full of pus, colon dark and ulcerated throughout.	Seaman—died in 9 days: lately had cholera.
35.	General ulceration throughout large intestine.	Seaman, ætat 36—died in 10 days.
36.	Cæcum and colon thickened, ulcerated, sloughing.	An immense abscess of liver.	Young Seaman—died in 26 days.
37.	Large intestines thickened and ulcerated throughout.	Healthy.	Large and soft.	Ætat 54—died in 19 days.
38.	Large intestine ulcerated throughout, chiefly in arch of colon.	Abscess of right kidney.	Recruit—died in 17 days.
39.	Great thickening of large intestine, with occasional spots of ecchymosis.	Liver large, pale and hard.	Kidneys pale and soft: some effusion at neck of bladder.	Seaman, age 18: ill about a month: passed bloody urine.
40.	Ulceration and sloughing state of cæcum extending along the colon.	Midshipman, ætat 14—died in 20 days.
41.	Cæcum and colon thickened and ulcerated throughout.	Blush on small intestines.	Mesentery injected, mesenteric glands enlarged.	Seaman, ætat 23—died in 20 days.

No.	State of large Intestines.	State of small Intestines.	State of Liver and Gall-bladder.	State of Mesentery, Mesenteric glands or other organs.	REMARKS.
42.	Cæcum much ulcerated.	Young European woman, much retching of coffee ground substance, and blood passed by stool.
43.	Large intestine, especially cæcum, a mass of ulceration. Ulcers varying from size of half crown to sixpence.	Liver large: gall-bladder empty.	Spleen large.	Recruit—died in 13 days.
44.	Large intestine thickened, ulcerated, and indurated throughout.	Stomach little affected.	Liver large, gall-bladder distended, with tarry bile.	Glands enlarged: spleen much enlarged.	Drummer: ætat 15—died in 22 days.
45.	Large intestines ulcerated in parts throughout.	Five small abscesses in liver.	Young Seaman—died in 34 days.
46.	Cæcum and colon greatly ulcerated—arch and descending colon little so.	Seaman, ætat 22—died in 29 days.
47.	Ulceration throughout the large intestines, and perforation.	Gall-bladder adhering to large intestine.	Seaman—died in 3 weeks.
48.	Large intestines thickened, ulcerated and covered with curdy matter.	Lower half of ileum livid externally, vascular internally.	Liver pale and enlarged.	West Indian—died in 12 days.
49.	Cæcum and neighbouring parts one mass of corruption: perforated.	Healthy.	Healthy.	Seaman—died in 3 months.

No.	State of large Intestines.	State of small Intestines.	State of Liver and Gall-bladder.	State of Mesentery, Mesenteric glands or other organs.	REMARKS.
50.	Cæcum and the rest of large intestine one mass of disease.	Healthy.	Young man : ill 15 days.
51.	Large intestines immensely thickened, ulcerated and perforated.	Healthy.	Healthy.	Recruit—died in 13 days.
52.	Cæcum and colon ulcerated throughout, and a pint of muddy fluid in abdomen from perforation of cæcum.	Liver very large, pushing up the diaphragm.	Glands hard and matted together.	Ætat 34—ill nearly two months.
53.	Large intestine ulcerated throughout.	Seaman—died in a fortnight.
54.	Large intestines ulcerated with fungous excrescences and lined with grumous matter throughout.	Small intestines discolored and vascular.	Mesentery discolored and vascular.	Female—died in 8 days : had phrenitis and vomiting : stools tarry.
55.	Scirrhus thickening of transverse and descending colon and rectum. Calibre of rectum would scarcely admit a finger : whole surface studded with fungous growths.	Old Soldier—died in 21 days.
56.	Large intestines ulcerated throughout in patches of irregular shape : contained bloody fluid and mucus.	Healthy.	Liver pale. Gall-bladder distended.	Recruit : ill about 40 days.

No.	State of large Intestines.	State of small Intestines.	State of Liver and Gall-bladder.	State of Mesentery, Mesenteric glands or other organs.	REMARKS.
57.	The whole large intestines ulcerated and sloughing, contracted in parts.	Liver natural in size, with cicatrices on surface : gall-bladder with healthy bile.	Glands enlarged.	Young stout man—died in 2 months.
58.	Whole large intestines ulcerated and sloughing : ileo-colic valve ulcerated.	Liver healthy. Gall-bladder full.	Ill 15 days.
59.	Large intestines one mass of disease.	Liver large, reddish flush on convex surface : gall bladder not distended.	Ætat 26—died in 10 days.
60.	Whole disease of intestines slight, except at cœcum.	Liver large, soft. Gall-bladder distended : duct pervious.	Mesentery injected.	Ætat 38—died in 12 days.
61.	Large intestines greatly altered, in some places very thin : chief seat of disease sigmoid flexure. Folds of transverse colon ulcerated.	Small intestines healthy.	A large and a small abscess of left lobe : small abscess of right lobe.	Meso-colon injected.	Pauper, ætat 30—died in 21 days.
62.	Cœcum and large intestines ulcerated throughout.	Ætat 14—died in 36 days.
63.	Large intestines one mass of disease.	Ætat 20—died in 4 weeks.
64.	Do. cœcum perforated.	Ætat 20,—ill 1 month.
65.	Large intestines diseased throughout, especially cœcum.	Small intestines generally healthy.	Liver, soft : a large and a small abscess in left lobe : gall-bladder half full.	Glands enlarged.	Seaman, ætat 32—died in a month.

No.	State of large Intestines.	State of small Intestines.	State of Liver and Gall-bladder.	State of Mesentery, Mesenteric glands or other organs.	REMARKS.
66.	Effusion of coagulable lymph on upper and outer surface : ulceration of ileo-colic valve. Large intestines diseased throughout : appearance here and there of a deposit of degenerated tubercular matter in the sub-mucous tissue.	Omentum tied down to small intestines.	Liver healthy : effusion of coagulable lymph on lower surface.	Old Seaman—died in 21 days.
67.	Generally ulcerated, especially the rectum.	Liver large with marks of old disease.	Glands enlarged.	Seaman, ætat 35—died in a month.
68.	Large intestines ulcerated throughout.	Seaman, ætat 27—ill a fortnight.
69.	Ditto.	Seaman, ætat 25—ill 1 month.
70.	Ditto. Sigmoid flexure perforated.	Liver healthy.	Glands enlarged.	Chief officer—died in 3 weeks.
71.	Highly diseased state : bowels breaking down under the fingers.	Seaman—died in 6 days : motions nearly pure blood.
72.	Generally ulcerated.	Small intestines healthy.	Liver large, hard and mottled.	Glands very large and hard.	Boy—died in 5 weeks : tubercles in lungs. Was a hard liver.
73.	Villous coat of large intestine red, and getting of deeper colour downwards : small ulcers at rectum of irregularly circular form, and rectum greatly thickened. Disease almost confined to rectum.	Healthy.	Healthy. ..	Healthy. ..	Seaman, ætat 33—died in 20 days.

No.	State of large Intestines.	State of small Intestines.	State of Liver and Gall-bladder.	State of Mesentery, Mesenteric glands or other organs.	REMARKS.
74.	Thickened, indurated and sloughing.	Pale and soft, slightly granular.	Pensioner, æt. 54—died in 18 days.
75.	Diseased throughout, with much ulceration, but chiefly at rectum, where much thickened: ulceration least in cœcum: villous coat destroyed and surface rough.	1½ foot of ileum had villous coat injected.	Liver healthy, adhesion to colon. Gall-bladder small, lined outside with rough false membrane.	Ill 14 days.
76.	Thickened and ulcerated, also covered with sloughs and coagula: latter chiefly in cœcum.	Pale and bloodless.	Liver pale and enlarged. Gall-bladder inordinately full of dark bile.	Seaman, æt. 37—died in 4 or 5 days: passed much blood.
77.	Irregularly shaped ulcers throughout, especially at the sigmoid flexure: intestine much thickened, contained a great quantity of bloody fluid, and at the cœcum was friable.	Healthy.	Healthy.	Seaman, in hospital 3 days; ill 3 weeks on board ship. Immense hemorrhage.
78.	Ulceration of rectum and also of cœcum, rest comparatively free from it: colon of dull purple colour.	For a few inches dull vascularity.	Liver somewhat larger than usual: five abscesses, one had about 3 ozs. of curdy pus.	Seaman, ill 3 weeks, in hospital 2 days: immense stools of blood.
79.	Ulcers occupying nearly whole calibre of intestine: worst at cœcum, right and left turns of intestine and sigmoid flexure, entirely destroying the villous coat, and in some places, the muscular, and leaving only the peritoneal, which gave way on being handled.	Healthy.	Healthy. ..	Omentum of dull pink colour, and adhering to cœcum.	Stout man, ill a fortnight, 8 days in hospital. Bloody motions.

No.	State of large Intestines.	State of small Intestines.	State of Liver and Gall-bladder.	State of Mesentery, Mesenteric glands or other organs.	REMARKS.
80.	Large intestines ulcerated and thickened throughout, chiefly in transverse colon.	Small abscess in left lobe.	Spleen enlarged.	Seaman, ætat 25—died in 10 days.
81.	Ulcerated throughout.	Liver large, soft and pale.	Omentum inflamed.	Seaman, ætat 18—died in 20 days. Bloody stools.
82.	Suppurating ulcers chiefly at arch of colon.	Liver pale and granular.	Seaman, ætat 16—died in 3 weeks.
83.	Cæcum and descending colon thickened and ulcerated: rectum in state of slough.	Slight adhesions between the convolutions of small intestine.	Liver pale. ..	Omentum vascular.	Seaman—died in a fortnight.
84.	Large intestine gangrenous and ulcerated throughout: whole one sloughy mass: every where covered with ash coloured sloughs.	Healthy.	Attack succeeded jungle fever—died in 5 weeks.
85.	Cæcum and colon, as far as sigmoid, thickened and cartilaginous—internal surface one continuous sloughing ulcer.	Liver very large and mottled: rather pale.	8 days in hospital—ill some weeks.
86.	Colon dilated to a diameter of 4 or 5 inches: cæcum a mass of ulceration and thickening, but no ulceration of rectum.	Liver small, pale, yellow, dry, and granular.	Old Seaman, ill 10 days: 3 in hospital.
87.	Cæcum sloughing, intestines, ulcerated and cartilaginous throughout.	Healthy.	Seaman, ætat 29—died in 5 or 6 days.

No.	State of large Intestines.	State of small Intestines.	State of Liver and Gall-bladder.	State of Mesentery, Mesenteric glands or other organs.	REMARKS.
88.	Large intestines in state of slough throughout.	Surface of liver shewed recent cicatrices: extensive abscess of right lobe.	Seaman, ætat 26—ill a month.
89.	Large intestines lined with ulcers, some of them apparently healing: coats perforated in several places.	Liver hard and granular, with some tubercles. Gall-duct obstructed and gall-bladder full.	Lungs tubercular: some effusion into chest.	Old man,—ill for one month before death.
90.	Intestines at most depending point in state of slough: rectum thickened and ulcerated.	Liver small and turgid.	Spleen slightly enlarged.	Ætat 67—died in 12 days: had immense hernia.
91.	Large intestines thickened, ulcerated and lined with sloughs throughout.	Liver small and softened: small abscess.	Seaman, ætat 30—died in 17 days.
92.	Large intestines lined with fungous growth, thickened, especially near rectum.	Liver pale, granular and soft.	Soldier, ill about 10 days.
93.	Colon, purple spots outside—inside mass of ulceration and thickening with lichenoid granulations.	Healthy.	Healthy.	Seaman—died in 8 days, but had just been discharged from hospital cured of dysentery.
94.	Mass of ulceration and thickening of cæcum, gave way on being handled—in sigmoid flexure and rectum less disease.	Healthy.	Liver one small abscess, size of nutmeg, otherwise healthy.	Ill 20 days—ætat 54.

No.	State of large Intestines.	State of small Intestines.	State of Liver and Gall-bladder.	State of Mesentery, Mesenteric glands or other organs.	REMARKS.
95.	Universal ulceration of large intestine, which broke down under fingers.	Healthy.	Liver enlarged: nutmeg.	Old pensioner: ætat 46, ill for 32 days.
96.	Cæcum, ascending and transverse colon thickened, cartilaginous and ulcerated.	Liver somewhat soft.	Townsmen, ætat 33, ill 3 weeks.
97.	Cæcum and colon thickened: full of suppurating ulcers.	Intestines floating in yellow serum.	Liver granular and dark: considerable abscess in left lobe.	Pauper—ill 12 days.
98.	Ditto...	Liver large and soft: left lobe pale.	Seaman, ætat 33, ill for 18 days.
99.	Cæcum and colon tough, and ulcerated throughout, rectum on the point of sloughing en masse.	Liver hard, granular, and almost white.	Pensioner, ætat 50—died in a fortnight.
100.	Large intestine mass of phagedænic ulcers.	Liver large and pale.	Townsmen, ætat 24—ill say 3 weeks.
101.	Large intestine ulcerated throughout: colon almost in state of decomposition.	Healthy.	Boatswain, ætat 45—ill 15 days.
102.	Ditto ditto, lower intestines less affected: no coagula in bowels.	Healthy.	Seaman, ætat 31—died in 15 days. Immense quantity of blood in stools.
103.	Ditto ditto, ulcerated throughout.	Liver pale and small.	Spleen large.	Stout: ætat 42—ill 17 days.
104.	Cæcum and ascending colon a mass of sloughing ulceration.	Ditto.	Ætat 31—ill 12 days.

No.	State of large Intestines.	State of small Intestines.	State of Liver and Gall-bladder.	State of Mesentery, Mesenteric glands or other organs.	REMARKS.
105.	Colon and whole of the large intestine full of cicatrised ulcers: very few in a state of active ulceration.	Small intestines in many places injected and red: slight abrasion of mucous surface.	Healthy. ..	Spleen and pancreas healthy.	Seaman, ætat 17—died in 31 days.
106.	A good deal of detached ulceration along the colon, but no great extent of it: coats of intestines not thickened.	Healthy.	Mottled.	Seaman, ætat 42, ill 3 weeks—stools chiefly of bloody washings.
107.	Cæcum much thickened and ulcerated, with imperfect granulations: colon, few superficial ulcers.	Small intestines and stomach blanched: patches of red on inner side of stomach.	Liver a little enlarged and full of abscesses, from size of pea to that of nutmeg, especially on upper surface of right lobe.	Recruit, ætat 23, ill at Dum-Dum—died 5 days after admission.
108.	Large intestines enormously ulcerated throughout and filled with coagula.	Liver dark, mottled and granular.	Invalid Artillery man—ætat 21, ill for 11 days.
109.	Extensive and deep ulcers throughout cæcum and colon—transverse and descending colon somewhat contracted.	Healthy.	Healthy. ..	Many glands enlarged.	Soldier, ætat 39: ill six weeks—stools bloody.
110.	Large intestines thickened and ulcerated throughout—sloughy.	Several patches of inflammation in small intestines.	Liver large and scirrhus, and adhering to diaphragm.	Spleen and kidney healthy.	Ætat 27, ill six weeks.

No.	State of large Intestines.	State of small Intestines.	State of Liver and Gall-bladder.	State of Mesentery, Mesenteric glands or other organs.	REMARKS.
111.	No perceptible lesion.	Intestine near the cæcum a mere web, and destitute of mucous membrane.	Liver, contained an immense abscess.	Ætat 18, ill 45 days: passed 12 or 14 inches of thickened mucous membrane 3 days before death.
112.	Arch of colon a mass of ulceration.	Seaman, ætat 24—ill 15 days: stools dark putrid blood.
113.	Patches of inflammation in colon and incipient ulceration.	Healthy.	Healthy. ..	2 pints serum in abdominal cavity.	Pensioner, ætat 40—died in 12 days: pulse hard.
114.	Large intestines much attenuated: cæcum and ascending colon exhibited large black patches ulcerating.	Small, soft, flabby and pale.	Mesentery very vascular: spleen healthy.	H. C. M. ætat 19—57 days ill: recovered but got relapse after fever chronic.
115.	Superficial ulceration throughout whole extent, not much thickening, ulcers like abrasions.	Healthy.	Healthy.	Ætat 28—struck by lightning: ill a fortnight: doing well: got attack of cholera—recovered. Dysentery returned—died in 40 days.
116.	Externally congested: internally of reddish hue throughout, thickly studded with small red spots: no ulceration or thickening but sort of superficial abrasion.	Small intestines vascular, inner surface in many places congested.	Healthy.	Midshipman, ætat 17—died in 7 days.

No.	State of large Intestines.	State of small Intestines.	State of Liver and Gall-bladder.	State of Mesentery, Mesenteric glands or other organs.	REMARKS.
117.	Mass of corruption breaking down under the fingers.	Attacked in hospital—died in 7 days—ætat 45.
118.	Large intestines a good deal thickened and ulcerated, especially inferiorly, but to no very great extent.	Healthy.	Liver small.	Seaman, ætat 30—died in 35 days.
119.	Intestine thickened and ulcerated throughout, lined with mucous and bloody coagula.	Liver rather large and pale.	Seaman, ætat 18—died in 34 days.
120.	Rectum and sigmoid flexure in state of irritable ulceration.	Some fluid effused in pelvis.	Soldier, ætat 21—died in 5 days, complicated with scorbutus and fever.
121.	Cæcum and sigmoid flexure thickened, ulcerated and breaking under fingers.	Small intestines slightly glued together, lower portion filled with dark brown fluid.	Liver rather large and soft.	Spleen large.	Soldier, ætat 25—died in 17 days: stools pure blood, mixed with mucus.
122.	Upper part of colon and cæcum in state of sphacelus.	Lower part of ileum in state of sphacelus.	Liver enlarged, granular, grey degeneration.	Pensioner, ætat 44—ill 40 days.
123.	Cæcum distended with clots of blood, much effused blood along the whole course of colon.	Rather blanched but quite healthy internally.	Nutmeg liver.	French Sailor—ætat 24: 11 days in hospital.

<i>No.</i>	<i>State of large Intestines.</i>	<i>State of small Intestines.</i>	<i>State of Liver and Gall-bladder.</i>	<i>State of Mesentery, Mesenteric glands or other organs.</i>	<i>REMARKS.</i>
124.	Ulcerated and thickened throughout caput coli: cœcum studded with large unorganized projections from ulcerated base: ulcers in every stage: whole rectum raw and abraded. Appendix vermiformis very long, full of sanious fluid.	Seaman, ætat 24—died in 3 weeks.
125.	Large sloughy ulcers in cœcum, less in transverse arch, but bowel in sloughy state throughout.	Small intestines inflamed and partially agglutinated together.	Liver granular, more like lung than liver. Gall-bladder small, with inspissated bile.	Mesentery much inflamed.	Seaman, ætat 24—died in 14 days: 2 days in hospital.
126.	Cœcum and transverse colon ulcerated and sloughy.	Ileum inflamed.	Liver gorged.	Mesentery and meso-colon inflamed: right iliac fossa a bath of blood and inflammation.	Seaman, ætat 27—died in 11 days.
127.	Large intestines mass of ulceration and sloughing.	Stomach healthy: small intestines in places discoloured.	Liver nutmeg.	Mesentery healthy: spleen healthy.	Ætat 24—died in 20 days.
128.	Externally dark, internally purple, with more or less abrasion, scarcely amounting to ulceration: transverse colon and rectum much affected: no clots of blood.	Small intestines dark and congested: some superficial abrasion.	Veins of mesentery turgid, with dark blood.	Seaman, ætat 19—died in 18 days: motions bloody.

No.	State of large Intestines.	State of small Intestines.	State of Liver and Gall-bladder.	State of Mesentery, Mesenteric glands or other organs.	REMARKS.
129.	Colon much thickened and ulcerated throughout.	Mucous coat of stomach red and abraded.	Healthy. ..	Healthy. ..	Seaman, ætat 41—ill 8 days—in hospital 6 hours.
130.	Large intestines purple, with ulcerations throughout, breaking down under the fingers.	Healthy.	Healthy. ..	Healthy. ..	Ætat 31—died in 11 days.
131.	Extensive ulceration of large intestine, especially cæcum.	Red blush externally, but internal coat healthy.	Healthy.	Seaman, ætat 22—died in 26 days.
132.	Thickened and ulcerated throughout with black spots as from hæmorrhage.	Healthy.	Healthy.	Ætat 38—died in 3 weeks : passed much blood.
133.	Large intestines ulcerated throughout, towards rectum frequent bloody patches, and the last part of the gut sloughing.	Healthy.	Healthy.	Soldier, ætat 24—died in 3 weeks : rectum especially affected.
134.	Externally dark purple, ulcerated throughout.	Small intestines dark outside : mucous membrane dark red, inclined to ulcerate.	Healthy.	Seaman, ætat 22—died in 16 days : small intestines more than usually affected.
135.	Cæcum and rectum ulcerated, former lined with coagula.	Spleen large.	Ætat 26—died in a month.

No.	State of large Intestines.	State of small Intestines.	State of Liver and Gall-bladder.	State of Mesentery, Mesenteric glands or other organs.	REMARKS.
136.	Cæcum distended.	Old man—died in three weeks.
137.	Large intestines in sloughy state.	Granular.	Ill about a month, ætat 32.
138.	Cæcum and colon lined with slough.	Small, yellow and granular.	Pensioner, ætat 44, ill 18 days.
139.	In cæcum much ulceration, less in ascending colon: transverse free from ulcers, a few in sigmoid: ulcers looked indolent: villous coat puckered round them as if going to cicatrise.	Nothing abnormal.	Liver enlarged, adhering throughout to diaphragm: structure loose and friable: abscess containing $4\frac{3}{4}$ of sanious pus. Gall-bladder full of bile.	Ætat 36: dysentery came on 5 weeks before, after fever.
140.	Large intestine one continued ulcer.	Liver large, pale and dry.	Ætat 26—died in 7 weeks.
141.	Large intestines ditto.	Small intestines in parts of dark purple colour.	Rather large but healthy.	Æt. 27, about 40 days. Odd nervous symptoms.
142.	Cæcum thickened, cartilaginous and studded with small perforating ulcers.	6 inches of small intestines thickened, &c.	Pale, yellow, large, hard and granular.	Invalid: ill 16 days.
143.	Ulceration throughout large intestines.	Liver pale, yellow and granular.	Died in three weeks.
144.	Large intestine ulcerated, but to no great extent.	Healthy.	Slightly hardened: right lobe had 2 small abscesses: one healthy, one of sanious pus.	Pensioner, ætat 41—died in a month.

No.	State of large Intestines.	State of small Intestines.	State of Liver and Gall-bladder.	State of Mesentery, Mesenteric glands or other organs.	REMARKS.
145.	Ulceration of large intestines throughout.	Healthy	Healthy.	Engineer, ætat 28—ill a month.
146.	Ditto.	Ditto. ..	Ditto.	Pauper, ætat 28, ditto.
147.	Large intestines thickened and ulcerated throughout.	Healthy.	Healthy.	Seaman, ætat 26—died in 11 days: stools like coffee grounds.
148.	Considerable ulceration of large intestine: mucous membrane in places destroyed.	Ditto. ..	Ditto.	Ætat 24, ill about 14 days.
149.	Thickening and ulceration of large intestine, and especially rectum, which broke down on handling.	Ditto. ..	Ditto. ..	Vascularity and enlargement of glands.	Midshipman, ætat 16, ill 12 days.
150.	Ulceration throughout whole extent: whole surface rough and covered with whitish-yellow lichenoid substance, which could scarcely be detached.	Ditto.	Pauper, ætat 43, recovered from jungle fever: ill a month.
151.	Extensive ulceration: coats thickened and friable.	Soldier, ætat 26—died in 3 weeks.
152.	Thickened throughout with numerous small ulcers.	Inner coat of stomach injected.	Soldier, ætat 20—died in 10 days.

No.	State of large Intestines.	State of small Intestines.	State of Liver and Gall-bladder.	State of Mesentery, Mesenteric glands or other organs.	REMARKS.
153.	General oozing of blood from mucous surface throughout its whole extent; collections of blood in rectum and cœcum, latter had a few deep ulcers, no general ulceration.	Healthy.	Healthy.	Ætat 25—died in 16 days from active hæmorrhage : bled from nose and lungs.
154.	Ulcerated throughout : cœcum in state of sphacelus.	Ditto. ..	Ditto.	Soldier, ætat 28, ill 20 days.
155.	Generally ulcerated.	Pale.	Woman, ætat 38—died in 5 weeks.
156.	Ulcerations chiefly of cœcum and rectum : ulcers distinct, separate and large, not much thickening.	Healthy : one or two yellow discoloration on upper surface.	Seaman, ætat 22—died in 15 days.
157.	Much ulcerated and thickened throughout.	Liver yellow.	Midshipman, ætat 18, ill 3 weeks.
158.	One mass of thickening and ulceration throughout.	Rather pale.	Ætat 15, ill 30 to 40 days.
159.	Large intestines one mass of ulceration.	Small intestines injected, especially cœcum.	Liver with many cicatrices.	Seaman, ætat 40—died in 17 days.
160.	One mass of disease: mucous membrane swollen and gangrenous, with black patches of slough.	Healthy.	Healthy.	Soldier, ætat 22, ill 18 days.

SUMMARY OF TABLE OF ACUTE DYSENTERY.

Thus in 160 cases of Acute Dysentery—

The liver is found to be altered in *	84
to contain abscess in	21
is enlarged in	40
is gorged or turgid in	4
is small in	7
is pale in	26
is granular or nutmeg in	22
is soft in	12
is hard in	5
contains cicatrices* in	3

The gall-bladder appears to be almost always full, and to contain healthy or somewhat inspissated bile.

The ileum is noted as over-vascular or congested in 21

Slight ulceration and abrasion are seen in 3

It is in a state of sphacelus in 1

The stomach has its mucous coat over-vascular or

 somewhat softened in 4

 and ulcerated in 1

The large intestine is ulcerated in all, and chiefly at the cœcum, sigmoid flexure, and rectum.

The cœcum, transverse and descending colon were

 free from ulceration in 3

It was perforated (generally in the cœcum) in .. 8

There was ulcerative destruction of ileo-colic valve in 3

Suppuration of appendix vermiformis (though not uncommon) is only recorded in 1

There is thickening and stricture of the intestine in 4

 and dilatation in 1

The mesenteric glands are enlarged or inflamed in. . 17

The spleen is enlarged in 6

The kidneys are diseased in 2

* These cicatrices do not appear to be the sequelæ of abscesses.

CHRONIC DYSENTERY RUNNING INTO DIARRHŒA.

No.	Large Intestines.	Stomach and small Intestines.	Liver and Gall-bladder.		REMARKS.
1.	Extensive ulceration of sigmoid flexure.	Large abscess of liver.	Ill 2½ months.
2.	Colon in a sloughy state : coats gave way on handling.	Long suffered from dysentery.
3.	Lining of colon ulcerated throughout, greatest lesion at sigmoid flexure and rectum, where ulcers were almost continuous.	Liver enlarged, weighing 6½ lbs. attached to spleen by false membrane, substance soft : colour pale. Gall-bladder distended with green bile.	Spleen enlarged, indurated : mesenteric glands enlarged.	Invalid—died in 10 days of acute dysentery : supervening on chronic.
4.	Extensive ulceration of colon, orifice in sigmoid flexure, through which contents of bowels extravasated.	Slight attempt at inflammation of peritoneal surface.	Woman of 55—ill for six months.
5.	Intestines attenuated, several ulcers in cæcum and along arch of colon to sigmoid flexure. Ulcers in an atonic state : no raised edges, no increased vascularity.	Liver small and unhealthy.	Congestion of mesentery : some enlargement of mesenteric glands.	Soldier, ill for 7 weeks out of and 115 days in hospital.
6.	Large intestines ulcerated throughout ; transverse colon thinner than natural : mucous membrane, had honeycomb appearance.	Stomach distended with air.	Liver dark, granular, with some congestion. Peritoneal coat puckered in various places : gall-bladder distended : ducts pervious.	Soldier, admitted bed-ridden—died in 18 days.

No.	Large Intestines.	Stomach and small Intestines.	Liver and Gall-bladder.		REMARKS.
7.	Large intestines thickened and ulcerated, lined with pus.	Healthy.	3 abscesses in right lobe full of curdy matter.	Pauper, ill several months—died in six days.
8.	Contraction of colon, general superficial ulceration of bowel: deeper of rectum.	Mucous surface of stomach abraded towards pylorus and an inch down the duodenum.	Mesenteric glands enlarged.	Invalided for chronic dysentery—died after two months' treatment.
9.	Large intestine thickened and ulcerated throughout, cœcum nearly obliterated.	Cicatrized fissure on upper surface, from 4 to 5 inches in length $4\frac{1}{2}$ an inch deep. A large hydatid firmly attached to left lobe.	Old Soldier, from Cabul.
10.	Large intestines contracted in several parts, ulcerated in patches of circular and irregular form from cœcum to rectum, which was in a state of slough.	Liver healthy.	Abscess in cellular tissue between cœcum and psoas, which was dissected by it.	Portugese Doctor, ill 8 months.
11.	Large intestines ulcerated throughout.	Pale and enlarged.	Invalided for bowel complaint.
12.	Scirrhus thickening and ulceration throughout, but chiefly of rectum.	Liver pale and granular.	Pauper—ill 6 or 8 months.

No.	Large Intestines.	Stomach and small Intestines.	Liver and Gall-bladder.		REMARKS.
13.	Whole cœcum and descending colon in sloughy state, transverse arch not much affected: rectum cartilaginous.	Liver healthy. Gall-bladder empty.	Glands enlarged.	Pensioner, a long resident in India, had been in China : ill many months.
14.	Large intestine in state of slough throughout.	Mate of Floating Light, æt. 24, ill 1 year.
15.	Cœcum rotten and perforated, transverse colon ulcerated, sigmoid flexure healthy.	Abscess of liver communicating with ascending colon.	Invalid, had been several weeks in hospital : discharged, but returned with fresh attack.
16.	Large intestines ulcerated and sloughing throughout.	Spleen greatly enlarged.	Young man, ill about 9 months.
17.	Cœcum thickened with villous coat, destroyed and like tanned leather : colon and rectum both much diseased.	Gall-bladder gorged.	Effusion in abdomen.	Ætat 26, ill one year.
18.	Large intestine contracted, thickened and cartilaginous with fungous granulations.	Liver pale and soft.	Recruit, ill for one year.
19.	Colon adhering to stomach.	Small intestines glued to abdominal parietes & torn into shreds on separation.	Pale and small.	Pensioner, ætat 40, ill 4½ months.
20.	Large intestines one mass of ulceration.	Healthy.	Pale and gritty.	Ætat 62, ill some months—female.

No.	Large Intestines.	Stomach and small Intestines.	Liver and Gall-bladder.		REMARKS.
21.	Large intestines lined with fungous granulations, and calibre of gut in places contracted to size of finger.	Much attenuated.	Yellow, hard and granular.	Seaman, ætat 40, ill three months.
22.	Hæmorrhage external to bowels, which were purple internally, with little ulceration.	Liver cirrlosed.	Mesentery and meso-colon full of effused blood: spleen greatly enlarged.	Ætat 44, broken down—drunkard, died suddenly on close stool.
23.	Large intestine ulcerated throughout.	Healthy.	Pale.	Soldier, ætat 25. Ill many months.
24.	Large intestines slightly ulcerated, some thickening of cæcum.	Healthy.	Liver double its natural size, like a coagulum of blood: nine small abscesses containing 3 or 4 ozs. of pus.	Invalid, ætat 31—many months ill.
25.	Large intestines with patches of rather superficial ulceration, and of dark red discolouration.	Healthy.	Healthy.	Ætat 27 : 75 days ill—complicated with secondary syphilis.
26.	Colon not much thickened, extensive patches of ulceration and spurious granulations.	Pale. ..	Pale.	Seaman, ætat 27—ill 5 months.
27.	Large intestine ulcerated at cæcum, effusion of grumous matter on its surface, and dotted with spots of ulceration throughout.	Healthy.	Ætat 34, ill 5 months.

No.	Large Intestines.	Stomach and small Intestines.	Liver and Gall-bladder.		REMARKS.
28.	Internal surface of colon ulcerated and abraded throughout: not much thickening of coats.	Healthy.	Healthy.	Ætat 33—ill about 3 months.
29.	Large intestines purple outside, inside quite gangrenous, with those lichenoid excrescences or abortive cicatrization?	Healthy.	Healthy.	Soldier, ætat 34, ill many months.
30.	Coats of large intestine extensively ulcerated, especially cœcum and rectum.	Mesentery vascular, glands enlarged.	French Sailor, ill 3 months.
31.	Extensive ulceration, coats black, and in state of sphacelus.	Ætat 26, ill 2 months.
32.	Much thickened, inner coat livid and ulcerated.	Coats of stomach vascular, and abrasion of coat of small intestines.	Mesentery injected, glands swollen, with some hydatids in them.	Pensioner, ætat 46, ill 8 months.
33.	Purple inside, with superficial ulceration throughout.	Immensely enlarged, granular, with white patches on upper surface: cicatrices.	Soldier, ætat 23, ill some months on voyage out.
34.	Thickened, with a few ulcers—and general vascularity of mucous coat.	Healthy.	Ætat 31, ill 3 months.
35.	Coats thin and friable, extensive ulceration throughout.	Enlarged, large abscess on lower surface of right lobe.	Glands swollen.	Ætat 22, ill 7 weeks.

No.	Large Intestines.	Stomach and small Intestines.	Liver and Gall-bladder.		REMARKS.
36.	General ulceration and thickening.	Coats of stomach vascular.	Liver full of abscesses, large one in left lobe. Gall-bladder unusually large.	Mesentery vascular, glands enlarged.	Ætat 30, ill 3 months.
37.	General ulceration and thickening, especially of cœcum and rectum.	Ætat 28, ill about 3 months.
38.	Large intestines lined with bloody tenacious matter.	Liver small and hard.	Pancreas small and hard, omentum and mesentery discoloured: latter nearly black: glands were rather enlarged.	Ætat 63, ill 2 or 3 months: passed 2 or 3 pints of blood every day.
39.	A good deal of ulceration of large intestine.	Stomach healthy, ulcer at the pyloric orifice, where there was cancerous ulceration.	Healthy.	Sailor, æt. 40, says he had been ill 3 months—died in 16 days.
40.	Patches of purple discolouration on inner coat.	Pale. ..	Early stage of nutmeg degeneration.	Seaman, ætat 40, ill some months—died with odd head symptoms.
41.	Attenuated, not ulcerated.	Mucous membrane of stomach softened throughout: towards pylorus abraded and rough.	Pale.	Has secondary syphilis.

<i>No.</i>	<i>Large Intestines.</i>	<i>Stomach and small intestines.</i>	<i>Liver and Gall-bladder.</i>		REMARKS.
42.	Internal surface of cæcum covered with a red efflorescence, some spots of ulceration in the colon. Descending colon and rectum contracted, thickened and cartilaginous.	Intestines generally glued together.	Liver natural, contents of gall-bladder pale.	Glands enlarged and livid.	Invalid, ill long time.
43.	Towards sigmoid flexure a few patches of ulceration.	Congested ecchymosed state of pyloric orifice, coat partially abraded—this extending to duodenum—congestion of ileum: bowels on whole softened, œdematous, and tearing readily.	Small particles like tubercular matter, on surface of liver.	Glands enlarged.	Ætat 36—ill some five weeks.
44.	Rectum in scirrhous state, with ulceration extending up to sigmoid; some patches of ulceration in transverse colon and ileo-colic valve.	Ulceration extending some inches up the ileum.	Liver hard and nutmeg.	Glands enlarged and diseased.	Old Seaman, 3 months ill.
45.	Spots of ecchymosis on descending colon.	Small intestine thickened, in lower portion some ecchymosis.	Liver small: flaccid gall-bladder.	Mesenteric glands enlarged and injected.	Diarrhœa after cholera—died in five weeks.

No.	Large Intestines.	Stomach and small Intestines.	Liver and Gall-bladder.		REMARKS.
46.	Large intestine pale externally, slight ulceration inside.	Pale. ..	Liver rather pale.	Mesenteric glands enlarged.	Invalid, ill many months.
47	Throughout thickened and cartilaginous, at the arch and sigmoid flexure surface irritable.	Stomach distended with bilious fluid : small intestine doughy.	Liver small and pale. Gall-bladder full of dark bile.	Mesenteric glands enlarged, spleen large.	Invalid : long ill.
48.	Large intestine pale, and so much contracted as scarcely to admit of the introduction of the blades of a pair of scissors : destitute of lubricating mucus, with red efflorescences towards rectum.	Small intestines extremely attenuated and transparent.	Dark.	Soldier, from China. Ætat 20—ill four and a half months.
49.	Considerable chronic inflammation of mucous coat of large intestine. Coats a good deal thickened : no deep ulceration.	Chronic inflammation of mucous coat of stomach and small intestines.	Healthy. Gall-bladder full.	Mesenteric glands atrophied : spleen and pancreas small.	Ætat 30—about six months ill.
50.	Some ulceration of large intestines.	Healthy.	Small dark surface, studded with spots of organized lymph.	Ætat 36—2 years' standing. Complicated with lungs.
51.	Colon and rectum contracted, dry and cartilaginous, external surface superficially ulcerated.	Soldier. ætat 43—ill 7 months.

No.	<i>Large Intestines.</i>	<i>Stomach and small Intestines.</i>	<i>Liver and Gall-bladder.</i>		REMARKS
52.	Large intestines pale and blanched, otherwise healthy.	Coats of stomach bloodless: small intestines pale.	Liver yellow, mottled or nutmeg.	Invalid, ætat 53, had been drinking—much bilious vomiting: ill 3 or 4 months.
53.	Large intestines congested and abraded throughout—no deep ulceration.	Small intestines congested: and mucous surface abraded partially.	Healthy.	Seaman, ætat 18—ill 3 months.
54.	Pale and bloodless.	Pale and bloodless.	Light yellow colour, slightly enlarged.	Half starved Seaman—ætat 43, ill 2 months.
55.	Pretty general ulceration of superficial kind.	Small and contracted from former inflammation.	East Indian, ætat 59, ill many months.

SUMMARY OF TABLE OF CHRONIC DYSENTERY.

In 55 cases of chronic dysentery—

the liver was altered in ..	31
abscess was found in ..	6
hydatids in	1
cirrhosis in	1
enlargement in	5
diminution of bulk in ..	8
nutmeg alteration in ..	6
it was pale in	11
it was hard in	4
soft in	1
contained cicatrices? in	1

The gall-bladder and its contents seem generally to have been healthy: the bile occasionally rather pale coloured, but at other times the reverse.

The large intestines were ulcerated in	50
the colon contracted in	3
cæcum nearly closed in	1
colon perforated in	1

The stomach is noted as unhealthy in 6

There was chronic inflammation and softening in ..	2
increased vascularity in	2
abrasion of pylorus in	3
cancer of pylorus in	1

The small intestines are noted as unhealthy in .. 12

Ulceration or abrasion of ileum in 3

Mesenteric glands enlarged in 16

The spleen was enlarged in 4

As no distinction can be drawn between acute and chronic dysentery, which is not, to a great degree, arbitrary, so chronic dysentery and diarrhoea are divided by no marked line. The last 15 cases belong rather to diarrhoea than dysentery, and it will be observed, that in them the large intestine is paler than natural, and in several instances not ulcerated, and that in no instance was there abscess, though the liver was found altered in 11 out of 15 cases.

CHUSAN DYSENTERY.

Although a great many patients died in the General Hospital from Chusan dysentery, these cases are excluded from the preceding tables,—1st, because they are instances of a form of malarious dysentery, which is very different from the common disease of Bengal,—And 2nd, because ulceration of the large intestine, of a true dysenteric character, though sometimes met with in China, seems chiefly to have supervened after the patients arrived in Bengal. But the mortality from it was so great, and it was altogether so unmanageable a form of disease, that the accompanying abstract of the pathological appearances in those who died soon after their arrival in Bengal is worthy of being preserved.

<i>Large Intestine.</i>	<i>Stomach and small Intestine.</i>	<i>Liver and Gall-bladder.</i>	<i>Mesentery, &c.</i>
Generally attenuated, now and then inflammatory blush: in further advanced stage some ulceration.	Attenuated: rosy hue of pyloric orifice, some roughness at commencement of duodenum: frequent abrasion and softening of mucous coat of jejunum and ileum, ulceration of glands of Peyer.	Liver healthy or pale: gall-bladder full of fluid bile, often flaccid: bile pale and limpid.	Mesenteric glands always enlarged, containing curdy, scrofulous like matter: spleen sometimes enlarged.

Chusan differed from Bengal dysentery chiefly in these respects: in its setting in with much fever, which here is very unusual; in the slight degree in which the liver and large intestine were affected; and the great amount of mesenteric disease and affection of the small intestines. In many respects it resembled Cabul dysentery, which the men called "the white flux," especially in the general absence of blood in the motions.

COMPARISON OF ACUTE AND CHRONIC DYSENTERY.

On a general review of the appearances presented in cases of acute and chronic dysentery, it would appear, that the liver is most frequently altered in the latter: that abscess is about equally frequent in either form—that in acute dysentery the liver is frequently enlarged and soft, while in chronic it is more generally small and indurated.

The stomach and small intestines also suffer more frequently in the chronic form, and the mesenteric glands are more frequently altered in it.

The extent of disease in the colon is least in the cases bordering on diarrhœa, and whereas in 160 cases of acute dysentery the large intestine is perforated 8 times, in chronic it was only once in 55.

Nothing very distinct can be said about the gall-bladder or its contents, but in the acute form it appears to be generally healthy, although often loaded with inspissated bile; while in the most chronic form bordering on diarrhœa the gall-bladder is often flaccid, and its contents pale and fluid.

The proportion of hæmorrhagic cases is smaller in the chronic than in the acute form.

Enlarged spleen is perhaps more common in the chronic form, but this is to be looked on as an accidental com-

plication. No occasional mention of the state of the pancreas or kidney is of value, as they are certainly not implicated in the disease.

TRUE DESCRIPTION OF THE CHANGES PRODUCED BY BENGAL DYSENTERY.

It is not my intention to enter into any minute description of the state of the intestines, which has been faithfully described by both Twining and Raleigh, nor am I able to throw any fresh light on the nature of the dysenteric process. It has been compared to erysipelas by Siebert, and to the corrosion of mineral acids by Cruveilhier and Rokitansky. The mechanical theory of the irritation of scybala or accumulation of fæces acting on an inflamed surface, though generally abandoned, still finds some supporters. It has by many been attributed to the irritation of altered biliary secretion, or to its absence. Parkes considers it to be a process of ulceration universally commencing in the solitary glands of the large intestine. Others, with Raleigh, consider it to be a simple inflammation of the mucous coat of the large intestine, (if it were simple it would be more amenable to treatment). Whatever of truth or error there may be in these opinions, the appearance presented to us in simple Bengal dysentery is, that of an inflammation of the large intestine, which may be diffusive, ulcerative, purulent, hæmorrhagic, or gangrenous, according to circumstances. The disease in Europe and in India is essentially the same, and the best scientific descriptions of Bengal dysentery are those given by Dr. Baly of London and Rokitansky of Vienna, although the latter has not met with the amount of ulceration, which is common here. As seen here, the process is very generally one of mortification and sloughing, not of simple ulceration, i. e. the ulceration is often secondary, and occurs only after the sloughs are thrown

off. Inflammation and ulceration of the solitary glands is very unusual, or has been very carelessly observed, and I believe it may be stated generally, that in Bengal dysentery, they are not peculiarly or primarily diseased. It should be borne in mind, that the state of the solitary glands, as observed by Murray and Parkes, exactly corresponds with their usual appearance in cholera, and that all Murray's and most of Parkes's cases occurred in dysenteric patients suddenly carried off by that disease.

FREQUENCY OF COMPLICATION WITH DISEASE OF LIVER.

Perhaps I may as well offer a few remarks on the frequency of the occurrence of disease of the liver in connexion with dysentery. Our information on this subject is very meagre, as few observers have recorded any change of structure in the liver except that of the formation of abscess.

On the continent of Europe, Broussais, Gueretin, Thomas, Siebert, and Rokitansky, some of whom have written at great length on the pathology of dysentery, do not, as far as I can learn from abstracts of their writings, allude to hepatic complications. In the miasmatic form of dysentery, abscess is very unusual: it was rarely observed in the Continental wars of the 18th century* and in the Peninsula—not at all in various Austrian epidemics or in the Penitentiary. In the Burmese war, Waddell says he never saw any structural lesion of the liver. In China, Dr. Wilson found abscess only twice in 61 fatal cases. The only exception to this general rule seems to be Ireland, where Dr. Cheyne found it four times in 30 cases; but again Dr. Mayne never met with it.

* Pringle however found abscess in 2 out of 5 cases, of which he gives dissections, but elsewhere says that the liver was generally healthy.

The following data are sufficient to confirm the opinion, that abscess of the liver is a much less frequent complication of dysentery in Calcutta, than in Bombay or Madras. The reason I am not prepared to explain: but assuming, what there is every reason to believe, that abscess co-existing with dysentery is rare in the miasmatic form, and common in the other—that it is more common in the other presidencies than in Bengal, (where dysentery itself prevails so extensively in its worst shape,) and that it may be common one season and rare the next, it naturally follows, that some far more general cause must determine the formation of abscess than the limited one, of the absorption of pus from the ulcerated surface of the colon, or from suppurating mesenteric glands, assigned by Dr. Budd.

In New Orleans Dr. Robertson found hepatic abscess the common cause of death in dysentery.

In Madras, Annesley	in 51 cases finds 26 abscesses.
Dr. Shanks	96 .. 36 „
Parker and Innes	61 .. 13 „
Ballingall	35 .. 4 „
At Bombay, Dr. Morehead	30 .. 12 „
At Calcutta, Seaman's Hospital*	24 .. 5 „
Medl. College Hospital*	54 .. 14 „
Genl. Hospital	215 .. 27 „

It is however worthy of remark, that the liver has been found in the General Hospital to have been altered in 118 out of 215 cases: in the Medical College Hospital in 13 out of 30 cases, while Sir James MacGrigor found it in India altered 16 times in 22 cases, “and in Egypt as in India found it diseased.” In the Peninsula the liver was found generally healthy, but sometimes indurated and softened and sometimes the seat of abscess. In Dublin Dr. Mayne says, it was always healthy, but sometimes congested.

* If these cases were minutely examined, I think that the number of cases of abscess would be somewhat less.

While on this subject, I may add, though not prepared to enter on the question of the connexion between liver disease and dysentery, that, on analysing 46 cases in the General Hospital, returned as hepatitis and terminating in abscess, the large intestine was found ulcerated in 15, and the small intestine in parts over-vascular, or with its mucous surface abraded in 4:—14 patients had been suffering for sometime from dysentery, and 41 had been ailing for a week or two before admission.

Simple acute inflammation terminating in abscess would therefore seem to be comparatively unusual in this part of India, though I am unacquainted with what is termed duodenal dyspepsia by Dr. Parkes, and which he considers to be the usual precursor of abscess.

I would also venture here to intimate a doubt, whether the occlusion of the duct from enlarged glands of the capsule of Glisson, or inflammation of the gall-bladder in new arrivals, as described by Twining, be at all of common occurrence.

ON CERTAIN APPEARANCES MENTIONED BY AUTHORS.

Ere concluding, I would offer an observation or two on various appearances which have been described by authors. Scybala or accumulations of fæces appear to have been scarcely ever observed, during life, and never after death.

No sloughs of complete portions of the intestinal canal have been remarked, but during life large, tough, tubular portions of mucous membrane and effused lymph have come away frequently, and no doubt they are occasionally found in situ.

Ulceration of the ileo-colic valve is probably more frequent than it is represented in these tables to be, but no such thing as intus-susception of the small into the large intestine has been witnessed.

The appearance described by MacGrigor, as fungoid or lichenoid excrescence of the colon, is common enough.

No suppuration of the mesenteric glands or occurrence of pus in the vena cava, have been met with.

Displacements of the large intestine are frequent, but have not been noticed with sufficient accuracy ; however they are of slight practical importance.

The same must be said of adhesions and matting of the omentum.

The *lumbricus teres* is often found, especially in patients who have come off a long voyage.

Cicatrization of ulcers, which Raleigh considers so unusual, is far from infrequent.

The pouring out of blood in immense quantities from the commencement of the disease, as described by Twining, and stated to be very rare at Bombay by Morehead, occurred in 19 out of 160 acute cases.

On the subject of spasm of the intestine, post mortem examination throws no light.

CONCLUSION.

Were it not my object to make these remarks as short as possible, I should like much, to append the history of some of the most interesting cases, for instance of two or three cases of what has been clumsily termed perityphlitis, or abscess in the cellular tissue round the cœcum, of a case carried off by cholera, just after the process of cicatrization, marked by a dark coloured deposit, had been completed, of various hæmorrhagic ones, especially an instance in which hæmorrhage from the bowels was preceded by epistaxis and hæmoptysis, and most of all, a fatal case complicated with odd tetanic symptoms, the only one I have met with, which corresponds at all with the nervous symptoms described as occurring in some of the cases at the Penitentiary.

NOTES.

On Treatment.

1. Having alluded only very cursorily to the subject of treatment, I may be allowed to add that, after greatly increased experience of the disease, the opinions expressed by me in former years in the following extracts, still appear to me in the main just. Possibly, the value of the use of sugar of lead in practice among Europeans may be over-rated, and others do not speak so well of it, but English authors have never placed so much confidence in it as American and German ones. After all, our practice in bad dysentery is very unsatisfactory, and quite a contrast to that in fever:—would that other diseases were as manageable as the latter! The approximate percentage of deaths in the General Hospital will shew, that except in bad seasons, and when cases are brought in late, the mortality from it is small.

1841.	1842.	1843.	1844.	1845.	1846.	1847.	1848.	1849.
6.09	5.2	3.8	2.1	4.	10.4	6.	6.2	2.9

“It cannot admit of doubt, that calomel and drastic purgatives have been injudiciously used, and that a return to a milder mode of treatment will be attended with the most beneficial results.

There is no difference of opinion as to the propriety of free depletion in the earlier stages of this disease, followed up by the use of mild purgatives, among which castor oil is quite invaluable. The combination of blue pill, ipecacuanha, gentian and hyoscyamus, so commonly employed, is a most useful preparation, and opium is also a very important remedy, although the belief that it merely masks the disease is very prevalent. An opiate enema, or Dover’s powder, may, in most stages of the disease, be most usefully administered. In the dysentery of children, no medicine exceeds in value the Hydrarg. c̄ Cretâ, combined with other remedies.”—*Med. Gazette*, June 25, 1841.

“As the incipient stage is usually past before men are sent to Hospital, general depletion and the means commonly applied to check the onset of the attack, are often inapplicable. Indeed, as dysentery is essentially an inflammation of a mucous not of a serous surface, it may be doubtful whether local be not often as effectual as general depletion. As to the use of calomel, which is so commonly employed at its onset, the general feeling of the profession seems to be against its employment at a later stage, and indeed it is difficult to conceive what beneficial influence it can exert on an ulcerated surface.

The usual practice in the Seaman’s Hospital is the free exhibition of sugar of lead and opium, and it appears to answer well, and has never produced those disagreeable constitutional effects which are attributed to the use of the preparations of lead. As much as from nine to fifteen grains of sugar of

lead, combined with small quantities of opium, (one-half or a whole grain of opium to three of sugar of lead) are given within the twenty-four hours, and this treatment is continued for several days, along with the free use of leeches and opiate and sugar of lead enemata, with purgatives every other morning and milk diet. This mode of treatment has been tried very extensively among natives, and appeared to be very efficacious and convenient, as avoiding the chance of salivation. It would seem to be peculiarly adapted to that insidious form of hæmorrhagic dysentery, supervening on chronic disease, of which several cases have of late occurred in middle aged men, characterised by the pouring out of immense quantities of blood from the mucous surfaces, indeed compared by some to uterine hæmorrhage. But we do not advocate any exclusive mode of treatment, or assert the superiority of sugar of lead over various other remedies, and of course, after the dysenteric symptoms have begun to subside, various alteratives are useful in aiding the bowels to regain their healthy tone."—*Seaman's Hospital Report for 1844.*

On Dysentery in Children.

2. As the number of autopsies in cases of dysentery, occurring in children at the General Hospital, is extremely small, and as no such case is included in the preceding tables, I add the post mortem appearances, in a case lately treated by my colleague Dr. Cantor, and obligingly communicated by him to me. It will be seen that the case was one of simple ulceration, without any sloughing; also that the ulcers were uniformly diffused along the mucous membrane, and that the solitary glands of the cæcum were not specially implicated; the process of ulceration was still going on, and that of reparation had not commenced at any point.

Child ætat. 4—ill for 3 weeks, death preceded by convulsions.

Large intestine studded throughout with equally diffused ulcers. The earliest stage of the ulcer was a raised white point, with greyish contents, (enlarged mucous follicle?) which enlarged and ulcerated, destroying the mucous membrane. The ulcers varied in size from a pin's head to a six-pence, and had raised margins of a white finely fringed appearance. The cæcum and rectum were somewhat thickened: appendix vermiformis studded with numerous minute grey points. There were nine lumbrici in the large intestine.

The small intestines were healthy, slightly injected here and there.

Liver somewhat small. Spleen healthy.

There were clusters of enlarged mesenteric glands, and some single ones enlarged, while others were natural.

A P P E N D I X.

Use of large Enemata in Dysentery.

1. *Out of India.*—With the use of enemata in dysentery, the profession has been familiar, at least since the days of Celsus. That author recommends various soothing and oleaginous injections. And to come to more modern times, Sydenham used not merely small opiate clysters, but enemata of $\frac{1}{2}$ lb. of milk. In the systematic use of these three or four times daily, he had the utmost confidence, although he remarks, that really bad dysentery, with much structural change, will not yield to clysters, whether purgative, astringent, or emollient. Böerhave exhibited emollient clysters three or four times daily.

It would be tedious to enumerate the different authors who have recommended particular enemata in this disease. As fashion has varied, they have been in high repute, or fallen into comparative disuse. The French especially have been fond of their lavements and demi-lavements, and have thought them peculiarly efficacious in the commencement of the disease. And of late years, in the dysentery of children, Trousseau strongly urges the employment of enemata of nitrate of silver of 8 or 10 oz., to be thrown up in a child of two years old with the long tube, after the bowels have been washed out with a common warm water lavement. All these enemata were meant to act on the ulcerated surface of the

colon, and all authors seem to have believed that with a syringe used in the common way, injections may be made to pass up as far as the ileo-colic valve,—and recently, Boudin states, that he has had positive evidence of such enemata passing beyond it.

Some twenty years ago, O'Beirne proposed the introduction of a long tube above the sigmoid flexure, (on the idea of its often being the seat of spasm,) to clear out the intestinal canal more effectually, and published two cases of dysentery which he had cured by it, and at a later period a third one. Whether his plan has been acted on in England, where the disease is rare, I am unable to say, but in 1840, Dr. Symonds (Library of Medicine) calls attention to the practice, and in Ireland, where epidemics of it are still not unusual as in former centuries, and where there must have been free scope for its use, it seems not to have been employed.

The chief novelty of late years appears to be the use of injections of water and the albumen of eggs by Mondière. He injected this mixture thrice daily, and by his own account, with wonderful success. All writers, however, with the exception perhaps of O'Beirne and Mondière, seem to have regarded enemata only in the light of most useful adjuvants.

2. *In India.*—To turn to tropical dysentery, it would be difficult to find an author who does not recommend enemata in one shape or another. In 1639, Bontius our first writer is found using them,—in 1783, Mathews advises their employment “for sheathing the bowels and obtunding the fluids.” He gives clysters of bark to support the system, or of tobacco to clear out the bowels, and these he administers through a hookah snake. With so formidable an instrument it seems probable that his enemata were large. Since his day, I imagine that few practitioners have failed to have recourse to clysters, small when opiate, and of about one pint in bulk, when meant to act generally on the surface of the bowel.

Sir James MacGrigor, for instance, in the beginning of the century, records the employment of almost every possible variety of them.

In 1837 Dr. Jackson, of H. M. 6th Regt., (a great advocate for the employment of clysters in ordinary constipation, and who recommended six pints to be usually thrown up), mentions having passed up the long tube 15 inches beyond the anus in a fatal case of dysentery.* Large enemata of tepid water, and the injection three times daily of a pint of water with nitro-muriatic acid, are frequently recommended by Madras writers some ten years ago; and in 1840, the officiating editor of the Madras Journal mentions, that he has often, in dysenteric cases, pursued O'Beirne's plan, and given wonderful relief with large emollient enemata.†

Enemata of various sorts have been constantly used in Bengal. Acetate of lead seems to have been always a favourite. Twining used it often; and up to 1840, Mr. Raleigh employed it largely in the General Hospital, and believed that with a pint and a half of fluid he covered the surface of the large intestines: these injections he repeated every 2 or 3 hours in hæmorrhagic cases.

In the end of the year 1847, Mr. Hare again invited attention to O'Beirne's method, and to the systematic use of large enemata. He thinks that they have not been hitherto used in sufficient bulk or with sufficient frequency, and he endeavours to assign to them a position of primary importance in the treatment of dysentery. He seems with Jackson to have arrived at a maximum of six pints. He believes "that he may save the lives of hundreds who die under the present system," which he considers to be commonly salivation, and hopes to produce quite a revolution in practice. Vari-

* "No one has thought of making use of this method."—Mr. Hare on Dysentery, p. 14.

† This mode of treatment ought to have been noticed and applied in India.—Hare, p. 7.

ous medical officers have since published cases of the successful employment of large enemata, chiefly in chronic dysentery.

To raise them into this new position of independent value, it appears to be necessary that we should have a series of bad cases of dysentery treated successfully by the long tube, unaided by the various other modes of internal treatment.

Whether Mr. Hare's zeal and energy will be able to elevate them to this new rank, or whether the long tube will share the fate of the long-forgotten hookah snake of Matthews, it is for the experience of future years to determine, for like other remedies, injections vary in efficacy in different seasons and in different forms of the disease. Of great value as injections are, they certainly produced no diminution of mortality in the years, in which they were employed in the General Hospital most frequently and in largest quantity, namely, in 1848 and 1849: and the consideration of their past history does not encourage the hope that they will ever permanently retain a position higher than that of most useful adjuvants. The large ones can be used systematically only in Hospitals, and are never likely to become favourites in private practice.

Use of Quinine in Remittent Fever.

An apology may be deemed necessary for collecting at this time of day, evidence in favour of the free use in remittents, of one of the very few specifics we possess in the whole circle of the materia medica. Yet it is possible, that some may not have had the opportunity of carrying their investigations on the subject further than my friend Mr. Hare, who thus expresses himself: "I have searched every where indeed, and all that I can find in books and Magazines since Johnson's time till now is, bleed and give large calomel pur-

gatives, but be very careful not to give quinine too soon." *Hare on Fever*, p. 13. To such readers the following information may be interesting from its novelty, and to others perhaps, as being a connected historical sketch of the use of quinine in remittents in India. After all, the use of quinine in remittents is merely an extension of its use in intermittents, and in either case, only a substitute for the older treatment by bark ; but to trace all this would lead us into still further details, while I fear that even the following brief recital of facts may be tiresome to many.

1. *Out of India*.—In the year 1835, M. Maillot published a work on the fevers of Africa, and states that, by the exhibition of quinine he reduced the mortality in bad remittents from 1 in $3\frac{1}{2}$ to 1 in 20. He gave it without any reference to the stage of the fever, and in bad cases he gave from 1 to 2 scruples by the mouth, and a drachm as an enema. In this manner he in several instances gave as much as 148 grs. in 24 hours.

The following extract from Bouchardat's *Materia Medica*, published in 1839, will show the state of European opinion and practice at that period. "In bad remittents the salts of cinchona save the patient from certain death. In these cases we must act much more rapidly, and augment the doses. We must then, according to Torti, administer the bark on the first sign of remission, for there is often no intermission. Bretonneau goes still further : he commences the exhibition of bark in the middle of the paroxysm, and is not afraid of increasing the intensity of the access during which it is given ; for experience shows, that the medicine does not commence to act till several hours after it has been administered, i. e. during the remission. Bretonneau used bark, but we now greatly prefer quinine, which is more quickly absorbed, and acts more rapidly. We prescribe 30 grs. of it with 1 gr. of opium, and give it in 3 doses. From the moment that the access has been prevented or moderated, it is no longer ne-

cessary to give it in these large doses, but we may continue to give smaller ones for some days."

Dr. Hille in Casper's *Wochenschrift* in 1839 mentions, that quinine is the only medicine for the intermittents and remittents of Surinam; and in 1842, writing in more detail, says, it must be given in 12 to 40 gr. doses in all stages.

The American physicians seem to have been somewhat slow in following their Continental brethren, and although quinine was given in yellow fever at New Orleans in 1839 in scruple and scruple and a half doses,* I believe that Dr. Byrne (*Boston Medical and Surgical Journal*, July 1845) is the first who talks strongly in favour of quinine in remittents. "He found that quinine might be used with the greatest advantage even in the hot stage. As regards the dose, he observes, that in ordinary remittents it is a matter of indifference what dose is administered, provided 10 or 20 grains are introduced into the system a certain number of hours before the paroxysm. In severe cases, there is no longer any choice; but the quantity above specified should be given in 1 or 2 doses at most. We thus find, that by the middle of the year 1845, the use of quinine in large doses, without much reference to the stage of the remittent, was fairly established, and the year 1846 produced papers by Drs. Tuck, Upsher, and Professor Van Buren, and the year 1847, by Holmes,† all recommending the practice. It is unnecessary to allude to writers of later date.

In India.—The free use of quinine in remittents appears to have been slowly and cautiously adopted, and this invaluable medicine seems on the whole to have been more extensively used in Madras than in the sister presidencies. Some of the earliest notices on the subject are the following: Dr.

* Mr. Hare alludes to this, p. 15.

† Mr. Hare mentions Mr. Holmes, page 14.

Geddes in 1828 mentions giving quinine with the pulse at 108, and after extensive use of it, he in after years announced, that he preferred giving it in every instance to affecting the mouth with mercury. In 1833, Corbyn treated a bad remittent fever in Calcutta with most gratifying success, and says, that his routine practice was, six or seven hours after the administration a purgative to exhibit 7 or 8 grs. of quinine, and to continue this steadily every four hours without reference to paroxysms. He seems to have produced giddiness and deafness frequently. In 1834 Dr. Wright, in the Goomsur country, gave the sepoy 20 grs. daily, but used to watch for remissions: in 1835, Mr. Eyre gave it in the same country often in scruple and a half doses, of which the Superintending Surgeon disapproved; and, the supply of quinine running short, Lord Auckland sent down a supply of 20lbs. at his private expense,* when every one who could get quinine, employed it with an unsparing hand, and I believe with very little regard to remissions.

In 1835, Twining, in the second edition of his book, strongly recommends it in remissions; and Goodeve in 1837 says, that it has produced quite a revolution in practice, having almost brought back the days of bark: still he waits for remissions. In 1840 Mr. Green recommends its free use in remissions, and the Madras Journal of the same year calls attention to Maillet's large doses. In this year also, Dr. J. Murray, writing of the Malwa sweating sickness, says, there were few conditions that would prevent his giving quinine; that he was not deterred by headache, but if it were violent, he would use other measures previously. In 1841, Mr. Martin writes, "when the remissions are well marked, quinine should be given without waiting for every thing; if we wait for every thing, we shall often wait till it is too long or too late." In 1842, Dr. Bell in Persia gave quinine in small doses without any

reference to the stage of the fever, and in the same year the recorded practice in the General Hospital at Madras was as follows: "Quinine is given in 5 gr. doses every hour on the least tendency to remission, or change, however slight. No bad consequences, nor any aggravation of the symptoms, have been observed, even after it has been continued throughout the greater part of the exacerbation."

In 1843, Dr. Morehead expressed himself thus clearly regarding the use of quinine in remittents at the General Hospital of Bombay: "I think that even in cases where the remission is very imperfect, quinine should be tried, and repeated or not according to the effect." "Should the remission be distinct, dryness and brownness of the tongue offer no drawback to the use of quinine." "I do not think the presence of the phlogistic diathesis or the presence of fixed congestions of necessity contra-indications to the use of quinine." "We are frequently, it is feared, induced to intermit the remedy, because a febrile exacerbation may have followed after its use in the first or second remission, under the apprehension, that the febrile excitement has been produced by the remedy," "but we have no sufficient grounds," &c. In this year also, though he waits for a remission, MacGrigor talks of the paramount importance of keeping off the next paroxysm. About the same time the present writer published a case in which quinine was successfully administered during grave cerebral congestion, notwithstanding the presence of a black tongue and sordes about the teeth; and in 1844, in the report of the Howrah Seaman's Hospital, he attributes the great reduction of mortality from the 8 per cent. of preceding years to 1.7 in one year, and to 0 per cent. in the next, or no death in 127 cases, to abstinence from free venesection and the early use of quinine, always making allowance for what the Germans call the "*Genius epidemicus morborum*." In the same year, Eveleigh in the Calcutta Journal mentions his having

employed quinine in severe cases, in which the head and the liver were affected.*

Meanwhile, quinine was becoming more extensively used, and especially in parts of Madras. In the bad remittents in Goomsur it was given in very large doses, and the Government supply of quinine did not equal the demand. It used even to be reported facetiously, that the Madras Board had recommended the appointment of a Special Superintending Surgeon to controul the extravagant expenditure of quinine. In 1845, the Bengal Medical Board was applied to, and liberally sanctioned a small extra supply for the Hill Agency, but very many lbs. were procured from private sources at the expense of the Agent. Mr. Cadenhead regularly and constantly used it in the hot stage of fever, from the commencement of the year; and at the request of the Editor of McClelland's Journal, began to write a paper on the subject, which, owing to the distraction of political duties, was never finished. There was a regular schism among the practitioners in the Northern division regarding the stage at which it should be given; in short, the most impartial account of the state of the question at this time is given by Dr. Williams, of St. Thomas, writing in the same year. After alluding to the use of large doses of quinine in India and in various parts of the world in remittents, he remarks: "The battle still rages between those who would treat this disease symptomatically, and those who prefer the specific remedy."

Two years after this, or towards the close of 1847, Mr. Hare published a pamphlet, in which he most strongly advocated the use of quinine from the commencement in all miasmatic fevers. He advises its administration without any reference to the stage of the fever, and his specific directions are chiefly the same as Mr. Corbyn's treatment of the fever

* "This is the only approach to the truth I can find."—Hare, p. 12.

of 1833. Since then, Dr. Macrae of Howrah has published an account of many cases of fever treated similarly: this mode of treatment he has pursued since the middle of 1847, as shewn by the records of the Seaman's Hospital; and Dr. Ford of Madras states, that he has been following a similar practice since the year 1845, as testified by his reports to the Medical Board.

By this time pretty abundant evidence had accumulated on the subject, and I do not know that any more recent has been submitted to the profession in Bengal. A paper by Dr. Cameron of Ceylon, advocating the same views, has since appeared, but I have not seen it.

To the enquiry, what has been established by all these facts regarding the free use of quinine in tropical remittents during the last twenty years, the following would seem to be the natural reply.

It admits of no question, that the free and early employment of quinine in remittent fever is a great step in advance in practical medicine, and it is also important to know that there is no necessity for extreme caution in its use, but the late Dr. Williams, as already quoted, describes the state of the case correctly. It remains for practitioners to adopt the symptomatic or the specific mode of treatment: or a mixture of both, which may be termed the eclectic. Much must of course depend on the intelligence and experience of the individual, on the circumstances in which he may be placed, and on the character of the fever which he may be called on to treat. No practice should be indiscriminate, for such treatment is never scientific, and is not always safe.

OBSERVATIONS
ON THE
RECENT OFFICIAL REPORT
ON
FEVER AND DYSENTERY.

BY
JOHN MACPHERSON, M. D.,
FIRST ASSISTANT SURGEON, GENERAL HOSPITAL.

“ Was there any novel mode of practice introduced,—was the nature of that practice beneficial, and such as to render its general adoption desirable ? ”

Medical Board in 1816.

Calcutta :

R. C. LEPAGE AND COMPANY, BRITISH LIBRARY.

MDCCCLII.

F. CARBERRY, BENGAL MILITARY ORPHAN PRESS.

CONTENTS.

	<i>Page.</i>
LETTER TO DR. MOREHEAD.	v.
1.—NATURE OF THE EXPERIMENT IN THE WARDS OF THE GENERAL HOSPITAL,	1
2.—EXAMINATION OF THE TABULAR STATEMENTS,	7
1.—Tabular Statements Generally,	<i>ib.</i>
2.—Tables of General Hospital,	11
3.—Returns of H. M.'s 70th,	17
4.—Other Regimental Returns,	19
3.—SOME POINTS OF THEORY AND PRACTICE,	23
FEVER.— <i>a.</i> Cause,	<i>ib.</i>
<i>b.</i> Inflammatory Nature,	24
<i>c.</i> Use of Quinine,	26
<i>d.</i> Mixed Treatment,	28
<i>e.</i> Arsenic,	<i>ib.</i>
<i>f.</i> General Statistics,	29
DYSENTERY,— <i>a.</i> Quinine,	30
<i>b.</i> Mixed Treatment,	33
<i>c.</i> Long Tube,	34
<i>d.</i> Calomel Treatment,	37
<i>e.</i> General Statistics,	39
QUININE—ITS SAFETY?	40
CONCLUSION,	44

TO C. MOREHEAD, Esq., M. D.,

Bombay Medical Service.

&c. &c. &c.

MY DEAR SIR,

I TAKE the liberty of addressing the following remarks to you, as one of the most constant and successful cultivators of practical medicine in this country.

I wish that they were of a less controversial nature, but, however undesirable in our profession controversy may be, there are occasions on which it is impossible to avoid it.

On the first appearance of the Bengal Report on Fever and Dysentery, I published a statement, which showed how far its conclusions were borne out by the records of the General Hospital, and having then examined the Tabular Statements of the Report that referred to the Institution with which I was connected, I thought that I had done all that was necessary on my part. But though the Analysis was, I believe, satisfactory to those who were in some degree

conversant with the facts of the case, and though its statements remain at the end of seven months unchallenged, it has, I have been told, been generally considered very much too concise, and the following pages are meant to remedy that defect.

Although also I knew that there was a very general wish throughout the profession in Bengal, that a more extended reply should be given to the Report, I remained silent, hoping that some influential member of the Service would take up the subject. But as I now see no prospect of this, and as besides, I have access to information on this particular subject, which no one at a distance can possess, I again venture to come forward.

The following observations, in which the Report is treated, purely as a matter of medical science, and with as much freedom, as if it had not been issued in an official form, fall under three heads:—

1st.—An examination of the nature of the experiment within the wards of the General Hospital.

2nd.—A review of the Tabular Statements.

3rd.—An inquiry into certain points of Theory and Practice.

In the last part I shall not follow the Experimenter in all his speculations, but confine myself chiefly to some points on which absolute originality has been claimed, and of which the success, according to the official account, "has been triumphantly demonstrated."

I cannot pretend to absolute accuracy in all the numerical statements which I have made, but I believe that their general correctness may be depended on. For such inaccuracies as may be discovered (and I shall feel obliged to any one who will point them out to me) some apology will, I hope, be found in the great difficulty which there is, in dealing accurately with the crowded general tabular Statement appended to the Report.

I am,

My dear Sir,

Yours very truly,

JOHN MACPHERSON.

November 8th 1852.

I. ON THE NATURE OF THE EXPERIMENT IN THE WARDS OF THE GENERAL HOSPITAL.

" Equidem dicam ex animo, Patres conscripti, quod sentio, et quod vobis audientibus sæpe jam dixi."—CICERO IN PISONEM.

THE Medical Board, in an early part of their Report remark, "that the method of figures, extracted from a wide field of observation, is the least fallible of all methods."

True it may be in the abstract, that there is only one method that can give solid results in medicine, that which is based on numbers, and supported by exact observation and accurate induction. But then the numbers themselves must be arrived at with the utmost care and consideration: the figures must represent facts, which have been tested and indisputably proved. If the numbers are hastily assumed, the air of method that belongs to numerical calculations, is only an additional source of error and confusion. Yet even when the numbers are correct, and all the bearings of the cases they represent, have been well ascertained, they may not afford a basis wide enough for any general conclusion.

Hence it is not surprising, that many practical men look on this method as a very poor test for the comparative value of different modes of treatment, and that it has even been seriously said, that it is possible to prove anything regarding a given mode of treatment by means of numbers—the homœo-

pathists, for instance, referring to an array of figures as evidence of their success.

The numerical method has, in short, failed often, and in such instances of failure, mainly from its imperfect application.

There are two conditions for obtaining valuable results from any experiment in medical science—the experiment must be conducted with extreme care, and it must be made on a considerable scale. Both these propositions may be almost termed self-evident, but the first of them deserves more minute examination. I shall now therefore inquire, what points should be borne in mind in an investigation of this kind, and how far such points were attended to in the experiment at the General Hospital.

1st. It is of primary importance to ascertain and lay down definitely, what the particular mode of treatment is to be—for if it be a mixed one, it is extremely difficult for the most discriminating practitioner to ascertain what effects are really produced by the individual remedies which he employs.

What was the treatment in the Experimental Ward? Was it a pure and unmixed one? Solely Quinine and Arsenic, the long tube and copious injections? This I imagine is the common opinion. Yet general bleeding and leeching, purgatives, opiates, metallic astringents, &c. &c. were all used in that ward. Has it been ascertained to what extent they were used? What degree of influence over disease was to be attributed to them, and what to the first-mentioned remedies? Was it ascertained in bowel complaint how much was effected by quinine, how much by the other remedies, and how much by injections? Was it even ascertained how often on an average the long tube was passed, in each case? On all these points information is required, and until it is fur-

nished, the experiment must remain a crude and imperfect one.

2ndly. To make the results practically useful, it should be accurately ascertained for what forms and for what stages of disease, a given mode of treatment is peculiarly suited. But in the published results of this experiment we have no Tabular Statements, showing in what varieties of fever and of bowel complaint the treatment was most successful. All the cases of Fever appear under one head, and the cases of Diarrhœa and of Dysentery are also for the Experimental Ward not distinguished from each other, much less acute and chronic dysentery, or the complication of liver affection. As to treatment—"Plenty of quinine" is recommended for all fevers, and for malarious dysentery; it is apparently not recommended for acute dysentery, but then we are not told how to distinguish the two. In short the tendency of the report, and of the practice recommended in it, is, mainly owing to want of precision in defining the experiment, and stating its results, to encourage careless observation, and want of attention in making out the changing phases of disease and their periods, considerations very important as regards its safe and successful management. The serious mistakes that may arise from the want of precision alluded to in this and in the preceding paragraph, are well illustrated, by contrasting in another part of these observations the results of the quinine and non-quinine treatment in the Experimental Ward.

3rdly. For comparative purposes, it is necessary that the patients treated should be all generally of the same class, and that the admissions should take place continuously to all the wards, which are the subject of comparison, as we all know the effects of season in modifying disease. Now, so far as I am able to follow the Tabular Statements of the

Report, the cases of women and children (the latter a class usually yielding a high mortality) a description of patients not treated at all in the Experimental Ward, are included in the Returns with which those of the Experimental Ward are contrasted—and besides this, the admissions to the different wards were interrupted. At one time the Experimental Ward took in all the cases of fever and of bowel complaint, then again it ceased for a few days to take any in, while at another time it only took its share of them with the other wards, and at still another, it admitted a few cases of neither fever nor bowel complaint. All these are disturbing elements, and must detract from the value and accuracy of the results.

The absence of children would tell in favor of the Experimental Ward, the other irregularities might or might not be in its favor according to circumstances. Regarding the influence of chance in these matters some remarks were made elsewhere, and every day affords fresh illustrations of it.*

4thly. It is very necessary to ascertain the exact condition of a patient on leaving Hospital. Boudin, when conducting his experiment with arsenic before a committee, kept his fever patients in hospital for 15 days after they were cured, that there might be a certainty of no relapse. At the General Hospital no such condition was observed. No one saw the patients before they were discharged, except the Medical Officer who had treated them. The Hospital Returns give no list of partial recoveries: they have only the headings "Discharged" or "Died," for the headings "Transferred" and "Invalided" give little information. The discharged are all supposed to be cured. Yet how many of the discharged may have died soon after in other hospitals, and how many

* "Analysis," p. 11.

of them were re-admitted*? This remark of course applies equally to all the wards of the General Hospital.

5thly. A patient, if not rejected at once as an unsuitable one for the experiment, should be treated from beginning to end, in the ward into which he has been admitted. He is not to be set down as a case of fever or of bowel complaint, if he be not one of either, but he ought to be written off to the total general results of the ward into which he was admitted. Now at the General Hospital, (where I may explain that there were three wards, an Eastern, Western and Experimental, each under a separate Medical Officer,) this system was only partially adopted. Out of the 24 deaths occurring in the Experimental Ward, 16 are written off, as not being cases of fever or bowel complaint, and they consequently do not appear anywhere in the Report, while at the same time two dozen cases, one half of which proved fatal, were, after an average of four days' treatment, sent, as not being cases of fever or bowel complaint, to the other wards,† instead of being written off like the rest. This has necessarily led to inextricable confusion of cases; it has, however, afforded an opportunity of judging of the effects of the treatment in the Experimental Ward in diseases, other than fever and bowel complaint.

6thly. A committee, or at least one person besides the experimenter, should watch the progress of the cases from day to day, and be present at the *post mortems*. No such precaution as this, was, as far as I know, adopted, and the duty would no doubt have been an irksome one. Still to obtain satisfactory results some such measure was absolutely

* See *infra* Dysentery. A reference to the records of the Medical College and Seaman's Hospital is desirable.

† If this had been the uniform practice, it would have been, comparatively speaking, unobjectionable.

necessary, and has always been adopted in inquiries of a similar nature.*

Regarding the other condition which has been mentioned, the sufficiency of the scale on which an experiment is conducted, every one will admit generally that, in drawing inferences from facts, the fewer the number of them, the less valuable will be the conclusions to which they lead, and every one will also, I presume, admit that in the present instance the scale of the experiment was small. It seems therefore hardly necessary to enlarge on this topic.

And thus, the want of method in conducting the experiment, the short period of time over which it extended, and the smallness of the number of patients that were the subject of it, would appear to deprive the experiment of almost all value.

As, however, in the opinion of the Board, one season at the General Hospital has produced clear and positive results, it may be well, although enough has been said to show that the basis on which the experiment itself rests, is not a very secure one—to examine the Tabular Statements adduced in proof of its success.

* Many of these points were, during the course of the experiment, brought to the notice of the Medical Board by the experimenter or by myself.

II. TABULAR STATEMENTS.

“ Behaupten ist nicht beweisen.”

I. THE TABULAR STATEMENTS GENERALLY.

1. The most general expressions of the Medical Board on the subject of the success of the Experimental Ward are the following:—“ We have prepared tables to prove the absolute reality of the superiority, positive as well as comparative, which he claims for the methods of treatment pursued by him over those pursued by others,” and the experimenter remarks “ I have actually had only one-half and one-fifth the mortality of my neighbours,” *i. e.*, the Eastern and Western Wards of the General Hospital and of H. M.’s 70th.

2. The position of the Medical Board is supported by the only abstract which they make, that of the results yielded by the general comparative statement; and it is a comparison of the percentage of mortality of the two Experimental Wards, one in the General Hospital and one in that of H. M.’s 70th, added together, with that of the other wards of the General Hospital and of H. M.’s 70th, and of all other European troops serving in Bengal, added together. It is as follows:—

	<i>Diarrhœa.</i>	<i>Dysentery.</i>	<i>Fever.</i>
Experimental wards, ...	0	4·91	0·7
All others,	3·24	10·55	1·24

3. If the Board had only this general comparison of results before them, their strong commendation of the Experimental Ward is not surprising, as it is apparently fully borne out by figures. But it is just possible that in their anxiety “ to pre-

vent cavilling at the narrowness of the field of observation" they may have run into the opposite extreme, and have selected too "wide a field of observation." It seems improbable that they would have arrived at the foregoing conclusions, if they had first closely examined the facts occurring within the narrower field of observation, and then proceeded to the wider one, using the facts supplied by the latter chiefly as illustrations. This is the course which I shall endeavour to pursue.

The special examination of the returns will presently show that these results are based on inaccurate data. But apart from this, some general objections must be made here to the nature of this comparison.

4. It classes and compares the results of the treatment in the various wards of the General Hospital, with the general results of the army. But every one knows that the class of patients that frequents a General Hospital, is very different from that which fills Regimental Hospitals, and that therefore the two cannot be considered fair subjects of comparison. In like manner, the character of disease varies so much in different places in the same season, that the making the comparison too wide a one, may defeat the very object of the comparison. Thus to take by no means an extreme case, from the tabular statement, H. M.'s 3rd Dragoons, at Umbala, lost only 2 per cent. of its cases of bowel complaints, while H. M.'s 98th at Peshawur, lost 9 per cent. at the same time. Could this difference be attributed solely to difference of practice?

5. Another general objection to the returns is, that, for the Experimental Ward they appear to give the cases of dysentery and of diarrhœa, added together under the head of dysentery, as will be immediately shewn, but that in the other returns diarrhœa and dysentery are very properly classed separately. On this account I shall throughout

compare the returns under the head of bowel complaints generally—and this I do, not from choice, but from necessity, as under these circumstances it is the only mode, in which the comparison can be a just one. Though this is forced on me, I regret it less, for reasons which will be constantly recurring in these remarks, and here I must enter on what is somewhat of a digression, but is a matter of considerable practical importance, in ascertaining the mortality from bowel complaints.

6. There can be no doubt in the abstract that for scientific purposes and for defining modes of treatment, diarrhoea and dysentery should be treated separately, as also acute and chronic dysentery. But on a former occasion, when compiling some tables of the mortality from dysentery, I remarked* “of course the mortality in returns may be made to vary, according as cases are classed under the heads of diarrhoea or of dysentery, a point often requiring nice discrimination.” I even found extreme difficulty in separating hepatitis from dysentery, owing to their frequent association.

The following are some of the reasons why diarrhoea and dysentery cannot well be classed separately with much accuracy in general returns. (*a.*) One man (I know this practically) will often regard as a case of chronic dysentery, what another looks on as diarrhoea. (*b.*) Many cases first admitted into hospital under the head of diarrhoea, are, when their symptoms become more marked, transferred to the head of dysentery, and the fact may very easily be lost sight of. (*c.*) On looking over a series of years, as for instance in the tables of the General Hospital and of the Garrison, appended to the Report, it appears, that though the mortality from diarrhoea or dysentery taken singly may vary much, yet the results for bowel com-

* Bengal Dysentery, p. 4.

plaint in general are wonderfully uniform ; thus, in my opinion, showing the cases to be returned as dysentery or diarrhoea, very much according to the taste of the Medical Officer.

(d.) On looking over the general tabular statement attached to the Report, we find that much mortality in dysentery is frequently attended with small mortality in diarrhoea, and *vice versa*, a fact which probably also depends mainly on the same cause, as the uniformity of mortality in bowel complaints just alluded to ; for instance :

In H. M.'s 70th	the mortality for diarrhoea	was 2.2	for dysentery	42.
„ H. M.'s 60th,	„	1.3	„	18.
And <i>vice versa</i> , H. M.'s 98th,	„	8.5	„	9.2
A detachment of Recruits,	„	8.	„	7.

(e.) Another reason is, that on reviewing the results among the European troops on a large scale during the last 20 years, as given in the table at the end of the Report, it appears, that the number of cases of dysentery has been rather diminishing, while that of cases of diarrhoea, has increased, facts not easily explained by natural causes, but of which the possible explanation may be, that cases of bowel complaint are not set down as cases of dysentery, so much as a matter of course now, as they were formerly.

On the whole, therefore, I trust, that the reader will be inclined to agree with me in considering it useful, to correct any extreme results got for dysentery alone, by a constant reference to the mean average for bowel complaints.

7. As far as I am able to ascertain, the cases of women and children, (the latter a class always yielding a very high mortality), a description of patients not treated at all in the Experimental Ward, are included in the returns, with which those of the Experimental Ward are contrasted. The absence of children from any ward would of course reduce the mortality in it.

Having stated the preceeding objections to them as they now stand, I proceed to the special examination of the tables, and I shall first turn to those for the General Hospital.

II. TABLES OF GENERAL HOSPITAL.

1. THE tabular statement assigns the following comparative percentage of mortality to the Experimental, and to the united Eastern and Western Wards of the General Hospital.

	<i>Diarrhœa.</i>	<i>Dysentery.</i>	<i>Fever.</i>
Experimental,	0	7·3	0·79
Eastern and Western, ...	2·1	15.	1·4

But, on the returns from which these deductions are made, two preliminary observations must be made, one of paramount, and one of secondary importance.

2. According to the tables, not a single case of diarrhœa occurred in the ward devoted to fever and bowel complaints, during a period in which forty-seven cases of that disease occurred in the other wards.* This is at first sight a startling fact, but fortunately it can be easily explained, as the experimenter has in his own handwriting fully and fairly recorded in the public registers of the hospital, about thirty-five cases of diarrhœa and about sixty-six of dysentery, from which the ninety-eight cases of dysentery in the tables must have been obtained. It is obvious, therefore, that these ninety-eight cases cannot be regarded as so many cases of dysentery, but of bowel complaint generally, *i e.* diarrhœa and dysentery taken together.

* If I may venture on a conjecture, those forty-seven cases appear to me to represent fifteen cases of diarrhœa that occurred in the other wards, and thirty-two that occurred in the Experimental, added together by some strange confusion. The tabular statement gives the other wards eighty-two cases of dysentery, whereas I can find only thirty-eight in the hospital records,—and curiously enough, the addition of the forty-seven cases of diarrhœa to the thirty-eight of dysentery gives a close approximation to the number eighty-two.

3. The tables would also seem to imply that the Experimental Ward did not treat any military patients in the General Hospital, whereas there was no difference between the wards in this respect, of which I am aware; it will therefore simplify matters, if we first add together for the other wards the two separate columns headed military and non-military, and then contrast the results with those of the Experimental Ward, as has been done in the above abstract.

4. But taking the tables as they stand, the Board's emphatic commendation of the treatment in the Experimental Ward is hardly borne out, and the statement of the experimenter that "his mortality was only half that of his neighbours" is most certainly not supported by them, for the general tables assign (taking the totals in fever and bowel complaint) only a superiority of 1.52 to the Experimental Ward, this being the difference between 3.64, the percentage of mortality of the Experimental, and 5.16, that of the other wards. Even if the experimenter's statement be limited to bowel complaints (for with these tables the separate consideration of dysentery is out of the question), it is still equally inaccurate, for the tables assign the Experimental Ward a mortality of 7.3 and the others one of 10.2 per cent.

Now, under the circumstances of the case as already detailed, and especially the absence of women and children from the Experimental Ward, the superiority of the treatment in that ward appears to me, even according to the tables, to be wonderfully small.

5. But having discovered the mistake about the diarrhoea cases, and finding it, after many fruitless attempts, impossible to reconcile the figures in the hospital books with those in the tables, (for the tables say there were 180 admissions of dysentery, and 338 of fever, while the books say they were 88, and 268, respectively,) I scrutinized those records,

and from data furnished by them shewed (Analysis, Calcutta, March 1852) that not challenging the number of admissions assigned to the Experimental Ward, (as I can see how they are got, though I cannot see how the numbers are got for the other wards) the mortality in fever and bowel complaints was in the Experimental Ward 4.46 per cent., and in the other wards taken together 6.14. To this I am now able to add, that this somewhat smaller mortality in the Experimental Ward was caused solely by its not treating women and children, for if those classes of patients be struck off from the other wards,* their mortality was only 4.2 per cent. : consequently their mortality was absolutely somewhat less, than that of the Experimental Ward.

6. It was also shown that for fever and bowel complaints, the mortality in the Western Ward of 2.9, was considerably less than the 4.46 of the Experimental, being rather less both in fever and in dysentery.

7. Further, that the total mortality of cases of all diseases in the Experimental Ward, even after thirteen fatal cases were removed from it to the other wards, was 10 per cent., or much the usual average of the hospital for all diseases : and that there also was in the Experimental Ward a special and unusually high rate of mortality, not falling short of 80 per cent., from certain diseases, (such as apoplexy, epilepsy, ebrietas, scorbutus, pneumonia, hæmorrhoids, &c.) not being either fever or bowel complaint.

8. To this I can now add, that the total mortality from December 8th 1849 to October 31st 1850, of all cases first treated in the Experimental Ward, (and therefore including the fatal transfer cases) was 13.8, and the total mortality from all diseases treated throughout in the other wards (and therefore excluding the transfer cases) was 8.5, giving the large

* The mortality of children in the other wards from fever and bowel complaints was 13 per cent.

balance of 5.3 in favor of the other wards. As the other wards had to treat along with the common and less fatal maladies, the bulk of the most fatal diseases, such as cholera, hepatitis, delirium tremens, and also of an epidemic of small-pox, besides women and children, and as in the Experimental Ward considerably more than half the cases were of fever and diarrhoea, neither of which classes of disease yield a high mortality, the excess of mortality on the part of the Experimental Ward is remarkable. Even crediting the other wards with the fatal transferred cases, the balance is not turned, for they were still somewhat more successful, the numbers being then, 9.4 and 10 respectively.

This last comparison from its nature does not admit of being a very accurate one, but still it is by no means unimportant in its general bearings on the question.

9. I cannot consider, as some have done, that the last conclusions are foreign to the matter in hand, especially in the absence of accurate data for individual diseases. On the contrary, I regard them as of much importance in judging of the general results of the treatment in the Experimental Ward, for where is the practical gain, if a seeming diminution in the mortality from fever and bowel complaints is attended with an increased mortality in other diseases? Unless the fact of the two occurring simultaneously can be shewn to be a matter of accidental occurrence, it must materially diminish the general value of the treatment in the Experimental Ward.

In another point of view also one of the last conclusions is of much importance, for assuming the tables to be correct, it has been shewn that eighteen cases of diseases other than fever and bowel complaints, were attended with sixteen deaths.* Now, this is so highly improbable, that it seems to point to some radical fallacy in the numerical statements, or in the classification of diseases.

* Analysis, p. 5.

10. I have also thought it worth while to enquire whether the presence of the experimenter affected in any degree the total mortality of the General Hospital. As his number of cases was small, no very striking effect could well be produced; still his great success in fever and in bowel complaints ought in some degree, however trifling, to have reduced the total mortality of the season for all diseases. The total mortality for the last five years has been.

In 1847-8	9.2
48-9	13.4
49-50	11.3
50-51	10.8
51-52	9.4*

* It may be interesting to contrast these percentages of mortality with those of some European General Hospitals. The most important points to ascertain, and without a knowledge of which such comparisons are little more than matters of curiosity, are, the class of patients admitted, and the nature and the stage of their diseases, also the prevalence of local epidemics; thus the mortality in the Edinburgh Infirmary during an epidemic of typhus is at least doubled. The following data refer chiefly to periods before 1840, and are compiled from Quetelet, Porter, and Milne Edwards :

Paris—Hotel Dieu, ...	14.	England.	
Charité,	14.	Bartholomew's,	7.62
des Enfants,	22.5	London Hospital,	11.44
Pitié,	12.2	St. George's,	11.19
Lyons—Hotel Dieu, ...	9.	Dreadnought, (1849) ...	11.
Montpellier, all Hospitals, ...	10.	Manchester Infirmary, ...	7.16
Pesth,	16.6	Liverpool ditto,	5.57
Berlin,	16.6	Probably the class of patients in the Dreadnought comes, of all the foregoing, nearest to that in the General Hospital; but in it no doubt there is a comparative preponderance of pulmonary, in the latter, of abdominal, diseases. The average mortality in the General Hospital, when taken	
Vienna,	16.6		
Dresden,	14.2		
Munich,	11.1		
Brussels,	11.1		
Amsterdam,	12.4		
Milan,	16.6		

for large periods, seems to have remained nearly uniform since the commencement of this century. There has certainly been no diminution of mortality. It is curious to observe that the presence of cholera since 1817, does not seem

Now the experimenter was present during portions of the seasons 49-50, 50-51, and the considerable diminution of mortality in the first of these seasons as compared with the one immediately preceding it, and a further slight diminution in the second, would look as if his presence had reduced the mortality; but on looking back, we find that in the year 47-48 the mortality was less than in either of the two years during which he was present, and on looking forward, we find that after his departure the mortality was also smaller. Any first impression of his having diminished the mortality, is therefore at once dissipated.

11. I am quite aware that there are various ways of explaining these facts, and that as they at present stand, if we were to attempt to draw any positive conclusions from them, they would merely furnish an additional example of the fallacy of the numerical test, where every element has not been taken into consideration.

One conclusion, however, to which they point, ought not to be overlooked. In the years 49-50, 50-51, especially in the latter, I find that in the annual returns of the General Hospital there was a considerable diminution in the mortality from bowel complaints, although the general rate of mortality for all diseases was not affected. This bears out the result, which we had already arrived at by another process, that apparent diminution of mortality in certain diseases must have been accompanied by increased mortality in others, and it is not a conclusion very flattering to medical science.

to affect the general averages, nor could we guess the presence of a severe epidemic of small-pox from the returns of 1849-50. The General Hospital does not admit natives, and it has few Surgical cases; the mean number of patients treated annually during the years mentioned in the text, was 1336.

III. RETURNS OF H. M.'s 70TH.

1. According to the general table, the percentage of deaths in the Experimental and other wards of H. M.'s 70th was, as under :

	<i>Diarrhœa.</i>	<i>Dysentery.</i>	<i>Fever.</i>
Experimental	0	4.09	7
Others	2.2	42.	1.4

For the examination of these returns, I do not possess the advantages which I have with regard to those of the General Hospital. I have no means of examining the books of H. M.'s 70th, or of any other regiment, nor any means of scrutinising their returns. My examination of them must, therefore, be a comparatively meagre one. Yet, a very cursory glance at the general tabular statement, excites the suspicion that there must be some fundamental error in them, as in those of the General Hospital.

2. As I understand it, the medical charge of certain companies was given to the experimenter, while the remaining and main portion of the regiment was treated by its own surgeon. Now the tables state, that the Experimental Ward had only 1 case of diarrhœa, whereas during the same period the surgeon had 135 cases of that disease. They also assign to the Experimental Ward 248 cases of dysentery and 1 of diarrhœa. The fact, of 1 case of diarrhœa occurring in the returns of the Experimental Ward of H. M.'s 70th, is much more surprising to me, than that none should have occurred in that ward of the General Hospital. Because it looks as if the cases of diarrhœa and of dysentery had been carefully examined, and separated intentionally, and produced the singular result of 1 case of diarrhœa to 248 cases of dysentery in the Experimental Ward of H. M.'s 70th ! a fact if possible more surprising, than that all the diarrhœa cases but one in the 70th, should have fallen to the surgeon. Then

again, the surgeon is set down as having 18 deaths out of 42 cases of dysentery, or the unheard-of mortality of 42 per cent., a mortality that must have depended on some more general cause, than any mere difference in the mode of treatment. These facts alone are sufficient to convince any practical man, that the returns require revision, even setting aside the fact of minor importance, but a surprising one, which also appears from these tables, that the main body of the regiment did not furnish so many cases of bowel complaint or of fever, as the companies treated by the experimenter.

3. But even taking the tables, as they stand uncorrected, I do not see how they bear out the assertion of the experimenter, if he means it to be a general one, that he had only one-fifth the mortality of the rest of the regiment, for the mortality in fever and bowel complaints, taking the totals, is set down as 2.8 in the Experimental and 5.6 in the other wards. He therefore had one-half, not one-fifth of their mortality, or even limiting the remark to bowel complaints, he still had about one-third instead of one-fifth the mortality. In either case, he is shown to have been very inaccurate.

It is further evident that in making this statement, he cannot have regarded all his own cases, as cases of dysentery, for in that event he was entitled, according to the tables which give him a mortality of 4 per cent. and the surgeon 42 per cent., to have made a boast of having only one-tenth the mortality of the other portion of the regiment.

It ought to be easy to clear up these strange-looking statements of the Report, by a reference to the regimental books, as I suppose in the Regimental Hospital there was no confusion about transfer cases.

IV. OTHER REGIMENTAL RETURNS.

1. To turn to the regiments generally, for which the abstract of the large table given by the Board is sufficiently accurate, and which has already been given (p. 7), there is the one great error, which runs through all the statements, the setting down diarrhœa and dysentery, separately for the other returns, but in reality classing them together for the experimental ones, for no one can maintain, that a ward open for the treatment of bowel complaints could by any possibility have its cases all dysentery, and no diarrhœa. If the diarrhœa and dysentery cases be added together for the other returns, and then the general result for the whole tables be compared with the Experimental Ward, the superior success of that ward is at once reduced to 1.6 per cent. according to the tables as they stand, and without applying to them any correction.

2. In fever also, the simple addition to the returns of the Experimental Ward, of one overlooked fatal case of fever in the General Hospital,* at once reduces its apparent superiority to ~~22~~²² per cent.

3. But after all, if we examine them in detail, do the uncorrected tables "prove the absolute reality of the superiority of the treatment in the Experimental Ward?" Suppose that we rigidly apply the numerical test, and ascertain whether, in bowel complaints for instance, (as the Board considers the results in them most striking,) it was the most successful. The experimental treatment in H. M.'s 70th gives, according to the uncorrected tables, a mortality of 4.08 per cent.

* Analysis, p. 8.

The Artillery, . . . Benares had 49 cases, no death.

Ditto, Dum-Dum 66 cases, 1 death.

H.M.'s 3rd Drag. Umballa lost 2.04 per cent.

96th, Cawnpore „ 4.3

Artillery, Lahore „ 4.3

Possibly this list might be extended, but this is enough to shew that by the tables themselves, the mortality was less in some regiments, and scarcely greater in other corps, than in the Experimental Ward of H. M.'s 70th. In these cases, was the success attributable to improved modes of treatment? and would it be incorrect to apply to them the remark of the Board concerning the Experimental Ward "that they have indisputably proved that their method of treatment is absolutely superior to that commonly practised?"

4. But it may be said, that, although some regiments in other stations were more successful than the Experimental Ward, still its mortality in bowel complaints must have been much less than the ordinary one among troops similarly situated. Nevertheless, if we turn to the table at the end of the Report, of the mortality among regiments stationed in Fort William during the last twenty years, and therefore, except so far as difference of season is concerned, situated exactly as H. M.'s 70th, we find that in five of these years, 1837, 39, 41, 42, 44, the mortality among them from bowel complaints was less than the 4.08 of the Experimental Ward, and in the years 1830 and 1836 about the same. Nay that even looking on the cases of the Experimental Ward as cases of dysentery, in 1836, 41 and 42, the mortality from dysentery of troops was just the same. Therefore, as compared with the average of years, no one can maintain that there was any particular success in the Experimental Ward of H. M.'s 70th.

5. But, to make the comparison a more general one (and it is in accordance with the principle of the tabular state-

ment, although not in my opinion a correct one), the tables assign to the Experimental Ward (those of the General Hospital and H. M.'s 70th taken together) a general mortality of 4.9, which is at once increased by the addition of one overlooked case in the General Hospital* to 5.21.

The 2nd Europeans lost 5 per cent.

H. M.'s 87th 5.2

H. M.'s 22nd 5.2

H. M.'s 75th 5.3

and therefore were one of them more successful than, and the other two or three as successful as the Experimental Ward.

6 The question may be viewed in still another light. The treatment in H. M.'s 10th and 80th regiments is praised in the Report, as approaching in some measure to that of the Experimental Ward. We should, therefore, applying the numerical test, expect to find the returns for those regiments very favourable, yet on examination we find, that H. M.'s 10th having lost 7 per cent., is behind all the regiments above enumerated, and H. M.'s 80th having lost 8 per cent., is behind them, and also behind H. M.'s 32nd and 24th regiments. This is comparing the results of those regiments for bowel complaints generally, but if the comparison were limited to dysentery, the general results would not, I believe, differ materially.

7. The preceding remarks have been chiefly confined to bowel complaints, but the same numerical analysis might have been applied in a more striking way to fever. Thus a hasty glance shews that at least one dozen of the regiments mentioned in the table were more successful in fever, than the Experimental Ward of H. M.'s 70th, and also that the mortality of troops in Fort William from fever has, on more than one occasion, been less. But a minute enquiry into

* Analysis, p. 8.

these matters would fatigue the patience of most readers, and therefore I shall not enter on it, notwithstanding that the Report says : "The tables shew incontrovertibly that the experimental treatment of fever was more successful than the practice followed generally in European Hospitals in Bengal," a remark still more applicable to the regiments that were more successful than the Experimental Ward.

8. What conclusions then are we warranted in drawing from the preceding analysis? I trust that every one has been convinced, that there must be many very serious inaccuracies in the returns of both the General Hospital and of H. M.'s 70th, and, as respects the "triumphant demonstration," that, whether as compared with the other wards of the General Hospital, or with the Army generally, the Experimental Ward did not present any particular or singular success, even according to the tables, compiled "in the Medical Board Office, to prove the absolute reality of its superiority."

III. ON SOME POINTS OF THEORY AND PRACTICE.

Non obtusa adeo gestamus pectora Pæni :

Nec tam aversus equos Syriâ Sol jungit ab urbe—ÆNEID, Lib I.

THE assiduity and perseverance which the experimenter has displayed, are worthy of all commendation ; but bearing in mind the history of the doctrines and treatment advocated by him, the praise of originality cannot be accorded to him ; and I wonder that he should now have repeated his song of triumph,* and that the Board, though expressly declining to enter upon the question of originality, should, to a certain extent, have recognised it, by the use of the phrase, “demonstrated the perfect safety and wonderful efficacy of quinine on the heroic scale,” an expression scarcely applicable to anything that is not new. Let us now examine some of his statements respecting fever.

FEVER.

(a.) *Cause.*—One of the objects which the experimenter says in his final report he came to Calcutta to accomplish, was

* Report *passim*.—“It has been my glory to be the first to do this.” “In this I am original, and hold my own ground.” “This is my own, all my own, for who has ventured on it before ?” “*My system* contains truths of infinite importance to the health and safety of nearly the whole human race.” “I have come to open new and wide paths of discovery.” “I have done this as it seems to me almost by a *special Providence*.” Had the experiment been less successful, according to his ideas, would he have still appealed to a special Providence ? Such appeals are often very infelicitous, not to say profane. A Surgeon in a local school, addressing his pupils once said—“You thus see, Gentlemen, what Providence, backed by a skilful Surgeon, can effect.” !!

to "re-classify tropical disease," and in an earlier report he announces that, "it will be his endeavour to shew what fevers have malaria for their origin, and what depend on other causes." But of these objects having been attained, there is no indication in the Report. It was the more necessary for him to have set these matters finally at rest, as, according to his views, there would then be no difficulty in knowing when to give quinine, of giving which injudiciously, he remarks, "this accident, (namely, of great mortality in some cases of small-pox and measles,) did occur in my ward." But the discriminating remarks of the Board on this point, make it unnecessary for me to enlarge on the subject.

(b.) *Inflammatory nature.*—1. Treating of the complications of fever, the experimenter remarks—"How astonished will Mr. Martin be to find that all these congestions are not inflammations." Is this a fair representation of the views of the only man, who after leaving these shores has attained distinguished professional eminence at home? Mr. Martin writes—"almost all our complications in the fevers of Bengal are abdominal, whether they be of an inflammatory nature, congestive, or of mere irritation."—*Tropical Diseases*, p. 124. So far was he from expressing an opinion that all these complications were inflammations. The views of my friend Dr. Mackinnon are also strangely distorted, when he is made to assert absolutely "the inflammatory nature of fever"—whereas he expressly says in his very practical work, p. 234, "the other abdominal complications of remittent fever, are rather those of congestion, than of inflammation."

2. The experimenter has ascertained, by examining the records of the General Hospital, that the fatal cases of fever were "cases of extreme congestion, without a trace of the only mark which can be relied on of inflammatory action, viz., the effusion of plastic lymph." But he can scarcely, I

should suppose, have come to Calcutta, expecting to find deposits of plastic lymph. I imagine that none of us are so far behind the progress of medical science, as to regard the lesions of fever as simply those of acute inflammation. But supposing that these were discoveries at the General Hospital, that this was new to us in India, and that deaths from fever, with no *post mortem* appearances at all sufficient to account for death, were of unusual occurrence,* supposing that these truths had never before flashed on the mind of any one in India, but the experimenter, is he aware that he is merely repeating the established doctrines of Europe? The late Dr. R. Williams, a very intelligent writer, says in his *Elementary Principles of Medicine* :—"As a general rule in all tropical fevers, the traces of diseased structure are always trifling," and in this he merely agrees with the French, who of late years have become far more familiar with remittents than the English, from the intimate connexion of France with Northern Africa. Grisolles, (who says expressly, that quinine is the only specific for remittent fever,) in his popular manual, which had reached a 4th edition in 1850, treating of the management of what the French call "pernicious intermittent fevers," says—"We are fully persuaded that the acute pains and the extreme derangement of the organic functions which characterise them, are not allied to inflammation," and of the pathology of remittent, he writes—"Some have ascribed the origin of remittent fever to visceral irritation or inflammation, but the researches of pathological anatomy do not bear out this opinion—for the lesions of the stomach are not sufficient to account for the general symptoms, and no one would attribute the functional disturbance of the system to any alteration of the liver, for its condition

* I gave the results of some such *post mortems* in the Medical Gazette, for 1841.

most certainly is no more of inflammatory origin, than is engorgement of the spleen. In fine, the researches of Leonard and Folley, in 1845, on the composition of the blood in fever, exclude the idea of inflammatory action."

(c.) *Use of Quinine.*—But physicians in India and in all parts of the world have long shown, by their treatment of fever, their practical disbelief of the inflammatory nature of these congestions. They have been gradually abandoning general blood-letting, and using quinine more freely. To vary my authority, and quote a popular German handbook, "Very generally," says Oesterlen, "preparatory measures delay the cure quite unnecessarily, as the quinine is often the best means to remove various disturbances along with the fever; especially, do not be deterred by some vague symptoms, as the state of the tongue, or swelling of the spleen or liver, from using quinine at the earliest stage, if the fever be a bad one." In the West Indies among H. M.'s troops as early as 1832, Dr. Binns always observed the practice of giving quinine in the early stage of fever:* and it is hardly necessary to say, that besides its not infrequent use in this way for many years in the East Indies, including Ceylon and China,† a whole host of French and American writers have, during the last 15 or 16 years, been giving accounts of the exhibition of quinine in immense doses, regardless of the presence of congestions. As an instance of this change of practice, I may quote Celle, *Hygiène Pratique*, 1848. He practises in Mexico, and says, of what we should call jungle fever,—“Where we used to purge and bleed to the great detriment of the patient, a few doses of quinine effect a cure,”—and in 1847, Dr.

* *Lancet*, 1846.

† The reader will find some account of the history of the use of quinine in India in remittent in “*Bengal Dysentery*, 1850.”

Bartlett, the American author of a work on fevers, lays it down, that quinine is to be given at once, without waiting to remove congestions—adding, the quinine removes them; or to quote Oesterlen again—“In conditions of excitement and congestion, even in inflammation of various parts, so soon as they show an intermittent or even a distinct remittent character, quinine is of great service.” Exactly the doctrine laid down by the experimenter as something new, that if “quinine be given freely, these imaginary inflammations vanish.”

So general, however, do I believe the opinion in India to be, that the complications of fever are not simply inflammatory, and so generally diffused is the knowledge that the quinzation treatment,* *i. e.*, by very large doses of quinine, is one effective method of treating tropical fevers, that I should scarcely have thought it necessary to cite the preceding authorities, had not the proposal of the Report, on the treatment of fevers, been considered by some, a more decided step than had ever been previously taken by any systematic writer.

The question of the expediency of adopting that or any other indiscriminate mode of treatment, is not now under consideration; but before leaving the subject, it may be remarked, that the experimenter in his former *brochures*, so far as my memory serves me, appeared to be quite as well satisfied with the effects of 8 or 10 grain doses, as he now is with scruples, and that Dr. Dundas, the latest English writer on the quinine treatment of fevers, is confident of the success of 10 grain doses in intermittent, remittent and continued fever equally; and that in continued fever with the aid of general support, it seems already to have been found efficacious, like the old bark and wine treatment.†

* I prefer the word quinzation to cinchonism, or quininism.

† See Dublin Journal, 1852.

(d). *Mixed Treatment*.—The experimenter says—"His fevers were not cured by quinine mixed with calomel and purgatives, but by quinine alone;" again, "as for other remedies to assist, I have used none of them, blisters, leeches, purgatives." Now without criticising in detail his fever cases, I have looked over the Hospital records, in which his treatment is registered, and I think that if he had an opportunity of referring to the records kept by himself, he would be inclined to modify the statements quoted above, for he has noted in his own handwriting the use both of venesection and of local bleeding; *haustus purgans* occurs constantly in his journals; nor are *Pulv. Jalap. C.* and *Ol. Ricini* absent—and he also used *haustus diaphoreticus* frequently, apparently when he suspected the fever to be eruptive.

It is, however, practically speaking, a matter of no importance, that in fever the experiment was so mixed a one—as it was well known, before the experiment was made, in short as completely ascertained as any fact in medicine, that fever might be successfully treated with large doses of quinine, without bleeding, purgatives, or even aperients being thought necessary.

(e.) *Arsenic*.—This is a very old remedy, which was used by the Chinese and Hindoos in fever. There is a very natural prejudice against the use of so active a poison, but in practice its careful employment has generally been considered quite safe. The experimenter tells us gravely "that all experiments made on the treatment of skin diseases, show it to be innocuous." Very true, and besides its familiar use in cutaneous diseases, besides its having been a popular remedy in fever for the last three quarters of a century at least, in England and in Germany, its use was revived in France about 10 years ago, and since then it has been employed most extensively and successfully in that country, in Italy, Africa and the Brazils. What

does Masselot* say of the treatment of bad remittent?—"In the present state of our knowledge, I should give quinine, if I had to choose between it and arsenic. But if I had both remedies at my command, I should give the two together." But we Indians, who are said in the Report "never to use it," are not entirely ignorant of its virtues, and have employed it, seemingly unconscious that we were doing anything much out of the way, when the supply of quinine ran short, or when quinine failed to stop the fever.†

The experimenter also informs us that "he has good reason to think that arsenic gives immunity from future attacks of fever." This is just what Boudin, Maillot and other writers say, and mention as one of the particular advantages of its exhibition. Again "now all this, novelty as it is, is no conjecture." It is undoubtedly no conjecture, in fact no more conjecture than novelty, as hundreds of practitioners can testify, as was shown in the Hospitals of Brussels after Waterloo, and as it has been of late years proved on the large scale in France, when Boudin found it succeed in 2,947 cases of fever, 2,000 at least of whom had been previously treated with quinine. I need not enter here on the question of its proving an efficient substitute for quinine.‡

(f.) *Statistics.*—Before leaving the subject of fever, it is agreeable to be able to say, that all the tables of the Report, (that of all the European troops in Bengal for 20 years from

* Archives de Medicine, 1846.

† In 1841 I alluded in the Medical Gazette to its successful and safe employment.

‡ I would venture here to throw out a hint for the experimenter's future investigations. He has tried fever successfully by the quinine and by the arsenic test. He has tried dysentery, he considers successfully, by the quinine test. The arsenic one should, according to *his theory*, be next applied: if it succeeds as in fever, he may be able to banish quinine from India, and possibly even the long tube also.

the large number of facts it represents, is a valuable one,) agree in showing, that though the percentage to strength of cases of fever has not varied much, a distinct diminution of mortality in fever has taken place. The percentage of mortality in the 20 years ending with 1849, taken in periods of 4 years, has been,

General Hospital.	Garrison of Fort William.	All European Troops in Bengal.
9·5	4·5	3·06
8·56	3·1	2·52
10·5	2·2	2·40
5·4	3·1	2·32
6·5	3·4	2·07

In all these columns it is to be observed, that there is a smaller mortality in the last 4, than in the first 4 years of the period, but it is also apparent, that it is only on the large scale of the whole European Army in Bengal, yielding for those 20 years an average of nearly 9,000 cases annually, that the improvement is at all uniform. No one, I believe, will doubt that this gradual change, this diminution of mortality by one-third, is owing to the less free use of depletion, and the freer employment of quinine.

DYSENTERY.

(a.) *Quinine*.—1. The experimenter “has proved the marvellous effects of quinine in hæmorrhagic dysentery.” “This is my own, all my own, for who has ventured on it before?” Unfortunately for these pretensions, the idea that because dysentery is probably a malarious disease, therefore it should be cured by quinine, has long been entertained. But the efficacy of that remedy in this disease is only partially established. Dr. R. Williams writes “So uncertain are the laws of paludal poison, that as a general rule in dysentery, the

exhibition of quinine in any form or quantity has proved rather injurious than otherwise." I am happy to say that this is erroneous, for it has been for many years given with advantage in small doses in chronic dysentery at the General Hospital, and no doubt elsewhere,—and the present Report alludes to its frequent use in hæmorrhagic dysentery in H. M.'s 80th. And as to the statement, that no one has ventured on it before, "*pereant qui ante nos, nostra dixerint*," here again the pertinacious Frenchman, envious of the glory of *perfidè Albion*, has been beforehand with the experimenter. "Sulphate of quinine," says Grisolles, "is indicated, where dysentery shows the character of paludal fever. In such cases I have seen its heroic use as successful as in bad intermittent."

2. Great stress is laid on this successful treatment of dysentery by quinine. The Report says, "It is directly a practical matter of fact, and cannot be answered." "I must have had increased mortality if *my* theory is wrong, whereas I have had far less." Now, waiving the question whether his treatment must have had the effect he supposes, I find myself, to my great surprise, in a position that enables me to tell him, from his own books, that he has misapprehended the matter, that it is *not* a matter of fact, and that there *was* increased mortality in the General Hospital in his own hands, under that mode of treatment. The experimenter explains that by hæmorrhagic dysentery, he means "all forms of it in which much blood is voided by stool—not only the worst forms of it:" his statement therefore may be considered to be of pretty general application to Bengal dysentery.

3. The statement that scruple doses cure hæmorrhagic dysentery being a very important one, and such experiments as I had myself seen made with it being failures, I examined the books of the Experimental Ward, to see how its success-

ful results had been obtained, and the following results of the examination of them, are as near an approach to the truth, as I am able to make.

There were treated without quinine about 30 cases of dysentery; of these 3 died, 1 with hepatitis and diarrhoea, and 1 with hæmorrhage; 27 were discharged, of these 5 were re-admitted. There were about 32 cases of dysentery treated freely with quinine; of these 6 died in the Experimental Ward, one of them of hepatitis; and there were transferred to the other wards 5 cases, (2 of them fatal.) There were discharged, 13 absolutely cured, 4 convalescent, 4 well, but who had passed blood within two or three days of their discharge:* or to view the matter generally, among 30 cases treated without quinine, there were 3 deaths; among 32 cases treated with it, there were 6 deaths, or the mortality was actually increased during the quinine treatment,—I do not say by it.

The reader ought also to know, that at the same time there were some 23 cases of diarrhoea treated freely with quinine, of whom 21 were cured, 1 died of cholera, and 1 was transferred: about 12 cases of diarrhoea appear to have been treated without quinine, and with no deaths.

Or to take a still more general view of the whole case, 55 cases of bowel complaint treated with quinine were attended with 8 deaths, (excluding one of cholera,) and 42 cases treated without it, with 3 deaths.

* Patients may be discharged in this state from any of the wards. If a sailor is convalescent, and his Captain wishes to have him, on leaving port, it is often necessary to discharge him. The state in which many of the dysentery cases were discharged, according to the books, has some bearing on the vaunted quick cures of the Experimental Ward: but it is unnecessary to open up a question, on which, from the nature of the experiment, it has been seen that there can be no distinct facts.

(b.) *Mixed treatment.*—As in his remarks on his treatment of fever, so in those on the management of dysentery, the experimenter seems to have expressed himself carelessly: he says, “I have used no other remedies but washing out the bowels with water, and giving 6 or 7 scruples of quinine daily”: and this is the impression which the whole Report conveys, for though in one place he says, “for the acute form of dysentery bleeding is decidedly necessary,” and “a full dose of laudanum will be found of the greatest benefit,” he no where tells us, how to distinguish the paludal from the acute form, and this statement is soon forgotten, among such general expressions as these, “leeches, blisters, and ipecacuanha I know nothing of;” “neglecting all these, bleeding, &c.” The tabular returns also make no distinctions of bowel complaints, according to their treatment.

From the comparison that has been made in the preceding section, the inaccuracy of these statements might be inferred, and the books also shew, that besides using venesection more freely than it was practised in the other wards, the experimenter employed occasional leeching, and very frequently opium, morphia, nitrate of silver, acetate of lead, all in large doses, and chalk mixture. Are these remedies to be looked on as part of the treatment or not?—for, although in an early part of the Report, he says “he may show the effects of alam and such like harmless palliatives,” we do not afterwards hear anything more on the subject. If any positive conclusion can be drawn from so small a number of facts, and I am very far from supposing that it would be safe to do so, it follows from those adduced in the last section, that the quinine treatment was less successful than the other, both in dysentery and in bowel complaints generally. It is therefore to be regretted, that the experimenter did not analyze more carefully his own cases, before making such sweeping state-

ments, as to the efficacy of quinine, and the non-employment of other remedies.

(c.) *The Long Tube*.—The earlier part of the Report and the experimenter's previous *brochures*, are full of the long tube and of its virtues. In former times, and before he had discovered the use of quinine in dysentery, the long tube and doses of laudanum were his panaceas, and were talked of, perhaps more confidently than his more recent mode of treatment. He paid his professional brethren the compliment of saying that "he would cure hundreds of the slighter cases, that perish under the present treatment." In those days he expressed his belief in the common existence of scybala; he then seriously expressed his opinion, that, as the conjunctiva and the lining of the intestines were both mucous membranes, the application of fœcal matter to the inner surface of the bowels must be as irritating to it, as it would be to the conjunctiva of the eye. He also compared his large injections to the treatment of open abscesses, and said that his treatment was in fact turning the internal abscess of the colon into an external one.* But although he says "he need not repeat his already published opinions of the theory of acute and chronic dysentery," more extended experience seems to have made him practically drop those false analogies, as we hear nothing more of them. He now limits himself to the remark, "that you must wash out all the irritating matters which the large bowel contains, and which are the cause

* Mr. Hare, on Dysentery, pp. 8, 9, 10, 11, 21, 30, 23; in Bengal Dysentery, p. 50,—it has been stated that "scybala or accumulations of fœces appear to have been scarcely ever observed during life, and never after death." This as it stands, is a general expression of the truth, and a stray instance or two to the contrary serve merely to confirm the fact. The statement would however have been more exact, if the expression used had been, as was intended, "accumulations of (hardened) fœces." This explanation was necessary, as many people attach a very vague meaning to the term scybala.

of its sloughing." Passing by this opinion as to the cause of sloughing,* which we leave it to him to reconcile with the paludal theory of dysentery, we observe, that towards the close of the Report, he speaks less urgently of the long tube, and says "it need not be passed up oftener than once in eight or nine cases," "although he always prefers using it himself," and says "the necessity of injections has been still more impressed on him." We also hear less of the spasms or narrowing of the sigmoid flexure, on the existence of which the use of the introduction of the long tube mainly depends.

In fact, notwithstanding the Board's condemnation of those who do not carry out the experimenter's mode of administering enemata, the Report to my mind gives the impression that quinine is much more efficacious than the long tube. It is a pity that the Report has not been more explicit on this head, that the use of that instrument has not apparently been studied, apart from other remedies. No one can collect from the Report, how often

* Notwithstanding the opinion expressed by an intelligent reviewer in a late number of the British and Foreign, it still appears to me, that the ordinary process in acute Bengal dysentery is more one of infiltration, death and sloughing of the mucous coats, than one of regular progressive ulceration of the solitary glands, as described by Parkes and Murray. There is scarcely time for the latter process. The case quoted by the reviewer from "Bengal Dysentery," as probably giving a fair average representation of the usual appearances in Bengal dysentery, was quite an exceptional one, although I have seen similar ones.

The descriptions of the alterations of the bowels in dysentery given by Rokitansky, the Irish writers in the Dublin Journal, and by Dr. Baly, agree in all their main features with those of Bengal dysentery, the degree of liver affection alone excepted. Rokitansky's ulcerative inflammation of the follicles brought on by diarrhœa, appears to correspond with the dysentery of Parkes. It is a curious fact that in the earlier months of this year more than half the fatal cases of dysentery in H. M.'s 80th were attended with abscess of the liver, while in the neighbouring Hospital abscess was not common. I have never heard of liver abscess in children connected with dysentery.

it was passed up beyond the sigmoid flexure, how much was actually attributable to the use of the long tube, and how much to the other remedies which were applied.

In short, the Report appears to me calculated rather to advance the cause of quinine than that of the long tube, and it is curious to observe, how the present state of things in India agrees with that in Europe, as represented in the Library of Medicine, 12 years ago—"Dr. O'Beirne is anxious to induce his professional brethren to make use of the introduction of his favorite tube into the sigmoid flexure, believing that much mischief results from the detention of fœcal matter and of diseased secretions in the large intestine, mainly caused by a spasmodic closure of the upper annules of the sigmoid." Dr. O'Beirne, however, as far as I can judge from books, has, like the experimenter, scarcely remained true to his first convictions, for he now urgently recommends tobacco, but in the form of fomentations, therein at once agreeing with, and differing from Matthews, who also administered tobacco, but by means of a hookah snake, which seems to have been the great prototype of the long tube. For myself, after giving large injections a fair trial, I cannot assign to them a higher place in Bengal dysentery than that of frequently useful adjuvants: and however useful they may be in certain conditions of the bowels, as in constipation, or in the incipient stage of dysentery in the Upper Provinces, the general result of what I have heard from experienced practitioners, who have given them a fair trial in European Hospitals—(I believe I may mention particularly that of H.M.'s 80th, whose Surgeon is so favorably mentioned by the Board) agrees with the opinion of them that has been just expressed.

"Quin si morbus diutius excurrat, frustra erit (me iudice) qui medicari sataget, vel quavis methodo prædicta, vel clys-

teribus, abstergentibus, agglutinantibus, atque astringentibus, quæ pro variis ulceris injici solent.”—*Sydenham Obs. Med.*, iv. 3, 19. And this, I fear, is every practical man’s view of the case to this day.

(d.) *Calomel Treatment of Dysentery*.—1. As the mercurial treatment is not a recommendation of the report, it is unnecessary to allude at much length to this subject; still I shall examine one or two statements of the Report on this head. The experimenter, who in a former *brochure* spoke “of the dreadful results produced by the system of salivation, which is still the common routine in India,” now expresses himself thus: “The mortality is proved by these tables to have increased by regular progression, and the case books also show that the number of cases salivated has also decreased in the same proportion, till now, when salivation is never resorted to, the mortality is actually doubled.” “How greatly is the demonstration strengthened, when the same fact occurs in the Regimental Hospital with a class of patients entirely distinct in every circumstance and habit from the former.” It may possibly be thought a sufficient reply to this, to quote the words of the experimenter at the close of the Report. Alluding to modern practice he says, “The mortality has increased to what it was when scruple doses of calomel were in use.” But as the experimenter has just stated, that since salivation was given up, the the mortality has been doubled, and now says that under either system it is the same, what are we to regard as his real opinion on the subject?

2. However, leaving him to reconcile those warring opinions, I proceed to the facts of the case, and have compiled from the experimenter’s own tables, the percentage of mortality from bowel complaints for the last 20 years, in periods of 4 years, for here as elsewhere diarrhœa and dysen-

tery classed together, will probably give the most trustworthy general results.*

	Year 1830-35.	1834-37.	1838-41.	1842-45.	1846-49.
Genl. Hospital,	10·6	19·6	17·8	17·7	18·1
Garrison,	9·6	6·1	5·3	6·1	6·6

Can any one trace a regular progression in these figures? So far from this, there appears to have been no change of any importance for the last 16 years. Again, do they show a progressive common deterioration? So far from this, we find, that the Regimental Hospital has been much more successful in the last 4 than in the first 4 years of the period, whereas it is exactly the reverse with the General Hospital.† But even if the returns had corresponded, the very fact of their doing so, while the Regiments in Garrison were constantly changing, each new Surgeon bringing his own particular practice with him, would have been sufficient to my mind to

* As some may wish to see how the figures stand for dysentery alone, I give them also:

	1830-33.	1834-37.	1838-41.	1842-45.	1846-49.	
Genl. Hospl.,	15·3	25	21·2	25·6	27·6	{ or accord- ing to my returns, }
Garrison, ...	11·1	6	6	12	16·1	
						26·4

In these figures can any one trace a gradual regular progression, or a correspondence between the two series, such as can be attributed to a progressive change of practice? So far from the mortality being regularly progressive, it was 14·8 in 1833 and 26· in 1834 in the General Hospital.

† Of these anomalous facts as they stand, I can offer no satisfactory explanation: probably, if any accurate data for the years immediately preceding 30, could be got, it might be possible to explain them. How Raleigh, writing in 1842, called the mortality 10 to 14 per cent. I don't know; in the eight previous years it was 22 per cent. Martin's tables make it 16·2. Raleigh says, that at the end of the last century, when calomel with large doses of opium and mercurial frictions was the most common practice, it was from 30 to 50, nay 75 or 80 per cent. It is much to be regretted that the labours of the white ants make it impossible to get any accurate account of the mortality in former years.

shew, that their accordance depended on some more general influences, than those of treatment.

3. Ere leaving this subject, it is curious to look back at the last experiment at the General Hospital, that of 1816. In those days Dr. Halliday, shocked at the abuse of calomel, of which he says 13,237 grains were given in the General Hospital in one month, thought that he had discovered a more successful mode of treating dysentery than by calomel. He thus compares his own, or the non-calomel treatment, with that of his colleague, Mr. Wood, or the calomel practice :

Dr. Halliday had 76 cases dysentery,	4 deaths,	mortality 5·2 per cent.
Mr. Wood „ 24 „ „ 9 „ „ 37 per cent.		

We find therefore that the non-mercurial treatment of 5·2 per cent. was more successful than that of the late Experimental Ward of the General Hospital, which was, according to the official report, 7·3, but that the mercurial treatment of Mr. Wood, with a mortality of 37 per cent., was scarcely so unfortunate as that of the Surgeon of H. M.'s 70th Regiment, who lost 42 per cent. These comparisons may appear somewhat ludicrous, but they are not in reality juxtapositions of more dissimilar things, or of more inaccurate data, than those which are classed together in the general tabular statement of the Report, and they afford additional illustrations of the extraordinary results that may be obtained by the comparison of figured statements, when not carefully conducted.

(e.) *General Statistics of Dysentery.*—But though, for reasons above stated, I cannot assent to the hastily assumed generalizations of the experimenter, it is a fact that there has been no progressive diminution of mortality in bowel complaints. On the contrary, it will be seen from the returns of the whole European army in Bengal, which for the last 20 years has furnished an annual average of 4,011 cases

of diarrhoea and dysentery, that the percentage is somewhat greater than it was 16 years ago, greater by about 1-6th.

1830-33.	1834-37.	1838-41.	1842-45.	1846-49.
5.04	4.99	6.63	6.3	6.2

It is extremely unsafe to generalise in statistics, but I cannot help remarking that 1838-41, the first period of increase, and the period of greatest mortality, comprehends the Affghan and China campaigns. It could also scarcely be expected, that there would be a diminution of mortality, as the troops have been much more engaged in campaigns, and in occupying new stations, during the last 12 years, than during the first 8 of the period: besides, there has been no important improvement made in the treatment of dysentery, at all events extended enough to affect general returns, as there has undoubtedly been in fever.

It is also well worthy of observation, that from the general returns for the whole European army, if we compare the first 4 years of the period of 20 years, with the 4 last of it, it appears, that the proportion to strength of cases of bowel complaint has considerably increased, that of diarrhoea having increased greatly, and that of dysentery having rather diminished, while, as we have already seen, the proportion of fevers to strength has remained nearly stationary.

QUININE—ITS SAFETY?

THE experimenter states, that "he has never seen any harm result" from reducing a patient to a state of quinization, and the Board re-echo "its perfect safety." This is the frequent testimony of writers who have used the drug freely. Bally, for instance, mentions giving 15-grain doses 4 times in the 24 hours, without the slightest inconvenience being produced, on which Trousseau and Pidoux remark, that either Bally observed his patients carelessly, or that his patients deceived him. But are there no exceptions to this rule? Are there not, besides many cases of more or less permanent deafness and blindness, several instances of still more disagreeable effects on record?* and have the French experiments in the treatment of rheumatism with it been forgotten? In Florida, in particular, where the system of monster doses has for some years been the routine in fever, disastrous results have not infrequently followed its indiscriminate exhibition, especially in children. The most commonly recorded effects on man, are amaurosis, deafness, delirium, muscular tremor and coma, and recent experiments on animals mention as some of the symptoms that precede death, great restlessness, speedily followed by muscular agitation or tremulous movements. To small animals it readily proves

* Piorry is said to have lost 6 cases of intermittent fever, and the other Paris Hospitals at least 12 more from its injudicious employment, and the same has happened to Giacomini and others in Italy. *Gazette des Hôpitaux*, No. 123, 1847. Alibert and Recamier have seen deaths caused by 15 grain doses. Oesterlen remarks, that he has always used quinine in moderate doses with perfect safety, but that its employment in a Parisian Hospital seems to be quite another, and a dangerous thing.

fatal, but only rarely to man. The experimenter mentions the frequent production of amaurosis and deafness, were they never the preludes to the further effects of the drug?

He says the man who died of fever in H. M.'s 70th "had been drinking, and had strong symptoms of delirium tremens." Does this mean that the man died of fever or of delirium tremens? and if of the latter, has he never observed quinine induce a state of tremor, a *delirium é quind*? strongly resembling *delirium é potu*, or at least has he not seen it aggravate an already existing tendency to delirium tremens?

One remarkable case was transferred from the Experimental to the other wards, as a case of delirium tremens, — who presented the symptoms of complete quinization, with deafness, tremor and coma. The patient died. In this case, after death the coats of the stomach were found pale, not even reddened, or with any abrasions of the mucous membrane, such as are almost always found in *delirium é potu*, which occurs usually in old drunkards with more or less diseased stomachs. This was a rather startling case, but there were some other striking incidents that deserve a short notice.

It is somewhat remarkable that in a ward devoted to fever and bowel complaints, two cases, both fatal, of ebrietas should have occurred, and also two fatal cases of epilepsy and of apoplexy, (of course there is nothing surprising in the case of apoplexy, if it was merely the termination of a fever, but, although the case was admitted as fever, it does not appear as such in the returns.) I also think it surprising, that of six cases of delirium tremens, which occurred in that ward, and which were treated more or less freely with quinine,* (for the two cases of ebrietas, I suppose must be counted as deli-

* All of these patients had taken 1 or more scruple doses.

- rium tremens,) no fewer than five terminated fatally in one or other of the wards.

The suspicion that quinine might have acted injuriously, has evidently crossed the mind of the experimenter, for in the case of apoplexy, just alluded to, he expressly explains, that the patient had stopped taking quinine, "as he carelessly thought him convalescent." But when does he seem to have stopped taking it? "This very stout man, who does not seem very bad," appears to have been taking from the 10th to the 12th, 4 scruples of quinine daily, and this was only stopped the evening before his sudden death on the 13th. This at least so far as the Hospital records show. The experimenter candidly admits the bad effects of large doses of quinine in eruptive fevers, the good effects of which in smaller doses many authors record; possibly on reflection he may admit, that it may also be injurious, where there is a tendency to delirium tremens, or other cerebral disturbance.

We have already seen that the total mortality among all cases that were treated at all in the Experimental Ward, exceeded that of the other wards, and until the cause of that mortality is satisfactorily explained, the effect of the incautious use of quinine offers itself as a possible explanation of it: and the possibility of this being the true solution, is strengthened by the fact, that the bowel cases admitted into the Experimental Ward, and treated with quinine, yielded a mortality of 14.5 per cent., while those treated without it gave only one of 7.1 per cent. (as almost all fevers were treated with quinine, it is only possible to make this comparison in bowel complaints.)

We have also just seen reason to suspect, that quinine proved deleterious in delirium tremens, and in tendency to cerebral congestion, as well as in eruptive fevers.

All this is matter for grave consideration, but I would

avoid hasty generalization from a small number of facts, and should hardly have felt called on to say so much on the subject, had not the Medical Board, notwithstanding that they had previously given a caution on the subject, "deploring the *possibility* of the occurrence of serious consequences," which in all probability had *actually* occurred under their eyes, taken this opportunity of assuring the profession, of the "perfect safety of exhibiting quinine in large or scruple doses repeated three, four, six times or oftener in the 24 hours, in fevers of all types, the product of the poison of malaria."

ἔμοι μὲν οὐ πιστὰ λέγοντες, ἄλλω δέ τινι.—HERODOT.

It is impossible, even with the limitation, which they have added, that such indiscriminate practice should not continue, as it has done heretofore, occasionally to produce disastrous effects, as it constantly will, disagreeable ones. They seem to think nothing of driving a patient distracted, or half mad (as he usually expresses himself) by singing in the ears, irritability of the stomach, deafness and partial blindness,* and this while there is every reason to believe, that smaller doses will produce every effect of the drug that we wish to obtain.

Though not agreeing with them in their opinion of its perfect safety, I would, however, go so far with them, and say, that it is certainly a very difficult thing to give quinine to an adult during a single attack of fever in permanently injurious doses, for large doses, such as they mention, make the head so uncomfortable that the patient will no longer take them, or if he does take them, they produce such irritability of stomach that each dose is usually rejected.

* In some cases in the General Hospital, patients have been treated expressly for the effects of quinzation.

CONCLUSION. .

BUT these remarks must draw to a close. After all that has been said, I trust that every one who has perused these pages, must feel satisfied, that the experiment was not conducted with sufficient care; that the tables are inaccurate, and do not bear out the interpretation of triumphant success, which has been put on them : that the treatment was a mixed one, not admitting of any accurate conclusions being drawn regarding the effects of individual remedies, or if, of any, of conclusions regarding bowel complaint, directly the reverse of those of the experimenter, and finally, that there is no originality in the views put forth in the Report.

In short a very distinct reply has been given to a question which was propounded by the Medical Board, in the year 1816, when Dr. Halliday's experiment was under consideration—"Was there any novel mode of practice introduced, was the nature of that practice beneficial, and such as to render its general adoption desirable"?*

What then, it may be asked, has the experimenter effected nothing?—have his enthusiasm and industry been thrown away? By no means: the agitation of the question of the employment of quinine has given a considerable impulse to its use in Bengal, and in so far it has done good. It is far better to give quinine too early than too late, too

* The reply given on that occasion was, that "Dr. Halliday's patients were discharged, often not cured, or even convalescent." In the present instance, there can be no distinct facts on this head, regarding any of the Wards of the General Hospital. The Official returns we have seen, afford a strong presumption, that the cases in the Experimental Ward of H. M.'s 70th must have been slight, because the portion of the regiment, treated by the experimenter, produced a larger number of cases, than the main body of it.

much of it rather than too little. It is a remedy which almost any one can give with safety, for, although I have seen with disapproval mothers giving their children 4-grain doses of it in dentition, and I have known patients, in dread of fever, often make themselves very uncomfortable with it, and seen a whole series of symptoms of deranged digestion and nervous disturbance follow its continued abuse, there is little risk of its being generally given in the monster doses of the Experimental Ward.

I would fain hope also, that there may be some foundation for the belief in the high efficacy of quinine in hæmorrhagic dysentery, for as yet I know of no medicine, that really makes an impression on its bad forms, and indeed a knowledge of the structural changes that take place, scarcely encourages the hope, that one will be discovered.

Next, as to the long tube, though that instrument is not likely to be often used, attention may have been awakened among those who were not familiar with their employment, to the use of enemata, remedies by no means to be neglected in the treatment of dysentery.

The experimenter also deserves credit for the industry with which he scrutinized the records of the General Hospital, and he has shown considerable ingenuity in the way in which he has endeavoured to support his theoretical opinions by numerical statements.

Before closing these observations, I must apologise to the reader for having in the last part of them, perhaps brought the experimenter too prominently forward, and so frequently examined his crude speculations, especially as I know that in this country *his system* has very generally been silently laughed at. My apology for this must be, that the Report consists mainly of his opinions, and that it was hardly possible to deal with the one apart from the others.