


GENERAL ZOOLOGY
or
SYSTEMATIC NATURAL HISTORY

by
GEORGE SHAW M.D.F.R.S &c.

WITH PLATES

from the first Authorities and most select specimens

Engraved principally by
MR HEATH.




VOL. IV. Part 2.

P I S C E S .

London Printed for G. Kearsley Fleet Street.

1803.



GENERAL ZOOLOGY.

VOLUME IV.—PART II.

PISCES.

LONDON.

PRINTED FOR GEORGE KEARSLEY, FLEET-STREET;
BY THOMAS DAVISON, WHITE-FRIARS.

1803.

XIII f. 13

CONTENTS

OF

VOL. IV.—PART II.

	Page		Page
ACANTHURUS GENUS		Band-Fish, rough-finned	191
.	374	Hermannian	191
Unicorn	374		
round-fronted		BODIANUS GENUS	569
.	376		
blue	377	Bodian, purple-backed	569
lanceet	379	five-spined	570
blackish	380	Aya	570
military	380	large-scaled	571
triradiated	381	spotted	571
guarded	381	Louti	572
violet-striped		palpebral	572
.	382	silvery	573
dusky	383	Apua	573
Achillean	383	guttated	574
lineated	384	Zebr	574
umbre	384	Rogas	575
speckled	385	lunated	575
broad finned	385	black-and-white	576
		star-eyed	577
Band-Fish, common	187		
rubescens	188	Bull-Head, river	254

Bull-Head, grunting	255	Chætodon, Butterfly	339
———— Lasher	257	———— chalybean	340
———— four-horned	259	———— arcuated	341
———— rough	269	———— bifasciated	342
———— insidious	260	———— horned	343
———— Marseilles	261	———— Bat	344
———— Madagascar	261	———— long-finned	345
———— black	262	———— red-striped	346
———— Southern	263	———— sickle	347
———— mailed	263	———— Charioteer	348
———— Japanese	264	———— Aruan	348
———— single-spined	265	———— Chinese	349
 CEPOLA GENUS	187	———— spotted	350
 CHÆTODON GENUS	323	———— biaculeated	351
———— imperial	324	———— golden-side	351
———— Paru	325	———— lunulated	352
———— Angel	325	———— golden	353
———— ciliated	326	———— Asfur	353
———— fasciated	327	———— geometrical	354
———— Surat	327	———— trifasciated	355
———— Kleinian	328	———— great-finned	355
———— eight-banded	329	———— Chili	356
———— annular	330	———— pleat-nose	356
———— collared	331	———— glaucous	357
———— black-and-white	331	———— rhomboid	358
———— Argus	332	———— bordered	359
———— bimaculated	333	———— tricolor	360
———— striped	334	———— speckled	361
———— divided	334	———— Peacock	362
———— single-spotted	335	———— Curassoa	362
———— wandering	336	———— Bengal	363
———— ocellated	337	———— rock	364
———— rostrated	337	———— Mauritian	364
———— Orb	339	———— punctated	365
		———— round	366
		———— single-spotted	366
		———— orbicular	367

CONTENTS.

v

Chætodon, silver	367	Dory, Brazilian	281
———— long-snouted	368	———— Indian	282
———— Boddaertian	368	———— ciliated	283
———— Sargoid	369	———— inædulous	284
———— canescent	369	———— common	285
———— scaleless	370	———— Opah	287
———— sordid	370	———— square	291
———— Lamark's	371	———— red	292
———— Plumier's	371		
———— constricted	372	Flounder, Holibut	295
———— armed	373	———— smaller	296
		———— Plaise	297
CORYPHÆNA GENUS	212	———— Dab	298
		———— Smear Dab	299
Coryphene, common	212	———— long-Dab	300
———— Brazilian	214	———— Flounder	301
———— Plumier's	215	———— rose-coloured	302
———— blue	216	———— Sole	302
———— Razor	217	———— Venetian	304
———— five-spotted	218	———— Pallasian	305
———— gilt-tail	219	———— Zebra	306
———— striped	220	———— slender-finned	306
———— Parrot	221	———— Carolina Flounder	306
———— Mackrel	221	————	306
———— undulated	223	———— ocellated	307
———— sailing	224	———— Rondeletian	307
———— lineated	224	———— platesoid	308
———— Japanese	225	———— silver	308
———— greenish	225	———— smooth	309
———— Chinese	226	———— bearded	309
———— flat-snouted	226	———— marbled	310
———— sharp-tailed	227	———— pavonian	310
———— half-finned	227	———— lineated	311
———— branchiostegous	227	———— Turbot	312
————	227	———— Pearl	315
———— clypeated	228	———— Whiff	316
		———— bilineated	317
COTTUS GENUS	254	———— ornamented	317

Flounder, large-scaled	318	Gobiomora, Southern	249
— sinistrous	318	— sleeping	250
— Argus	319	— Gronovian	251
— stellated	321	— Kolreuterian	251
— Carolina	321	— Pisonian	252
— Japanese	322	— great-headed	253
GASTEROSTEUS GENUS	60	GOMPHOSUS GENUS	479
GONIUS GENUS	232	— blue	479
		— variegated	480
Goby, common	232	Gurnard, grey	619
— lance-tailed	233	— Piper	620
— Boscian	234	— Cuckow	620
— blue	235	— Asiatic	621
— Bronssonet's	236	— sapphirine	621
— spotted	236	— flying	622
— blue-finned	237	— springing	622
— Boddaertian	238	— Lucerna	623
— hare-lipped	239	— Carolina	624
— Paganel	240	— Japanese	624
— red-fronted	240	— minute	625
— Ouheitee	241	— Cavillone	625
— Arabian	242	— punctated	626
— black	243	— lineated	626
— black-finned	243	— pine	627
— clouded	244	— mailed	628
— Chinese	245	— Chabrontera	629
— Plumier's	246	GYMNETRUS GENUS	193
— pectinated	246	— Ascanian	193
— Schlosserian	246	— Russelian	195
— Eel	247	— Blochian	197
— Carp	247	— Cepedian	198
— Nilotic	248	HOLOCENTRUS GENUS	553
— black-tailed	248	— Sogo	553
GOBOMORUS GENUS	249		

CONTENTS.

vii

Holocentrus, Schraetzer	554	Holocentrus, pied	568
Radula	554		
Gaterina	555	Imminset, long-tailed	229
greenish	555		
five-lined	555	Knigh Fish, American	387
Bengal	556		
Tigrine	556	LABRUS GENUS	481
decussated	557	Hepat	481
striated	557	eared	482
argentine	558	Oyena	482
black	558	falcated	482
acerine	559	operculated	483
blueish	559	Ceylonese	483
variegated	559	rufous	484
Cottoid	560	gray	484
Philadelphic		Jaculator	485
.	560	lunar	485
great	560	long-snouted	486
Forskal's	561	hooped	486
Tauvin	561	hirsute	487
Ongo	561	half-red	487
golden	562	trilobate	487
four-lined	562	Scare	488
fasciated	562	chain	488
punctated	563	black-bellied	489
spur-gilled	563	bifasciated	489
Surinam	564	four-spined	490
African	564	demi-disk	490
Japanese	565	forked	490
Merra	565	six-banded	491
tortoise-shell		bellied	491
.	566	filamentous	491
marginied	566	angular	492
Sonnerat's	567	eight-striped	492
lanceolate	567	punctulated	492
blue-spotted		Commersonian	493
.	568	smooth	493

Labrus, large-finned . . .	494	Labrus, parterre . . .	510
—— fifteen-spined . . .	494	—— Parrakeet . . .	510
—— great-headed . . .	494	—— blood-spotted . . .	511
—— Plumier's . . .	495	—— full-eyed . . .	511
—— Gouan's . . .	495	—— Ascanian . . .	512
—— nine-spined . . .	495	—— beautiful . . .	512
—— red-lined . . .	496	—— bull-eyed . . .	513
—— broad-striped . . .	496	—— blue-finned . . .	513
—— undulated . . .	497	—— chequered . . .	514
—— annulated . . .	497	—— small-scaled . . .	514
—— Brazilian . . .	497	—— girdled . . .	515
—— Peacock . . .	498	—— Diana . . .	515
—— Ballan . . .	498	—— great-toothed . . .	515
—— ancient . . .	499	—— Neustrian . . .	516
—— varying . . .	500	—— perdic . . .	516
—— green . . .	501	—— soft-backed . . .	517
—— striped . . .	501	—— Guaza . . .	517
—— bimaculated . . .	502	—— obscure . . .	518
—— trimaculated . . .	502	—— Chinese . . .	518
—— Comber . . .	503	—— Japanese . . .	518
—— Cook . . .	503	—— linear . . .	519
—— Cornubian . . .	504	—— lunulated . . .	519
—— Leopard . . .	504	—— reticulated . . .	520
—— marginal . . .	505	—— guttated . . .	520
—— ocellate . . .	505	—— guttulated . . .	521
—— black-eyed . . .	505	—— Bergylt . . .	521
—— Nilotic . . .	506	—— carinated . . .	522
—— browed . . .	506	—— white-banded . . .	522
—— cinereous . . .	506	—— punctated . . .	522
—— mixed . . .	507	—— pleat-lipped . . .	523
—— dark . . .	507	—— Onite . . .	523
—— fulvous . . .	507	—— anile . . .	524
—— fuliginous . . .	508	—— Norwegian . . .	524
—— fuscous . . .	508	—— Jurella . . .	525
—— marbled . . .	509	—— Parrot . . .	526
—— large-tailed . . .	509	—— Karut . . .	526
—— crenulated . . .	509	—— cupreous . . .	527

CONTENTS.

ix

LONGHURUS GENUS	531	Mackrel, Carang	599
bearded	531	red	600
		chlorotic	600
MACROURUS GENUS	229	Kleinian	601
		Lippus	601
Mackrel, common	577	yellow-tailed	602
Coly	580	Ferdau	602
Japanese	580	beautiful	603
gilded	581		
Tunny	581	MULLUS GENUS	611
Thazard	583		
Germon	584	OPHICEPHALUS GENUS	529
Sarda	584		
salient	585	PERCA GENUS	545
Pilot	586		
Zebra	587	Perch, common	545
toothless	587	Sandre	547
Bonito	588	Volga	547
Commerson's	589	Ruffe	546
Alatunga	590	Whiting	548
Madagascar	590	Arabian	549
Botla	591	Lophar	549
Leopard	591	Nilotic	550
mottled	592	ocellated	550
black	593	small	550
Ascension	593	Zingel	551
small	594	Zingelin	551
aculeated	594		
spur	595	PLEURONECTES GENUS	294
rhomboid	595		
saddled	596	Remora, Mediterranean	201
brownish	596	Indian	209
black-necked	597	lineated	211
Scad	597		
Rottlerian	598	SCARUS GENUS	394
Plumier's	599	Cretan	394
purse-eyed	599	green	395

Scarus, rivulatus . . .	396	Sciæna, Plumier's . . .	537
— stellatus . . .	396	— Mauritian . . .	537
— red . . .	397	— Coro . . .	538
— Parrot . . .	397	— ciliated . . .	538
— purpled . . .	398	— Cappa . . .	538
— black . . .	398	— scale-backed . . .	539
— Schlosserian . . .	399	— single-spotted . . .	539
— blue-striped . . .	399	— gibbous . . .	539
— ferruginous . . .	400	— silver-spotted . . .	540
— sordid . . .	400	— red . . .	540
— rostrated . . .	401	— long-spined . . .	541
		— Jarbua . . .	541
SCORPÆNA GENUS . . .	267	— gold-striped . . .	542
— porcine . . .	267	— black . . .	542
— rufus . . .	268	— large-scaled . . .	543
— horrid . . .	269	— spotted . . .	543
— Plumier's . . .	270	— cylindric . . .	544
— Commersonian . . .	271		
— Gape . . .	272	SPARUS GENUS . . .	402
— bicapillated . . .	273	— Gilthead . . .	403
— brachiated . . .	274	— Rajan . . .	404
— aculeated . . .	274	— Lintian . . .	404
— American . . .	275	— red-tailed . . .	405
— spiny . . .	276	— Bream . . .	405
— bearded . . .	276	— Mendole . . .	406
— flying . . .	277	— annular . . .	406
— antennated . . .	278	— great-eyed . . .	407
— didactyle . . .	279	— rose . . .	407
		— Pugre . . .	408
SCIÆNA GENUS . . .	533	— rough . . .	408
— bearded . . .	533	— falcated . . .	409
— Basse . . .	534	— Salp . . .	410
— two-spined . . .	535	— violet . . .	410
— pike-headed . . .	535	— anchor-toothed . . .	411
— lineated . . .	536	— dog-toothed . . .	411
— punctatod . . .	536	— fasciated . . .	412
		— Abilgaard's . . .	412

CONTENTS.

xi

Sparus, green-tailed	413	Sparus, Commersonian	428
— red-finned	413	— golden-lined	429
— Boar	414	— black-tail	429
— gold-tailed	414	— leaping	429
— blue-banded	415	— poisonous	430
— Oriental	415	— salin	430
— rufous-banded	416	— Juba	431
— spined	416	— dusky-backed	431
— bull-eyed	417	— Nippon	431
— Sarb	417	— blue	432
— two-banded	418	— scaly-tailed	433
— Smare	418	— moon	433
— black-tail	419	— bilobate	434
— silver	419	— Cardinal	434
— Synagre	419	— Chinese	435
— broad	420	— bufonite	435
— virgated	420	— Parrot	436
— Hassara	420	— Virginian	436
— Berda	421	— Anthia	437
— Chili	421	— Ascension	437
— golden-eyed	422	— stigma	438
— spine	422	— pentagram	438
— Mormyre	422	— silvery	438
— brownish	423	— Serran	439
— Osbeck's	423	— Squirrel	439
— variegated	423	— Hamrur	440
— Marseilles	424	— diagram	440
— Castaneola	424	— silver-eyed	441
— Brunnich's	424	— striated	441
— Mahsena	425	— Vosmaer's	441
— Harak	426	— Niptic	442
— Ramak	426	— Japanese	442
— great-eyed	426	— hexagonal	443
— silver-eyed	426	— lunulated	443
— gold-green	427	— gold-banded	443
— Mylostome	427	— naked-headed	444
— Mylio	428	— triangle	444

Sparus, small-mouthed	445	Sparus, Adriatic	462
—— insidious	445	—— magnificent	463
—— Orph	446	—— Polymna	463
—— Maroon	446	—— palpebral	464
—— rhomboid	447	—— black	464
—— reticulated	447	—— golden-finned	465
—— Galilæan	448	—— Mediterranean	465
—— rock	448	—— vittated	465
—— Peacock	449	—— Scribe	466
—— radiated	449	—— Chinese	466
—— livid	450	—— spear	466
—— variegated	450	—— saddled	467
—— black-finned	451	—— bident	467
—— green-finned	451	—— marked	468
—— zonated	452	—— Linkian	468
—— punctulated	452	—— greenish-finned	469
—— sanguine	452	—— long-headed	469
—— Atlantic	453	—— bifasciated	470
—— black-banded	454	—— Argus	470
—— Acara	455	—— Tranquebar	471
—— Nochunda	455	—— tortoise-shell	471
—— hemispheric	456	—— semi-fasciated	472
—— Panther	456	—— trilineated	472
—— brachiated	456	—— patched	473
—— Meaco	457	—— Oriental	473
—— Desfontaine's	457	—— Cepedian	473
—— Norwegian	458	—— perculine	474
—— Surinam	458	—— sealed	474
—— ten-spined	459	—— climbing	475
—— Scina	459	—— chaetodontoid	476
—— Lapina	459	—— two-spined	476
—— ramentous	460	—— Cayenne	477
—— ocellated	460	—— trident	477
—— gibbous	461	—— trilobate	478
—— olivaceous	461		
—— Brunnich's	462	Surmullet, red	611
—— Marseilles	462	—— striped	613

CONTENTS.

xiii

Surmullet, oriflamme . . .	613	Stickleback, Canada . . .	609
—— Indian . . .	614	—— skipping . . .	609
—— Bandi . . .	615	—— Japanese . . .	610
—— Japanese . . .	615		
—— banded . . .	616	TRACHYCHTHYS GENUS	630
—— trifasciated . . .	616	—— Southern . . .	
—— bifasciated . . .	617	630
—— spotted . . .	617		
—— radiated . . .	618	TRICHOPUS GENUS . . .	388
—— gold-striped . . .	618	—— Goramy . . .	388
—— beardless . . .	618	—— Arabian . . .	390
		—— Satyr . . .	391
SCOMBER GENUS . . .	579	—— Pallasian . . .	392
		—— monodactyle . . .	392
Stickleback, common . . .	604		
—— smaller . . .	606	TRIGLA GENUS . . .	619
—— fifteen-spined . . .	607		
—— minute . . .	608	VANDELLIUS GENUS . . .	199
—— two-spined . . .	608		
—— Carolina . . .	608	Vandel, Lusitanian . . .	199

Directions for placing the Plates in vol. IV. part II.

The Vignette represents the *Chætodon geometricus*, described at p. 354.

Plate 26 to face page 186	Plate 50 to face page 375
27 ————— 192	51 ————— 377
28 ————— 194	52 ————— 378
29 ————— 196	53 ————— 384
30 ————— 197	54 ————— 386
31 ————— 200	55 ————— 389
32 ————— 213	56 ————— 394
33 ————— 229	57 ————— 396
34 ————— 233	58 ————— 408
35 ————— 255	59 ————— 411
36 ————— 257	60 ————— 415
37 ————— 261	61 ————— 434
38 ————— 266	62 ————— 435
39 ————— 269	63 ————— 436
40 ————— 271	64 ————— 438
41 ————— 280	65 ————— 441
42 ————— 280	65* ————— 453
43 ————— 296	66 ————— 455
43* ————— 303	67 ————— 478
44 ————— 304	68 ————— 484
45 ————— 319	69 ————— 491
46 ————— 325	69* ————— 496
47 ————— 331	70 ————— 496
48 ————— 344	71 ————— 499
49 ————— 357	72 ————— 501

Directions for placing the Plates.

Plate 73 to face page 511	Plate 83 to face page 571
74 ————— 512	84 ————— 576
74* ————— 527	85 ————— 589
75 ————— 528	86 ————— 601
76 ————— 531	87 ————— 605
77 ————— 533	88 ————— 612
78 ————— 538	89 ————— 617
79 ————— 544	90 ————— 618
80 ————— 552	91 ————— 623
81 ————— 557	92 ————— 631
82 ————— 564	

ERRATA.—VOL. IV. PART II.

- P. 410, l. 13, for *Sparus chrysops*, Bloch. read *Lutianus Chrysops*, Bloch.
P. 518, l. 22, place the *asterisk* l. 22, at line 1, after the word *mud*.



HERMATIAN BASTARD-FISH



HERMATIAN BASTARD-FISH

FISHES.

ORDER

THORACICI.

CEPOLA. BAND-FISH.

Generic Character.

Caput breve: dentes curvati, acuti.

Corpus longissimum, compressum, abdomine brevissimo.

Membr. branch. radius sex.

Head short: teeth curved, sharp.

Body very long and compressed: abdomen extremely short.

Gill-Membrane six-rayed.

COMMON BAND-FISH.

Cepola Tænia. *C. argentea*, pinnis rubris, capite obtusissimo, cauda attenuata.

Silvery Band-Fish, with red fins, very obtuse head, and attenuated tail.

Cepola Tænia. *C. pinna caudæ attenuata, capite obtusissimo.*

Lin. Syst. Nat. p. 445.

Tænia auctorum. *Artedi gen. 83. syn. 114.*

Cepola capite truncato. *Block. pl. 170.*

THIS remarkable species is of a form so thin and flat in proportion to its length as to have obtained among the ancient ichthyologists the title of Tænia or ribband-fish. It is a native of the Mediterranean sea, and is sometimes found of the length of three or four feet, though the generality of specimens do not exceed the length of eighteen inches or two feet: the head is short, and rounded, or rather truncated in front: the mouth is wide, opening downwards; the lower jaw somewhat exceeding the upper in length: both are armed with moderately large, sharp, and curved teeth, placed in a single row in the upper, and in a double one in the lower jaw: the tongue is broad and rough: the eyes very large, with silvery irides: the abdomen, properly speaking, is so very short as scarce to exceed the length of the head, and from this part the body is continued of equal diameter for a great extent, gradually tapering as it approaches the tail: the sides are extremely compressed, and the body, both above and below, sharpens into a kind of carina or ridge: the pectoral fins, which are placed immediately beyond the gills, are ovate, and of moderate size: beneath them are situated the ventral fins, which are of nearly similar form: the dorsal, which is of moderate breadth, and furnished with pretty numerous distinct rays, com-

commences from the back of the head, and is continued as far as the tail, which is of moderate size and of an oval shape: the vent fin commences at a small distance from the ventral ones, and like the dorsal is continued to the beginning of the tail: the colour of the body is bright silver, with a dusky tinge above, and marked along the sides by a few obscure, distant, and rather large reddish spots: the lateral line runs straight through the middle of the body from the head to the tail: the fins are all of a pale red colour, and the skin is covered with extremely small scales. This fish is said to swim with great rapidity, in the manner of the Trichiure, and to present a beautiful spectacle by the undulating flexures of its body. It is of a predacious nature, living principally on the smaller kind of crabs and shell-fish, the remains of which are said to be usually found in its stomach. From its extreme thinness, which is such as almost to permit the vertebræ to be seen through the sides, it is scarcely considered as an eatable fish. It is observed to frequent the neighbourhood of the shores, in order the more readily to obtain its food, and is occasionally used by the fishermen for the purpose of a bait.

RUBESCENT BAND-FISH.

Cepola Rubescens. *C. rubescens*, pinna caudæ attenuata, maxillis acutis.

Reddish Band-fish, with attenuated tail, and pointed jaws.

Myrus alter, sive *Serpens rubescens* Rondeletii. Aldr. pisc. p. 367.

Ophidium macrophthalmum. Lin. Syst. Nat. ed. X. p. 259.

THIS species, which seems to be far rarer than the preceding, and which is not very distinctly described by those authors who have seen it, is said to have a pointed rather than a rounded head, and to be not so thin in the body: to be of a pale red colour throughout, marked transversely by numerous, narrow, and somewhat undulated streaks: lateral line as in the preceding species: back and vent fin placed as in the former, but shallower, and with more disjoined or separated rays: indeed from the expression of Rondeletius one might doubt whether he meant that the rays were connected by any membrane or not; ("In dorso a capite ad caudam velut capillamenta tenuia a sese disjuncta; in ventre similia.") It is questioned, and not without reason, by Willughby whether the fish be any thing more than some variety of the former kind; and so far as can be judged from the rude figures of Rondeletius and Aldrovandus, it can scarcely be considered as a distinct species. It is said to be a native of the Mediterranean.

ROUGH-FINNED BAND-FISH.

Cepola Trachyptera. C. capite declivi, maxilla utraque fornicata, pinnis aculeatis serratis scabris. Lin. Syst. Nat. Gmel. p. 1187.

Band-fish with sloping head, arched jaws, and prickly, serrated, rough fins.

NATIVE of the Mediterranean: observed in the Adriatic Gulf: described as having a strait lateral line, with a single row of scales, and the first rays of the fins strong, sharp, and rough.

HERMANNIAN BAND-FISH.

Cepola Hermanniana. C. capite angulato, oculis minimis, ore cirroso, cauda subulata.

Band-Fish with angular head, extremely small eyes, bearded mouth, and subulate tail.

Le Ténioïde Hermannien. *Cepede, 2. p. 533, pl. 14.*

DESCRIBED by Cepede, who ranks it in a distinct genus, under the title of *Tænioides*. He observes that though of the same general form, thinness, &c. yet it differs in many points from the true *Cepolæ*. The eyes are so small as to be with great difficulty discovered, and consist of two small black points: there is no caudal fin, the tail terminating in the same manner as in the *Trichiures*, viz. in a fine point: the dorsal fin, which is very shallow, with soft rays, commences at no great distance from the head, and is continued to the extremity of the

tail: the vent is situated opposite the origin of the back fin, and from it commences a very short and shallow vent fin, continued only to a small distance beyond the vent: the pectoral fins are moderately large, rounded, and furnished with numerous, soft, setaceous rays: the head, which is of moderate size, is as it were marked into facets, as represented on the plate, and at the mouth are placed a few cirri or beards of unequal length: there are no visible scales on the skin, and the disposition of the muscles may be traced through it: the pectoral and anal fins are marked by a few small black specks.

This fish takes its specific name from Professor *Hermann* of Strasburg, so well known by his learned labours in natural history. It seems to have been described from a dried specimen, and in point of natural affinity, may be considered as equally allied to the genera of *Cepola* and *Trichiurus*. No mention is made of its colour or its native region.



ASTACIAN GYNOPTRIS

1888, *Proc. Zool. Acad. London*, published by G. Knapley, 1888, Street.

GYMNETRUS. GYMNETRUS.

Generic Character.

<i>Corpus</i> longissimum, compressum.	<i>Body</i> extremely long, compressed.
<i>Dentes</i> numerosi, subulati.	<i>Teeth</i> numerous, subulate.
<i>Membr. branch.</i> 4—5—radiata.	<i>Gill-membrane</i> four or five-rayed.
<i>Pinna analis</i> nulla.	<i>Anal fin</i> wanting.

ASCANIAN GYMNETRUS.

Gymnetrus Ascanii. *G. argenteus*, corpore longitudinaliter fusco punctato, cirri ventralibus apice dilatatis.

Silvery Gymnetrus, speckled longitudinally with brown points, and with the ventral cirri dilated at the tips.

Regalecus Glesne. *Ascan. ic. rer. nat. t. 11.*

THIS extraordinary fish seems to have been almost unknown till within the space of a few years past; nor are its characters and history yet so distinctly ascertained as might be wished. It is a native of the northern seas, and seems to have been first described by Professor Ascanius in his work entitled *Icones rerum naturalium*, &c. The length of the specimen was ten feet, and the diameter, which was equal throughout the whole length,

about six inches: the head short, the mouth small, and the eyes rather large: on the upper part of the head, before the commencement of the dorsal fin, were situated seven or eight upright, naked rays or processes, of moderate length: the dorsal fin, which was rather shallow, commenced at a small distance beyond these, and running along the whole length of the back, formed by its continuation the tail-fin, which was carried to a very small distance beneath the body, there being, properly speaking, no vent-fin: the pectoral fins were very small, of a slightly ovate or rounded shape, and situated at a small distance from the head: the ventral fins, if they can be said to deserve the name, consisted of a pair of extremely long single rays or processes terminated by a small ovate expanded tip or finny extremity: the gill-covers appeared to consist of five or six radiated laminae: the colour of the whole body was bright silver, with a blueish cast diffused over the upper part of the back: the lateral line was strongly marked, and ran from the gill-covers to the tail, and the sides of the body were marked by several longitudinal double rows of slightly extant, very small, dusky specks: the forehead was white; the fins pale brown.

This fish is said to be generally seen either preceding or accompanying the shoals of herrings in the northern seas, for which reason it is popularly known by the title of *King of the Herrings*.

RUSSELIAN GYMNETRUS.

VAR?

Russelian Gymnetrus.

DESCRIBED from a drawing in the possession of Dr. Russel. General proportions similar to those of the first described species, except that the dorsal fin is considerably wider: from the top of the head proceeds an extremely long process, passing to a considerable distance over the back: this process is very broad at its base, where it appears to be strengthened by four stout rays, which gradually coalesce in their progress till at length they form a single filament: at some little distance beyond the base is a second process of similar form, but much shorter and consisting but of one ray gradually lessening to its point: the pectoral fins are very small: the ventral processes are of great length, but terminate in mere filaments, without any dilated extremities, and the tail, which consists of a lanceolate membrane strengthened by two external or bounding rays, is furnished with a similar terminating filament: the colour of the whole body is bright silver; the fins pale brown; the dorsal fin edged with a dusky border.


The length of the above specimen, which was taken near Vizagapatam in the year 1788, was two feet eight inches from head to tail. It might therefore be considered either as a variety, or a sexual difference of the preceding species in a younger

state; the differences being not such as absolutely to contradict this supposition.

From Dr. Russel's memorandums on this subject, politely communicated during my description of this genus, it appears that no teeth were visible in either jaw, whereas in the Ascanian fish the teeth are expressly said to have been of a subulated form: the opercula consisted of two oblong, flexible, streaked plates: the abdomen was very short, and the vent placed near the head: (in the Ascanian fish it was situated towards the middle of the abdomen.) The caudal fin was not united with the dorsal, and consisted of four rays, connected at the base by a thin membrane, and afterwards joined together and ending in a setaceous thread. The gill-membrane had five rays; the dorsal fin 320: (in the Ascanian fish, notwithstanding its vast length, there were only 120.) The pectoral fin had eleven rays, the ventral two, and the caudal four*.

In the British Museum is a dried specimen of a fish which appears to be nearly allied to the preceding, but which is not sufficiently perfect to admit of very exact description: the body is much shorter in proportion: the lateral line extremely strong or distinct: the colour of the whole animal, except the fins and processes (which are red), a bright silver; and from the top of the

* I must here observe that, (regarding the number of rays in the fins of fishes, and more especially in those which have long and shallow fins, as a character on which no very great dependence is to be placed,) I have not been solicitous to introduce it, except in a few instances, into the present work.



BLOCHIAN GYMNETRUS.

seen July 1 London. Published by G. Edwards Fleet Street.



LEPIDION LEWINI (LYMNPTERIS)

Lepidion lewini published by G. Brachet, Paris, 1894

head proceeds a very strong horn-shaped process of the length of several inches, gradually tapering into a slender extremity: the length of this specimen is about four feet and a half, and the breadth about five inches, exclusive of the back fin.

BLOCHIAN GYMNETRUS.

Gymnetrus argenteo-cærulescens, fasciis linearibus obliquis, maculisque rotundatis fuscis, pinnis rubris, ventralibus quatuor.

Blueish-silvery Gymnetrus, with oblique, linear, brown bands, and rounded spots, red fins, and four ventral processes.

Gymnetrus Hawkenii. G. pinna ventrali biradiata. Bloch. pl. 423.

THIS, which is a native of the Indian seas, and which appears also to be occasionally seen in those of Europe, is described by Dr. Bloch from a drawing communicated by J. Hawkins, esq. In its general appearance it is much allied to the preceding kinds, but appears to be furnished with two pair of ventral processes, which are of considerable length, and terminate in large, dilated, finny extremities of an oval form: the back-fin is continued as far as the tail, which, in the figure given by Dr. Bloch, is represented as of the usual form in the generality of fishes, (a circumstance probably owing to a mistake of the draughtsman*.) The colour of this species is silvery, with a blueish cast

* I am assured by Mr. Hawkins that this is really the case, the specimen from which the drawing was taken having been defective in that part.

on the upper parts, and several transverse, alternate, brownish shades continued along the body; accompanied by large, distant round spots of a similar colour: the fins and processes deep crimson: the pectoral fins pretty large in proportion.

It appears from a print, published in the year 1798, that a specimen of this fish was thrown on the coast of Cornwall in the month of February in the same year. Its length was eight feet six inches, its breadth in the widest part ten inches and a half, and its thickness only two inches and three quarters: the tail in this specimen was wanting: the colours the same as in the specimen figured by Dr. Bloch.

CEPEDIAN GYMNETRUS.

• Le Regalee lanceolé. *Cepede pisc.* 2. p. 219.

This, which must be considered as a doubtful species, is described by Cepede from a coloured Chinese drawing belonging to the Museum of the Prince of Orange. The author refers it to the present genus from the circumstance of its wanting the anal fin, and gives it the trivial name of lanceolatus from the shape of its tail, which is pointed at the tip. This fish is of a gold-colour, shaded with brown, and is destitute of the ventral fins or appendages so remarkable in the other species.

VANDELLIUS. VANDEL.

Generic Character.

<i>Corpus</i> longissimum, ensi- forme.	<i>Body</i> extremely long, sword- shaped.
<i>Membr. branch.</i> radiis quin- que vel sex.	<i>Gill-membrane</i> five or six- rayed.
<i>Dentes</i> subulati, anterioribus majoribus.	<i>Teeth</i> subulate, those in front largest.

LUSITANIAN VANDEL.

Vandellius Lusitanicus. V. argenteus, cauda furcata.

Silvery Vandel, with forked tail.



Trichiurus ensiformis. Vandelli. Brit. Mus.

THIS fish, of which a dried specimen, in good preservation, occurs in the British Museum, is found, though very rarely, in the Mediterranean and Atlantic seas; has been sometimes taken near Lisbon, and has been considered by Dr. Vandelli of Coimbra as greatly allied to the genus *Trichiurus*.


The length of the specimen in the British Museum is four feet eight inches; the breadth (exclusive of the back fin, which is about an inch

wide) three inches and a half; and the thickness very slight in proportion. The head is of the usual form in the generality of fishes, and the lower jaw somewhat exceeds the length of the upper: the mouth is moderately wide, and furnished, both above and below, with a row of middle-sized, sharp teeth, but of which the two front ones in the upper jaw, as well as two lateral ones at no great distance from them, far exceed the rest in size: the tongue is cartilaginous, spatular, and about three quarters of an inch in length: the eyes very large, with silvery irides: the gill-covers consist of a large single piece on each side, and the gill-membrane is furnished with five rays: the dorsal fin is continued from the hind part of the head throughout the whole length of the back, as far as the tail, which is small and very much forked, with sharp tips: the pectoral fins are of moderate size, thin, ovate, or rather subacute, with about twelve rays: the anal fin is much shallower than the dorsal, and commences at the distance of but a few inches from the tail, to the base of which it is continued: the lateral line commences from the upper part of the head, and runs straight to the tail: the colour of the whole fish, except on the fins and tail, which are pale brown, is rich silver, with a dusky blueish tinge towards the back, and without any appearance of scales: the dorsal fin has 105 pretty strong rays; the pectoral fins about 12; the anal 20, and the tail 20, subdivided at their tips.

MEDITERRANEAN REMORA.



INDIAN REMORA.



ECHENEIS. REMORA.

Generic Character.

<i>Caput nudum, clypeo ovato plano transversimulcato.</i>	<i>Head furnished above with a flat, ovate, transversely sulcated shield.</i>
<i>Membr. branch. radiis sex.</i>	<i>Gill-membrane six-rayed.</i>
<i>Corpus nudum.</i>	<i>Body without scales.</i>

MEDITERRANEAN REMORA.

Echeneis Remora. *E. fusca, striis capitis octodecim, cauda sub-furcata.*

Brown Remora, with about eighteen bars on the shield, and subfurcated tail.

Echeneis Remora. *E. cauda bifurca, striis capitis 18. Lin. Syst. Nat. p. 446.*

Remora Imperati & Aldrovandi. *Will. p. 119.*

Echeneis pinna caudali semilunata. *Bloch. pl. 172.*

THE extraordinary faculty which this fish possesses of adhering at pleasure with the utmost tenacity to any moderately flat surface, was not unobserved by the ancients, and is described in terms of considerable luxuriance by Pliny in particular, who, giving way to the popular prejudices of his time, represents the Remora as

possessing the power of stopping a vessel in full sail, so as to render it perfectly immoveable in the midst of the sea.

“ Ventum est ad summa Naturæ, &c.”

I shall give the translation in the words of Philemon Holland.

“ Having so far proceeded in the discourse of Nature’s historie, that I am now arrived at the very heighth of her forces, and come into a world of examples, I cann’t chuse but in the first place consider the power of her operations, and the infinitnesse of her secrets, which offer themselves before our eyes in the Sea: for in no part else of this universal frame is it possible to observe the like majestie of Nature: insomuch as we need not seeke any farther, nay we ought not to make more search into her divinitie, considering there cannot be found any thing equall or like unto this one Element, wherein she hath surmounted and gone beyond her own selfe in a wonderfull number of respects. For first and foremost, Is there any thing more violent than the Sea, and namcly when it is troubled with blustering winds, whirlepuffs, storms and tempests? or wherein hath the wit of man been more employed (seeke out all parts of the whole world) than in seconding the waves and billows of the Sea, by saile and ore? Finally, is there ought more admirable than the inenarrable force of the reciprocall tides of the Sea, ebbing and flowing as it doth, whereby it keepeth a current also, as it were the stream of some great river?

" The current of the sea is great, the tide much, the winds vehement and forcible, and more than that, ores and sailes withall to help forward the rest, are mightie and powerfull: and yet there is one little sillie fish, named Echeneis, that checketh, scorneth, and arresteth them all: let the winds blow as much as they will, rage the storms and tempests what they can, yet this little fish commaundeth their furie, restraineth their puissance, and maugre all their force ~~as~~ great as it is, compelleth ships to stand still, a thing which no cables be they never so big and able as they will, can performe. She bridleth the violence and tameth the greatest rage of this universall world, and that without any paine that she putteth herselfe unto, without any holding and putting backe, or any other meanes save only by cleaving and sticking fast to a vessell: in such a sort as this one small and poore fish is sufficient to resist and withstand so great a power both of sea and navie, yea and to stop the passage of a ship, doe they all what they can possible to the contrarie. What should our fleets and armadoes at sea, make such turrets in their decks, and forecastles? what should they fortifie their ships in warlike manner, to fight from them upon the Sea, as it were from mure and rampier on firme land? See the vanitie of man! alas, how foolish are we to make all this adoe? When one little fish, not above half a foot long, is able to arrest and stay perforce, yea and hold as prisoners our goodly tall and proud ships, so well armed in the beake-head with yron pikes and brazen

tines; so offensive and dangerous to bouge and pierce any enemie ship which they doe encountre. Certes, reported it is, that in the naval battaile before Actium, wherein *Antonius* and *Cleopatra* the queene were defeited by *Augustus*, one of these fishes staied the admirall ship wherein *M. Antonius* was, at what time as he made all the hast and meanes he could devise with help of ores to encourage his people from ship to ship, and could not profaile, untill he was forced to abandon the said admirall and goe into another galley. Meanwhile the armada of *Augustus Caesar*, seeing this disorder, charged with greater violence, and soone invested the fleete of *Antonie*. Of late daies also, and within our remembrance the like happened to the Roiall ship of the Emperour *Caius Caligula*, at what time as he rowed backe, and made saile from Astura to Antium; when and where this little fish detained his ship, and (as it fell out afterward) presaged an unfortunate event thereby: for this was the last time that ever this Emperour made his returne to Rome: and no sooner was he arrived, but his own souldiours in a mutinie fell upon him and stabbed him to death. And yet it was not long ere the cause of this wonderfull staie of his ship was knowne: for so soon as ever the vessel (and a galliace it was, furnished with five bankes of ores to a side) was perceived alone in the fleete to stand still, presentlie a number of tall fellows leapt out of their ships into the sea, to search what the reason might be what it stirred not?

and found one of these fishes sticking fast to the very helme: which being reported unto *Caius Caligula*, he fumed and fared as an Emperour, taking great indignation that so small a thing as it, should hold him back perforce, and check the strength of all his mariners, notwithstanding there were no fewer than foure hundred lustie men in his galley that laboured at the ore all that ever they could to the contrarie. But this prince (as it is for certain knowne) was most astonied at this, namely, that the fish sticking only to the ship, should hold it fast; and the same being brought into the ship and there laid, not worke the like effect. They who at that time and afterward saw the fish, report that it resembled for all the world a snaile of the greatest making: but as touching the form and sundrie kinds thereof, many have written diversly, whose opinions I have set downe in my treatise of living creatures belonging to the waters, and namely in the particular discourse of this fish: neither doe I doubt but all the sorte of fishes are able to doe as much: for this we are to beleeve, that Pourcellans also be of the same vertue, since it was well knowne by a notorious example, that one of them did the like by a ship sent from *Periander* to the cape of *Gnidos*: in regard whereof, the inhabitants of *Gnidos* doe honour and consecrate the said Porcellane within their temples of *Venus*. Some of our Latin writers do call the said fish that thus staieth a ship, by the name of *Remora*."

The real fact is, that the Remora, being a fish of very weak powers of fin, takes the advantage of occasionally attaching itself to any large swimming body, whether animate or inanimate, which it happens to encounter; adhering to ships, as well as whales, sharks, and many other of the larger fishes: it has even been observed by Commerson, that the Remora is so ill calculated for supporting a long and laborious course in the water, that when left to its own exertions, it generally swims on its back, and that in an unsteady and feeble manner. It is therefore necessary that it should avail itself of the occasional assistance of some larger floating body. For this purpose the upper part of the head is wonderfully constituted; presenting a large, flat, oval shield or area, traversed by numerous dissepiments or partitions, each of which is fringed at the edge by a row of very numerous perpendicular teeth or filaments, while the whole area or oval space is strengthened by a longitudinal division or septum. So strong is the power of adhesion which the fish by this apparatus is enabled to exert, that we are assured by Commerson, whose observations on this subject are detailed by Cope, that, on applying his thumb to the shield of a living Remora, it was affected not only with a strong stupor, but even with a kind of paralysis, which continued for a considerable time after withdrawing his hand. When attached, as is frequently the case, to the skin of a shark or other large fish, it quits not its hold when the former

IMP 92665 DE 29.10.10

राष्ट्रीय पुस्तकालय, कोलकाता
National Library, Kolkata

is drawn out of the water, but continues adhering after the death of the animal; nor is it easy for the strongest arm to effect its separation, unless it be pulled in a lateral direction, so as to force it to slide along the surface of the skin. When a great many of these fishes are thus adhering at once to the sides of a ship, they may in some degree retard its motion, by preventing its easy passage through the water, in the same manner as any other extraneous substances are known to do; nor can it be thought improbable that the adhesion of these fishes in considerable numbers to the side of a small canoe in the earlier ages of mankind, may have operated still more powerfully, and not only have impeded its progress, but even have caused it to incline towards one side; and the tale once related, might have gradually grown into the exaggerated powers afterwards ascribed to the animal.

A second reason of the Remora's thus attaching itself to the larger fishes and to vessels is, that it may be in readiness to avail itself of the occasional remains of the prey of the former, or of the offals thrown out from the latter; being naturally voracious, and by no means delicate in its choice, and frequently following vessels in great numbers in order to obtain occasional supplies of food. The Remora also, especially in stormy weather, adheres to rocks, like the Lump-Fish and some others.

The Remora is principally an inhabitant of the Mediterranean and Atlantic seas. Its general co-

lour is an uniform brown without any material difference of shade either on the upper or lower surface. It sometimes however varies in colour, and Commerson assures us that when a great number of these fishes are attached either to the side of a vessel or to a large fish, it is not very uncommon to see one or two which differ from the rest in being of a whitish colour. The skin is smooth and destitute of scales, but marked with numerous impressed points or pores: the mouth is large, and furnished with very numerous small teeth; and the lower jaw is rather longer than the upper: the eyes are small, with yellow irides: the lateral line commences above the pectoral fins, and from thence pretty suddenly descending, runs strait in the tail, which is of a slightly forked, or rather lunated form. The number of transverse divisions on the shield of this species varies from sixteen to twenty, but the most general number is eighteen. The fish grows to the length of about eighteen inches.

INDIAN REMORA.

Echeneis Neurates. E. olivacea, subtus albida, striis capitis viginti quatuor, cauda integra.

Olive-green Remora, whitish beneath, with twenty-four bars on the head, and ovate tail.

E. Neurates. E. cauda integra, striis capitis 24. Lin. Syst. Nat. p. 446.

Iperuquiba et Piraqueba Brasiliensibus. *Marcgr. bras. p. 180.*

Echeneis cauda rotundata. Bloch. t. 71.

THIS species may be easily distinguished from the preceding by its more lengthened or slender shape, as well as by the different form of the tail, which is ovate instead of crescent-shaped. It also differs from the former in its colours; the upper parts of the body being olive-green, and the under parts whitish: the fins are yellowish brown, with dusky edges. In its habits it resembles the preceding species, but is more frequently seen in the Indian and American seas than in those of Europe. The Count de Cepede informs us, from the manuscripts of Commerson, that it is also very common about the coasts of Mozambique, where it is sometimes made use of for the following very singular manner of catching turtles. A ring is fastened round the tail of the fish, in such a manner as to prevent its escape, and a long cord fastened to the ring. When thus prepared, the fish is carried in a vessel of seawater, and when the boatmen observe a turtle sleeping, as is the frequent custom of those ani-

mals, on the surface of the water, they approach as near as possible without disturbing it; and then, throwing the Remora into the sea, and giving it the proper length of cord, it soon attaches itself to the breast of the sleeping turtle, which is thus easily drawn to the boat by the fishermen. If we may depend on the accuracy of this description, it seems to give some degree of confirmation to a mode of fishing recorded by Aldrovandus and others, in which a fish of a very unusual form, and which seems to be not very distinctly described, is said to be employed in a similar manner by the American Indians, and to be naturally furnished with a large pouch at its head, with which it seizes and draws up its prey.

This species is said to arrive at a much larger size than the preceding; being found of the length of two or three feet, and even, according to a description quoted by Dr. Bloch, to that of seven feet. The usual number of divisions on its shield is from twenty-two to twenty-four.

LINEATED REMORA.

Echeneis Lineata. *E. cauda cuneata, striis capitis decem, lineis albis utrinque duabus longitudinalibus.* Menz. Lin. Trans. 1. p. 187. pl. 17. f. 1.

Remora with cuneated tail, ten streaks on the shield, and two longitudinal white lines on each side of the body.

This species is an inhabitant of the Pacific ocean, and is thus described by Mr. Archibald Menzies, in the first volume of the transactions of the Linnæan Society.

“ The body of this fish is about five inches long, subulate, smooth, and of a dark brown colour, dotted all over with minute darker spots, and ornamented with two whitish longitudinal lines on each side, which begin at the eyes, and end in the tail; the under mandible is a little longer than the upper, and both are furnished with minute teeth: the clypeus on the top of the head has but ten transverse streaks, which is the chief distinction of this species.”

In its general proportions this fish approaches very near to the *E. Neurates*, but seems to be a truly distinct species; differing not only in the white lines along the body, but much more remarkably, in the far smaller number of divisions on the shield, which, in a young specimen of the *E. Neurates*, I have observed to be as numerous as in the more advanced animal. The specimen described by Mr. Menzies was found adhering to a turtle.

CORYPHÆNA. CORYPHENE.

Generic Character.

<i>Caput</i> maxime truncato-de-	<i>Head</i> sloping suddenly
clive.	downwards.
<i>Membr. branch.</i> radiis quin-	<i>Gill-membrane</i> five-rayed.
que.	
<i>Pinna dorsalis</i> longitudine	<i>Dorsal fin</i> the length of the
dorsi.	back.

COMMON CORYPHENE.

Coryphæna Hippurus. C. viridi-cerulea, aurantio-maculata,
radiis dorsalibus sexaginta, cauda furcata.


Blue-green Coryphene, with orange-coloured spots, sixty rays
in the dorsal fin, and forked tail.

Coryphæna Hippurus. C. cauda bifida, radiis dorsalibus sexa-
ginta. Lin. Syst. Nat. p. 446.


Coryphæna Hippurus. C. pinna uni radiis viginti quinque.
Bloch. pl. 174.

Hippurus. Gesn. Rondel, Aldr. Will. &c.

THE genus *Coryphæna* is, in general, distinguished by a peculiar splendor of appearance most of the species being of highly vivid and elegant colours. The common Coryphene (popularly known by the name of Dolphin) is an inhabitant of the mediterranean, Indian and Atlantic seas, where it often appears in large shoals, and is sometimes observed to follow ships, devouring with



COMMON CORYTHENE.



PLUNIER'S CORYTHENE.

avidity any occasional articles of food which may happen to be thrown over-board: it will even swallow substances of a very different nature; and we are informed, on the authority of Plumier, that in the stomach of a Coryphene which he examined, were found four iron nails, one of which measured more than five inches. The general length of the common Coryphene is about three feet; but it is often seen of four or even five feet in length: its colour is a bright and beautiful blue-green, accompanied by golden gloss, gradually altering into a silvery cast as it approaches the abdomen: the upper parts are marked by pretty numerous round orange-coloured spots: the head is large, but short and compressed; the lips strong, the mouth wide, and armed with four rows of small curved teeth: the scales are of moderate size, and the whole body is of a compressed form: the fins are green with a tinge of yellow on the rays: the dorsal fin is continued throughout the whole length of the back. When taken out of the water, the beautiful combination of colours fades as the fish expires; the lustre vanishing by degrees, with partial restorations, till at length it becomes of a dull greyish or cinereous cast, without any remains of its former splendor. This gradual evanescence of colour in the dying Coryphene is contemplated by sailors with as much delight as the ancient Romans are said to have exhibited on viewing similar changes in the expiring Mullet, when brought to their tables before the feast began.

The Coryphene is a fish of a very strong and

vigorous nature, swims with great rapidity, is extremely voracious, and is observed to be perpetually engaged in the pursuit of the smaller fishes: it is considered as one of the most cruel persecutors of the flying-fish (*Exocoetus*), which, in order to avoid its pursuit, avails itself of its power of temporary flight, or rather of springing at once to a considerable distance over the surface of the water; in which action however it is often observed to fall a prey to some of the larger gulls, who seize it during its passage. The flesh of the Coryphene is said to be excellent, and it is therefore often taken both by the line and net. It is observed to frequent the shores in spring and autumn in order to deposit its spawn, and sometimes approaches the mouths of the large rivers: the young are said to be of uncommonly quick growth, as is proved from observations made on such as have been kept in reservoirs for any length of time.

BRASILIAN CORYPHENE.

Coryphæna Equiselis. C. viridi-aurea, cauda bifurca, radius dorsalibus quinquaginta tribus.

Golden-green Coryphene, with forked tail, and fifty-three rays in the dorsal fin.

Coryphæna Equiselis. C. cauda bifurca, radius dorsalibus quinquaginta tribus. Lin. Syst. Nat. p. 447.

Guaracapema. Marcgr. bras. p. 160.

This species is so nearly allied to the former that it has been sometimes considered rather as a variety

than truly distinct. It seems to have been first described by Marcgrave, who informs us that it is known to the Brasilians by the name of *Guaracapema*, that it grows to the length of six or seven feet, and is of a silvery green colour on the head and upper parts, variegated with numerous blue spots of different sizes, but all rather small than large, and that the belly is of a whitish colour. He adds that it is a very swift swimmer, and is considered as an excellent fish for the table.

PLUMIER'S CORYPHENE.

Coryphæna Plumieri. C. subfusca, lineis curvis cæruleis variata, subtus argentea, cauda lunata.

Brownish Coryphene, variegated with curved blue lines; beneath silvery, with crescent-shaped tail.

Coryphæna Plumieri. C. pinnae analis radiis quinquaginta quinque. Lin. Syst. Nat. Gmel. p. 1191. Bloch. pl. 175.

This elegant species takes its title from the celebrated Father Plumier, who appears to have been its first describer. It is a native of the American seas, and in its general form is much allied to the common Coryphene, but the head is longer in proportion: the general colour of the fish is bright yellow, with a silvery cast on the abdomen; while the back is brown, variegated with pretty numerous and somewhat irregular blue streaks, forming so many transverse bars on that part of the animal: the fins are yellow, the tail crescent-shaped and

edged with blue: the whole body is covered with moderately large scales, none being visible on the head. The specimen figured in the manuscripts of Plumier measured about eighteen inches in length, but the fish probably grows to a much larger size.

BLUE CORYPHENE.

Coryphæna Cærulea. *C. tota cærulea, squamis magnis, cauda lunata.*

Blue Coryphene, with large scales, and crescent-shaped tail.

Coryphæna cærulea. C. tota cærulea. Lin. Syst. Nat. Gmel.
1191. *Bloch. pl.* 148.

Novacula cærulea. Catesb. car. N. pl. 13.

THIS is a native of the American seas, and is easily distinguished from most other species by its uniform blue colour, as well as by its great depth or breadth of body and the superior size of its scales: the head is very large, but short, and is covered with smaller scales than those of the body. It is found of the length of eighteen inches or two feet, and may probably grow to a much larger size. In its manner of living it resembles the rest of the genus. It appears to have been first described and figured by Plumier, in whose manuscripts occurred the representation given by Dr. Bloch, and repeated in the present work. It is also tolerably well represented by Catesby in his Natural History of Carolina.

RAZOR CORYPHENE.

Coryphæna Novacula. *C. rufo-flavescens*, capite pinnaque anali lineis cæruleis cancellatis.

Reddish yellow Coryphene, with the head and vent-fin cancellated with blue lines.

Novacula piscis Rondeletii. Will. p. 214. t. O. 2. *Pesce Pettina*. Salo.

THIS, according to Salvian, its most accurate describer, is a small species, scarcely exceeding a palm in length: the head is very large and compressed, as is the whole body, rising into an edge both above and below: the mouth is rather small, and furnished with small and very sharp teeth, and with four larger ones in front: the eyes are small, and situated on the upper part of the head: the dorsal fin, which is of moderate breadth, commences from the back of the head, and is continued nearly to the tail: the vent is placed nearer the head than the tail, and from it is continued the vent fin, which terminates, like the dorsal, at a small distance from the base of the tail: the tail itself is broad, and nearly even at the end: the head, jaws, and gill-covers, are marked by numerous blue lines; while the anal fin and tail are beautifully variegated by a kind of cancellated alternation of green and yellowish lines: the dorsal fin is red, and marked with a few scattered blue spots: the rest of the animal is of a reddish yellow colour, and in habit resembles the *C. Cærulea*, being very deep in proportion to its length. It is a native of

the Mediterranean, where it chiefly frequents the shores, feeding on the smaller fishes and sea-insects, and is numbered among the superior kinds of edible fish.

FIVE-SPOTTED CORYPHENE.

Coryphæna Quinque-maculata. *C. subfusca, subtus argentea, maculis utrinque versus caput quinque nigris.*

Brownish Coryphene, silvery beneath, with five black spots on each side towards the head.

Coryphæna pentadactyla. *C. maculis nigris quinque versus caput longitudinalibus.* *Lin. Syst. Nat. p. 447. Bloch. pl. 173.*

THE five-spotted Coryphene is a native of the Indian seas, and is of similar habit with the preceding, having a large abrupt head, and being very deep in proportion to its length, as well as greatly compressed on the sides: the mouth is of moderate width, and each jaw is furnished with a row of sharp teeth, with two larger ones in front: both head and body are covered with middle-sized scales, and a small distance beyond the eyes, on each side, are five black round and slightly ocellated spots in a longitudinal direction. The usual length of this species is about twelve inches. It is of a gregarious nature, and is said to assemble in such vast shoals about the Molucca islands, that it furnishes a branch of commerce among those people almost as important as that of the Cod fishery among the Europeans.

GILT-TAIL CORYPHENE.

Coryphæna Chrysurus. *C. aurea, corpore maculis parvis cæruleis variato.*

Gold-yellow Coryphene, with the body besprinkled with small blue spots.

Le Coryphene Chrysure. *Cepede. 3. p. 186.*

OBSERVED by Commerson in the equatorial parts of the Pacific, and described, from his manuscripts, by Cepede. Greatly allied in general appearance to the common Coryphene, but is of a more compressed and lengthened form, and differs in the number and disposition of the teeth, which are very small, short, and sharp, and stand separate from each other: the mouth also is much less wide: the general or ground colour of the animal is bright gold, the tail exceeding the other parts in brilliancy: on the throat and breast is a gloss of silver, and on the back are a few blueish clouds: the fins are tinged with blue, and the whole body is sprinkled over with lenticular bright blue spots. In its manners this elegant species resembles the common Coryphene, and is of equal excellence as an article of food.

STRIPED CORYPHENE.

Coryphæna Pompius. *C. dorso supra lineam curvam fasciis
flavescentibus picto.* *Lin. Syst. Nat. p. 147.*

Coryphene with the body marked by transverse yellowish
stripes over the curved lateral line.

Coryphæna cauda æquali, linea laterali curva. *Artedi gen. 16.*
syn. 29.

THIS species is thicker and shorter in proportion than the common Coryphene, with a smaller head and wider mouth: the lower mandible is longer than the upper, and the sides of the head are marked by several small cavities or impressed spots. the lateral line is curved; the pectoral fins very sharp-pointed, and the tail but slightly lunated: over each eye is a gold-coloured spot, and the sides of the body are marked by numerous transverse narrow bands of a similar hue. Native of the Mediterranean and Atlantic seas, where it is said occasionally to accompany vessels in the manner of the common Coryphene.

PARROT CORYPHENE.

Coryphæna Psittacus. C. capite variegato, corpore macula versicolori rhombea.

Coryphene with variegated head, and body marked by a variously coloured rhomboid spot.

Coryphæna Psittacus. C. linea laterali intercepta, pinnis longitudinalibus colore lineatis. Lin. Syst. Nat. p. 448.

OBSERVED by Dr. Garden in the American seas : head beautifully variegated : irides flame-colour edged with blue : on the middle of the body towards the back a rhomboid purple spot, accompanied by green, yellow, and blue variegations : vent in the middle of the body : dorsal and anal fin linear ; the former commencing from the head and reaching, like the vent fin, to the tail, which is even at the end : the beautiful colours of this species, like those of the common Coryphene, are observed to vanish as the fish expires.

MACKREL CORYPHENE.

Coryphæna Scomberoides. C. argentea, dorso subcæruleo, pinna dorsali analique versus caudam sinuatis.

Silvery Coryphene, with blueish back, and the dorsal and anal fin sinuated towards the tail.

Le Coryphene scomberoide. Cope 3. p. 192.

DESCRIBED by Cope from the manuscripts of Commerson, who observed it in the South seas. Size between that of a Mackrel and a Herring :

colour bright silver, tinged with brownish blue, on the back, and with a deeper cast of the same on the top of the head, where it is accompanied by a gilded tinge: all the fins are entirely brown except the ventral, which are white on the exterior sides, while the pectoral are slightly gilt: the lower jaw is longer than the upper, and both are beset with such a number of small teeth in a reversed direction, that the fish may be suspended by introducing the tip of the finger into the mouth: the tongue is large, and shaped somewhat like the human nail; and on its middle is a squarish rough bone: the gill-covers consist of two large, smooth, and rounded laminæ: the lateral line is marked by several undulations, which decrease as they approach the tail: the ventral fins are united at their base by a membrane communicating with a longitudinal furrow beneath the belly, and in which the animal can at pleasure conceal them: they consist each of five or six rays: the dorsal fin reaches from the back of the head to the tail, and is scalloped towards that part, so as to bear a resemblance to the small or spurious fins in the Mackrel tribe: the vent fin is of similar structure: the lower part of the tail is bounded by a sharp carina. Vast multitudes of these fishes were observed by Commerson to follow the French ships for many days together: they appeared to prey chiefly on young flying-fishes, which at intervals sprang around the ships like so many butterflies, which they scarcely surpassed in size.

UNDULATED CORYPHENE.

Coryphæna Fasciolata. *C. lacteo-argentea, circulis transversis fuscis a fasciis pinnae dorsalis super latera defluentibus.*

Silvery-white Coryphene, with transverse brown circles descending on the body from the bands of the dorsal fin.

Coryphæna fasciolata. *C. lacteo-argentea, circulis transversis fuscis a fasciis pinnae dorsalis defluentibus, in dorso passim coherentibus, in ventre evanidis.* *Lin. Syst. Nat. Gmel. p. 1193.*

Pall. sp. zool. 8.

DESCRIBED by Dr. Pallas: length about two inches: colour silvery white, with several transverse brown circular marks descending on each side the body from the back fin, and gradually losing themselves in the whiteness of the sides: head conical, flat above: eyes large: irides gold-coloured: mouth wide: tongue flat, round, and smooth: opening of the gill-covers very large: gill-covers consisting of two rounded laminae: lateral line strait: pectoral and ventral fins hyaline; dorsal and anal variegated by dusky bands: tail deeply forked, and marked by a dusky crescent. Native of the seas about Amboina. Perhaps not arrived at its full size. In the dorsal fin are fifty-four rays.

SAILING CORYPHENE.

Coryphæna Velifera. *C. ex cinereo-argentea, pinnæ dorsali & anali maximis, ventralibus minimis.* *Lin. Syst. Nat. Gmel.* p. 1193. *Pall. sp. zool.* 8.

Silvery-grey Coryphene, with the dorsal and anal fin very large, and the ventral fins very small.

A small species, described by Dr. Pallas. Native of the Indian seas: colour silvery grey; body gradually tapering to the tail, and covered by eleven longitudinal rows of rather large, thin, slightly striated scales, each of which is marked at the base by a small recumbent spine, and is emarginated at the tip: dorsal and anal fins very large; ventral very small: in the dorsal fin are fifty-five rays.

 LINEATED CORYPHENE.

Coryphæna Lineata. *C. capite picto lineis transversis coloratis.* *Lin. Syst. Nat. Gmel.* p. 1195.

Coryphene with the head variegated with transverse, coloured stripes.

NATIVE of the American seas, observed by Dr. Garden: size not mentioned: head naked, compressed: two sharp front teeth longer than the rest in both jaws: head, dorsal and anal fin marked by coloured streaks: tail rounded: body covered with large scales.

JAPANESE CORYPHENE.

Coryphæna Japonica. *C. lutea, branchiarum apertura rima transversa.* *Lin. Syst. Nat. Gmel. p. 1194. Houtuyn act. Haarl. 20. p. 315.*

Deep-yellow Coryphene, with the branchial aperture transverse.

LENGTH about six inches: colour deep yellow, scales moderately small and tender: described by Houttuyn in the Haarlem Transactions. Native of the Japanese seas. The dorsal fin has twenty-four rays.



GREENISH CORYPHENE.

Coryphæna Virens. *C. virens, pinnis appendiculis filiformibus.*

Greenish Coryphene with filiform appendages to the fins.

Coryphæna Virens. *C. pinnis appendiculis filiformibus.* *Lin. Syst. Nat. Gmel. p. 1194.*

GENERAL colour greenish: dorsal and anal fin terminated by a lengthened filiform process: native of the Indian seas: the dorsal fin has twenty-six rays.

CHINESE CORYPHENE.

Coryphæna Sinensis. *C. viridi-argentea, pinna anali brevissima.*
 Silvery-green Coryphene, with very short vent fin.
Le Coryphene Chinois. *Cepede 3. p. 209.*

THIS is described by the Count de Cepede merely on the authority of a Chinese drawing, and is said to be of a green colour, more or less deep on different parts of the body, and accompanied here and there by a gloss of silver: the dorsal fin is very long, but the vent fin very short; and the tail is of a rounded shape: the body, gill-covers, and tail, are covered with large scales: the lower jaw is longer than the upper, and turns upward in a slight degree.

The volume of Chinese drawings from whence this species is described made a part of the collection of the Prince of Orange, lately ceded to France, and at present in the national museum.

PLAT-SNOUTED CORYPHENE.

Coryphæna Sima. *C. cauda integra, labio inferiore longiore.*
Tin. Syst. Nat. Gmel. p. 1194.
 Coryphene with the lower lip longer than the upper, and even tail.

SIZE uncertain: native of the Indian seas: in the dorsal fin are thirty-two rays.

SHARP-TAILED CORYPHENE.

Coryphæna Acuta. *C. cauda acuminata, linea laterali convexa.*
Lin. Syst. Nat. Gmel. p. 1194.

Coryphene with sharp-pointed tail, and convex lateral line.

SIZE uncertain : native of the Indian seas. In the dorsal fin are forty-five rays.

HALF-FINNED CORYPHENE.

Coryphæna Hemiptera. *C. maxillis subæqualibus, pinna dorsali brevi.* *Lin. Syst. Nat. Gmel. p. 1194.*

Coryphene with nearly equal jaws, and short dorsal fin.

AN uncertain species ; native of the Indian seas. The dorsal fin has but fourteen rays.

BRANCHIOSTEGOUS CORYPHENE.

Coryphæna Branchiostega. *C. apertura branchiarum transversa.*
Lin. Syst. Nat. Gmel. p. 1194.

Coryphene with the branchial aperture transverse.

ALLIED to the *C. Japonica*. Native of the Indian seas. In the dorsal fin are twenty-four rays.

CLYPEATED CORYPHENE.

Coryphæna Clypeata. *C. lamina ossea inter oculos.* *Lin. Syst. Nat. Gmel. p. 1195.*

Coryphene with a bony plate between the eyes.

Thus species, which seems at present but obscurely known, is said to be distinguished by a long bony process or lamina situated between the eyes, and is a native of the Indian seas: the dorsal fin has thirty-two rays; the pectoral fourteen; the ventral five; the anal twelve, and the caudal seven.

MACROURUS. IMMINSET.

Generic Character.

<i>Caput grande: oculi magni.</i>	<i>Head large: eyes large.</i>
<i>Corpus postice attenuatum</i> <i>in caudam.</i>	<i>Body at the hind part at-</i> <i>tenuated into the tail.</i>

LONG-TAILED IMMINSET.

Macrourus Rupestris. *M. argenteo-cinereus, squamis caruatis, pinnae dorsalis prioris rari, primo retro-dentato.*

Silvery-grey Imminset, with carinated scales, and the first ray of the first dorsal fin toothed at the back.

Macrourus rupestris. *Bloch, pl. 177.*

Coryphæna rupestris. *C. dorso dipterygio, pinnae dorsalis primæ radio primo retro-dentato.* *Linn. Syst. Nat. Gmel. p. 195.*

Mull. prodr. zool. Dan. p. 43.

THE head of this remarkable fish is large and thick; the upper jaw projecting above into the form of an obtuse snout: the eyes are very large: the mouth wide, with five rows of small, curved teeth in the upper jaw, and two rows in the lower: the tongue white, thick, short, and cartilaginous: beneath the tip of the lower jaw hangs a beard or cirrus: the body gradually tapers from the middle part, and at length is continued into a very long, slender, and pointed tail: the whole fish is covered with moderately large, rounded scales, each of

which is furnished in the middle with a strong, toothed carina, ending in a pointed tip: this causes a remarkable roughness of surface; so that the hand is wounded by drawing it over the fish from the tail towards the head: the carinæ of the scales are much stronger on the head than on other parts: the first dorsal fin is situated near the middle of the back, and is furnished with nine or ten rays, of which the first is very strong, and serrated on the back: the second dorsal fin, which is shallow, commences at a small distance from the first, and running to the tip of the tail, is united with the vent fin, which is in a similar manner continued from the tip of the tail to the vent, near the middle of the body: the pectoral fins are rather small and sharp-pointed: and the ventral are of similar form: the lateral line, which is nearer to the back than to the abdomen, commences at the upper part of the gill-covers, and is from thence continued to the tail: the colour of this fish is a silvery grey, deepest on the upper parts: its usual length is about three feet, but it is occasionally seen of larger size: it is a native of the northern seas, and is mostly seen about the coasts of Greenland and Iceland, and is numbered among the edible fishes of the Greenlanders. It is said to swim very swiftly, and when first taken, to struggle with great violence; endeavouring to defend itself by lashing with its tail; its large eyes projecting at the same time to a surprising degree. It is known in some places by the name of Berg-lax, or Mountain Salmon, and

among the Greenlanders is called Ingminniset*. It was first described by Egede in his natural history of Greenland in the year 1741, and has by some naturalists been considered as a species of Coryphene.

* It is from this word, with a slight alteration in the spelling, that the English generic name has been taken.

GOBIUS. GOBY.

Generic Character.

<i>Caput</i> parvum : Oculi approximati.	<i>Head</i> small : Eyes approximated.
<i>Membrana branch. radiis</i> quatuor.	<i>Gill-Membrane</i> four-rayed.
<i>Pinnæ ventrales unitæ</i> in cavam.	<i>Ventral fins</i> united into the form of a funnel.

COMMON GOBY.

Gobius Niger. *G. albedo nigroque varius, pinna dorsali secunda radiis quatuordecim.*

Goby with black and whitish variegations, and about fourteen rays on the second dorsal fin.


Gobius niger. *G. pinna dorsali secunda radiis quatuordecim.*
Lin. Syst. Nat. p. 449.

Gobius ex albo & fusco varius, pinnæ dorsi secundæ radiis sedecim. *Bloch. pl. 38.*

Black Goby. *Penn. Brit. Zool. 3.*

THIS species grows to the length of six inches: the body is soft, slippery, and of a slender form: the head rather large: the cheeks inflated: the teeth small, and disposed in two rows: from the head to the first dorsal fin is a small furrow: the first dorsal fin consists of six rays: the second according to Linnæus, of fourteen, and according to Dr. Bloch, of sixteen: the pectoral of sixteen or seventeen,

LANCE-TAILED GORY.



COMMON GORY.



Small Goby.

closely set together, and the middlemost the longest: the others on each side gradually shorter: the ventral fins coalesce, and form a sort of funnel, by which these fish are said to affix themselves immoveably to the rocks; for which reason they are called by the name of *Rock-Fish*; the tail is rounded at the end: the general colour of the fish is dusky or blackish; but this, on close inspection, is owing to numerous small dusky or blackish specks, accompanied by brown or olive coloured bars and clouds disposed on a whitish ground: the dorsal and anal fins are of a pale blue; the rays marked with minute black spots. This fish is a native of the mediterranean and northern seas, and sometimes enters the mouths of the larger rivers, particularly in the beginning of summer, at which season it deposits its spawn on stones near the shores. It is in the number of edible fish, but is in no particular estimation.

 LANCE-TAILED GOBY.

Gobius Lanceolatus. *G. fusco-flavescens, squamis versus caudam majoribus, cauda elongato-acuminata.*

Yellowish brown Goby, with the scales increasing in size towards the tail, which is large, elongated, and sharp-pointed.

Gobius lanceolatus. *G. cauda longissima acuminata.* Bloch. t. 38. f. 1. *Lin. Syst. Nat. Gmel. p. 1203.*

THIS species is distinguished by the peculiar form of its tail, which is large in proportion to the animal, and sharp pointed at the tip: the body is

of a lengthened shape, and nearly of equal diameter throughout: the head is oblong, and truncated in front; the jaws of equal length, and armed with sharp teeth: the gill-covers consist of two small laminæ, and the opening of the gills is large: the vent is situated much nearer the head than the tail: the body is covered with scales, of which those toward the tail are much larger than those on the upper parts. This is a West-Indian species, it is found in the rivers of Martinique and some other islands.

BOSCIAN GOBY.

Gobius Boscii. *G. cinereus, fusco-punctatus, fuscus septem transversis albidis.*

Grey Goby, speckled with brown, and marked by seven whitish transverse bands.

Le Gobie Bosc. Cepede 2. p. 556. pl. 16. f. 1.

NATIVE of the American seas, and observed by Bosc in the bay of Charles-Town. Head broader than the body: both jaws of equal length: teeth very small: eyes prominent: nostrils elevated: gill-covers terminating acutely: four first rays of the first dorsal fin prolonged into a small filament: body without scales: general colour grey, spotted with brown, and marked by seven irregular transverse whitish or pale bands, fins brown: length fifty-four millimetres; breadth thirteen. Described by Cepede from the communications of Mr. Bosc.

BLUE GOBY.

Gobius Cæruleus. *G. cæruleus, cauda rubra nigro marginata.*

Blue Goby, with red tail bordered with black.

Le Gobie bleu. *Cepede 2. p. 560.*

DESCRIBED by Cepede from Commerson. A highly beautiful, though very small species: colour fine blue, rather paler beneath: tail red, with a black border: length about a decimetre: mouth obtuse; teeth in the lower jaw sharp, and rather longer than those of the upper: eyes rather more distant than in others of the genus: body covered with small rough scales: first dorsal fin triangular, with the rays terminating in lengthened filaments: second dorsal terminated by a ray of twice the length of the rest: vent placed nearly in the middle of the body: tail rounded. Inhabits the seas about the eastern coasts of Africa, where it is used by the negroes as a bait for other fish. From the brilliancy of its colours it appears, when swimming in a calm sea, during a bright sunshine, like a small tube of sapphire, tipped with carbuncle.

BROUSSONET'S GOBY.

Gobius Broussonetii. *G. corpore elongato, compresso, pinna dorsali unica, cauda lanceolata.*

Goby with elongated and compressed body, single dorsal fin, and lanceolate tail.

Le Gobioides Broussonet. *Cepede* 2. p. 380. pl. 17. f. 1.

DESCRIBED by *Cepede* from a specimen in the Stadtholder's Museum. Body very long, much compressed, and covered with a skin so transparent as to admit a view of the disposition of the muscles: jaws furnished with very small teeth: ventral fins rather long, and by their union forming a deep funnel: dorsal fin single: pectoral fins small and rounded: dorsal and anal pretty wide, and reaching as far as the tail, which is of a lanceolate shape: rays of the dorsal and anal fins project beyond the membrane. Native of the African seas.

SPOTTED GOBY.

Gobius Minutus. *G. albidus, ferrugineo maculatus, oculis caeruleis.*

Whitish Goby with ferruginous spots and blue eyes.

Gobius minutus. *G. albicans, ferrugineo maculatus, radiis dorsalibus & caudalibus ferrugineo obsolete striatis.* *Lin. Syst. Nat. Gmel.* p. 1199. *Pall. spic. zool.* 8. p. 4.

THIS species is a native of the European seas, and, according to Mr. Pennant, frequents the sandy shores of our own island during the summer months;

its length is about three inches: the head flat; the eyes large, and very prominent; with blue irides: the tongue large; and both jaws are furnished with teeth; the first dorsal fin consists of six rays: the second of eleven, and is placed at some distance from the other: the vent fin has eleven rays; the tail is even at the end: the body is of a whitish colour, obscurely spotted with ferruginous, and the rays of the dorsal fins and tail are barred with the same colour.

BLUE-FINNED GOBY.

Gobius Jozo. *G. fuscus, subtus albidus, pinnis subcæruleis, radiis dorsalibus eminentibus setaceis.*

Brown Goby, whitish beneath, with blueish fins, and projecting setaceous dorsal rays.

Gobius Jozo. *G. radiis dorsalibus eminentibus setaceis. Lin. Syst. Nat. p. 450.*

Bloch. t. 107. f. 3.

THIS species grows to the length of four or six inches, and is principally distinguished by the blue colour of its fins, and the streaks on the first dorsal fin: the jaws are of equal length, and armed with small, sharp teeth: the lateral line runs in a straight direction along the middle of the body. It is a native of the Mediterranean and the Baltic, and commonly frequents the muddy shores, living on sea-insects, &c. It deposits its spawn on the soft mud; and though very prolific, is not observed to

be very numerous; owing to the small fry becoming the prey of other fishes: as a food it is held in little or no esteem.

BODDERTIAN GOBY.

Gobius Boddaerti. *G. fusco-ceruleus, subtus gilvus, lateribus fusco maculatis.*

Blueish-brown Goby, buff-coloured beneath, with the sides of the body spotted with brown.

Gobius Boddaerti. *G. pinnæ dorsalis anterioris radius cirriformis, tertio longissimo.* *Lin. Syst. Nat. Gmel. p. 1201.*
Pall. spic. zool. 8. p. 11. t. 2. f. 4. 5.

NATIVE of the Indian seas: length about five or six inches: colour blueish brown above, and reddish white beneath: head varied with brown and white: lips thick and fleshy: body covered with small scales, and marked above by seven brown spots in a longitudinal direction; beneath which are seven other spots of the same colour but speckled with white: vent surrounded by a black ring, and terminated by a fleshy protuberance: lateral line marked with minute papillæ and snow-white spots: first dorsal fin blackish blue speckled with white, and furnished with long projecting filaments: second dorsal fin traversed by five or six white lines: tail blueish white.

HARE-LIPPED GOBY.

Gobius Lagocephalus. *G. fuscus, labiis emarginatis, superiore crassissimo.*

Brown Goby, with emarginated lips, the upper one very thick.

Gobius lagocephalus. *G. maxilla superiore hemispherica, lingua lineaque laterali nulla.* Lin. Syst. Nat. Gmel. p. 1202. Pall. spic. zool. 8. p. 14. t. 2. f. 6. 7.

DESCRIBED by Kolreuter in the Petersburg Transactions. Body round, compressed towards the tail: size of a finger: colour grey or dark brown: covered with minute scales: head short, thick, without scales: mouth transverse; upper jaw very thick; lower with a few larger teeth, which are slightly hooked: lips cleft in the middle, doubled, the upper fleshy and very thick, reaching much beyond the jaw; palate with numerous crowded small teeth on the margin: eyes distant, small, covered with the common skin: rays of the first dorsal fin simple, of the other and the tail branched: tail oval-rounded: pectoral lanceolate-oval: ventral with very short crowded rays: vent in the middle of the body.

PAGANEL GOBY.

Gobius Paganellus. G. subflavescens, fuscò punctatus, pinna dorsali secunda basi subpurpurea, margine lutea.

Yellowish Goby, speckled with brown, with the second dorsal fin purplish at the base and yellow at the margin.

Gobius Paganellus. G. pinna caudali dorsalique secundu basi purpurascens, priori linea lutea terminali. Lin. Syst. Nat. p. 449.

LENGTH from three, or four, to six inches: body rather compressed; general colour white, more or less tinged with yellow, and sometimes with a cast of green: body marked with small blackish spots: first dorsal fin edged with bright yellow: the second dorsal and the anal tinged with purple at the base: tail nearly strait. Native of the Mediterranean, frequenting rocky coasts.

 RED-FRONTED GOBY.

Gobius Cruentatus. G. albidus, fasciis transversis fuscis, capite punisque rubro maculatis.

Whitish Goby, with transverse brown bands; the head and fins spotted with red.

Gobius cruentatus. G. ore rubro-pustulato, pinnarum dorsalium radiis ultra membranam prominentibus. Lin. Syst. Nat. Gmel. p. 1197.

NATIVE of the Mediterranean; much allied to the *G. Paganellus*, but the rays of the dorsal fins project beyond the membrane: mouth, gill-covers,

throat, and most of the fins spotted with red: general colour dull white, with a few transverse brown bands: ventral fins blueish: tail rounded, and marked with obscure blackish bands. This species has been sometimes regarded as a variety of the *G. Jozo*.

OTAHEITEE GOBY.

Gobius Ocellaris. *G. olivaceus, subtus glaucus, pinna dorsali priore ocello nigro prope basin notata.*

Olivaceous Goby, glaucous beneath, with the first dorsal fin marked towards the base by a black ocellate spot.

Gobius Ocellaris. *G. maxilla superiore longiore, pinna dorsali prima ocellata sexradiata.* *Lin. Syst. Nat. Gmel. p. 1201.*
Brouss. ichth. dec. 1. t. 2.

NATIVE of the rivulets of Otaheitee. Body compressed lanceolate, covered on the hind part with imbricate ciliate scales, obscurely clouded with olive and black, beneath glaucous. Head a little compressed, subconic, blackish, slightly curved above, obtusely carinate in the middle, beneath strait, flat; eyes looking obliquely upwards, pupil black, iris glaucous; teeth unequal, minute, those of the lower jaw less: lateral line nearly in the middle, strait, obscure; vent placed a little behind: fins generally with divided rays, the connecting membrane diaphanous: pectoral and ventral fins blackish; first dorsal short, with olive and brown undulations, and simple rays, setaceous and recurved at their tips: second dorsal fin long, greenish,

tinged with dull red at the tip ; the rays annulated with greenish and brown ; the first ray simple : anal fin long, greenish-brown, with transparent tip : tail oval, greenish, with simple rays.

ARABIAN GOBY.

Gobius Arabicus. G. olivaceus, maculis confluentibus violaceis punctisque caeruleis variatus, radiis dorsalibus quinque elongatis.

Olivaceous Goby, variegated with confluent violet spots and blue specks, with five of the dorsal rays elongated.

Gobius Arabicus. G. pinnae dorsalis primae radiis quinque posterioribus filo rubro terminatis membrana duplo longioribus.

Lin. Syst. Nat. Gmel. p. 1198. Forsk. arab. p. 23.

A SMALL species, about the size of the little finger, and of a greenish brown colour, variegated with numerous violet-coloured spots and blue specks ; the violet spots becoming confluent on some parts : the skin is very soft, and covered with extremely small, close-set scales : all the fins are spotted, and the tail is of a cuneated shape. Observed by Forskal in the Arabian seas.

BLACK GOBY.

Gobius Ater. *G. ater, squamis parvis carinatis, genis prominentibus.*

Deep-Black Goby, with small carinated scales, and prominent cheeks.

Le Gobie noir. *Cepedc. 2. p. 568.*

DESCRIBED by Cope from Commerson : length two or three inches : colour deep black with a bluish or greenish cast : scales very small, carinated : head swelling out on each side : upper lip longer than the lower : teeth small : tongue short and fixed : eyes much approximated : on each side the branchial aperture is a tubercle, and at the vent a small process. Native of the Indian seas ; often entering the mouths of small rivers, particularly where there is a muddy bottom : considerably esteemed as a food, and used by convalescents.

 BLACK-FINNED GOBY.

Gobius Bicolor. *G. fuscus, pinnis omnibus nigris. Lin. Syst. Nat. Gmel. p. 1197. Brünn. pisc. mass. p. 30.*

Brown Goby, with all the fins black.

NATIVE of the Mediterranean : habit of *G. Paganellus* : length three inches ; rays of the fins project in a small degree beyond the connecting membrane.

CLOUDED GOBY.

Gobius Nebulosus. *G. albidus, maculis confluentibus fuscis, squamis magnis rhombeis.*

Whitish Goby, variegated with confluent brown spots, and covered with large rhomboid scales.

Gobius nebulosus. *G. pinnae dorsalis primæ radio secundo filo membrana duplo longiore nigro terminato.* *Lin. Syst. Nat. Gmel. p. 1198. Forsk. arab. p. 24.*

OBSERVED by Forskal in the Arabian sea. Length three inches: colour whitish, variegated with confluent brown spots: beneath plain: scales large, rhomboid, and rough: pupils of the eyes blueish, irides white: pectoral fins glaucous, variegated at the base by obscure brown spots: ventral fins brownish: dorsal and caudal hyaline, spotted with brown: anal hyaline bordered with brown.

 CHINESE GOBY.

Gobius Eleotris. *G. albidus, squamis magnis lævibus, macula prope opercula magna violacea.*

Whitish Goby, with large smooth scales, and a large violet spot near the gill-covers.

Gobius Eleotris. *G. pinna anali radiis novem.* *Lin. Syst. Nat. p. 449. Gobius Chinensis. Osb. it. p. 260.*

NATIVE of China: colour whitish, and covered with large, round, smooth scales: on the back, near the gill-covers, is a large violet-coloured spot: both the dorsal fins are of equal height, and the tail is rounded.

PLUMIER'S GOBY.

Gobius Plumieri. G. fusco-flavescens, subtus albidus, maxilla superiore prominente.

Yellowish-brown Goby, whitish beneath, with thick, projecting upper lip.

Gobius Plumieri. G. maxilla superiore prominente. Lin. Syst. Nat. Gmel. p. 1203. Bloch. t. 178.

NATIVE of the American seas: described by Plumier: length three or four inches: habit plump or fleshy: colour yellowish brown above, yellow on the sides, and white beneath: body covered with small scales: upper lip obtuse and longer than the lower: rays of the first dorsal fin project beyond the membrane: pectoral fins rather large: tail rounded: considerably esteemed as an article of food.

PECTINATED GOBY.

Gobius Pectinirostris. G. dentibus maxillæ inferioris horizontalibus. Lin. Syst. Nat. p. 451.

Goby with the teeth of the lower jaw horizontal.

Apocryptes Chinensis. Osb. it. p. 130.

NATIVE of China: size rather small: particularly distinguished by the position of its teeth in the lower jaw, which project horizontally in such a manner as to represent a small semicircular comb: in the first dorsal fin are five rays, in the second twenty-six, in the pectoral fins nineteen, in the ventral ten, in the anal twenty-six, and in the tail fifteen.

SCHLOSSERIAN GOBY.

Gobius Schlosseri. *G. nigricans, subtus albidus, capite magno, corpore squamis magnis, coriaceis, rotundatis.*

Blackish Goby, whitish beneath, with large head, and body covered with large, rounded, coriaceous scales.

Gobius Schlosseri. *G. ex fusco nigricans, subtus exallidus, pinnae dorsalis primæ radius spinosis.* Lin. Syst. Nat. Cimel. p. 1201. Pall. sp. zool. 8.

THIS species is a native of the island of Amboina, inhabiting fresh waters, and generally residing on the soft mud, in which it conceals itself on the approach of danger: it is about a span in length, of a somewhat compressed form, and tapering but very slightly towards the tail: the head is longish, and much broader than the body; the eyes protuberant, and placed in front of the head: the body is covered with moderately large, coriaceous, rounded scales, and is of a dusky colour above, and whitish beneath: the pectoral fins are attached, on each side the base, to a kind of fleshy arm or foot-stalk. According to Cope the *Gobius barbarus* of Linnæus is in reality no other than the above species: it is thus distinguished in the *Systœna Naturæ*. *G. pinnis pectoralibus flabello insistentibus, pinna dorsali priore radiis 12, posteriore 13.*

EEL GOBY.

Gobius Anguillaris. *G. elongatus fuscus, pinnis rubris, dorsali unica.*

Elongated brown Goby, with red fins ; the dorsal single.

Gobius anguillaris. *G. pinna dorsali unica, cauda rubra.* *Lin. Syst. Nat. Gmel. p. 1201.*

THIS species differs from the generality of the Gobies by its more elongated form, approaching in some degree to that of an eel : the dorsal fin is single, and reaches nearly to the tail, the vent fin being equally extended in the opposite direction : the pectoral fins are very small and rounded : the skin of the body is smooth and slippery ; the colour pale brown, and all the fins red. Native of the Chinese coasts. The dorsal fin has fifty-two rays, the pectoral fins twelve, the ventral ten, the anal forty-three, and the tail twelve.

 CARP GOBY.

Gobius Cyprinoides. *G. cinereus, corpore cyprinacco, squamis magnis subciliatis.*

Grey Goby with carp-shaped body, and large subciliated scales.

Gobius cyprinoides. *G. squamis magnis subciliatis vestitus, pinna caudæ rotundata : radiis membrana fusco-tessellata nexis.* *Lin. Syst. Nat. Gmel. p. 1202. Pall. sp. zool. 8. t. 1. f. 5.*

NATIVE of Amboina : length about three inches and a half : habit resembling that of the Carp tribe rather than that of the Gobies : colour grey-brown,

whitish beneath : scales large, and slightly ciliated at the edges : head rather broader than the body, and covered by a soft skin, crossed with very fine lines : teeth small : eyes lateral ; between them a blackish, triangular, longitudinal crest, resembling a fin in miniature : behind the vent is a tubular process imbedded in a small channel.

NILOTIC GOBY.

Gobius Aphyæ. *G. albidus, nigro-punctatus, fasciis pinnarum fuscis.*

Whitish Goby speckled with black ; the fins barred with brown.

Gobius Aphyæ. *G. fasciis etiam pinnarum fuscis.* Lin. Syst. Nat. p. 450.

A VERY small species, found in the Nile, and in some parts of the Mediterranean sea : length scarce more than an inch : colour whitish varied by a few small black spots : dorsal and anal fin marked by brown bands : eyes very small : the second dorsal fin has seventeen rays.

BLACK-TAILED GOBY.

Gobius Melanurus. *G. pinna dorsali unica, cauda nigra.* Lin. Syst. Nat. Gmel. p. 1201. Brouss. ichth. dec. 1.

Goby with single dorsal fin, and black tail.

A SMALL species : sufficiently described by its specific character : native country uncertain.

GOBIOMORUS. GOBIOMORE.

Generic Character.

Habitus generis præcedentis. || *Habit* as in the preceding
genus.

Pinnæ ventrales distinctæ. || *Ventral fins* distinct.

THE genus *Gobiomorus* is instituted by the Count de Cèpede, and contains those species which differ from the genuine Gobies in having the ventral fins separate, instead of being united, as in that genus, into the form of a funnel : in the Gmelinian edition of the *Systema Naturæ* they are associated with the genus *Gobius*.

SOUTHERN GOBIOMORE.

Gobiomorus Australis. *G. thalassinus, rubro-maculatus, subtus albus, capite flavo pinnis rubro variatis.*

Blue-green Gobimore with red spots ; whitish beneath ; the head variegated with yellow, and the fins with red.

Gobius strigatus. *G. pinna ventrali bipartita, dorsali prima serrata.* *Lin. Syst. Nat. Gmel. p. 1204. Brouss. ichth. dec. 1.*

Le Gobimore Taïboa. *Cèpede 2. p. 587.*

NATIVE of the Southern ocean, and observed about the coasts of Otaheitee : body compressed, much elongated, and covered with squarish scales :

head compressed, but wider than the body: upper jaw rather longer than the lower: teeth unequal; tongue and palate smooth, but the throat roughened by small sharp teeth: first dorsal fin furnished with very long rays: tail large and rounded: colours of the whole animal very elegant; the back being of a blueish green, and the abdomen whitish: the head yellow, varied with green, and marked by dusky streaks and specks: sides of the body varied with reddish spots: dorsal fin transparent, greenish, with red rays and transverse bars: pectoral fins greenish; ventral white; anal green tinged with red: tail yellowish green, with red rays, and marked on each side by a broad, oblique, curved, red stripe.

SLEEPING GOBIOMORE.

Gobiomorus Dormitor. *G. maxilla superiore longiore, cauda rotunda.*

Gobiomore with the upper jaw longer than the lower, and round tail.

Le Gobiomore Dormeur. *Cepede 2. p. 589.*

NATIVE of the fresh-water lakes and rivers of South America: lower jaw longer than the upper: tail very much rounded: eleven rays in the second dorsal fin: eight in the pectoral and anal fins: colour not mentioned by Cepede, who informs us that this species is figured in the drawings of Plumier: supposed to take its name from the slowness of its movements, or want of activity.

GRONOVIAN GOBIOMORE.

Gobiomorus Gronovii. *G. argenteus, corpore scombriformi, dorso subnigro, lateribus nigro-maculatis, cauda furcata.*

Silvery Gobiomore, with mackrel-shaped body, blackish back, sides spotted with black, and forked tail.

Le Gobiomore Gronovien. *Cepede 2. p. 584.*

Gobius Gronovii. *Lin. Syst. Nat. Gmel. p. 1205.*

DESCRIBED by Gronovius: native of the equatorial parts of the American seas: has some affinity to the genus Scomber: body broad or deep, and of a silvery colour, covered with very small, imbricated scales, and spotted on the sides with black: the back also is of a blackish tinge: the eyes are very large; the mouth small; furnished in both jaws, as well as in the palate, with pretty large teeth: gill-covers roundish; back slightly convex: lateral line following the flexure of the back: vent in the middle of the body.

KOLREUTERIAN GOBIOMORE.

Gobiomorus Kolreuteri. *G. albidus, pinnis ventralibus ovatis, brachio affixis.*

Whitish Gobiomore, with ovate ventral fins affixed to an arm or process.

Gobius Koelreuteri. *G. pinna ventrali bipartita, dorsali prima undecim-radiata. Lin. Syst. Nat. Gmel. p. 1205. Pall. sp. zool. 8. t. 2. f. 1. 3.*

Le Gobiomore Koelreuter. *Cepede 2. p. 590.*

DESCRIBED by Kolreuter in the Petersburg Transactions. Length about a span: colour greyish

white: allied in habit to the *Gobius Schlosseri*, having the ventral fins attached to a kind of fleshy arm or process on each side: eyes very much approximated, and situated on the top of the head: lips double and fleshy: teeth unequal and conic: upper jaw furnished on each side with a tooth much longer than the rest: lateral line appearing as if impressed: vent situated towards the middle of the body: tail somewhat lanceolate: first dorsal fin brown, bordered with black: second dorsal fin yellowish, with a longitudinal brown stripe: pectoral fins ovate or fan-shaped: processes on which they are seated very broad.

PISONIAN GOBIOMORE.

Gobiomorus Pisonis. *G. maxilla superiore longiore, pinna dorsali unica.*

Gobionore with the upper lip longer than the lower, and single dorsal fin.

Gobius Pisonis. *G. pinna ventrali bipartita, maxilla inferiore longiore.* *Lin. Syst. Nat. Gmel. p. 1206.*

Amore Pixunia. *Pis. bras. 3. p. 72.*

Le Gobiomoride Pison. *Cepede 2. p. 593.*

A SMALL species, described by Piso in his *Natural History of Brasil*: lower jaw longer than the upper; and both armed with several rows of pretty strong and sharp teeth: head both compressed and depressed, and covered with scales of nearly similar size with those on the back: tail rounded: dorsal fin single. *Cepede* places this in a separate genus, which he calls *Gobiomoroides*.

GREAT-HEADED GOBIOMORE.

Gobiomorus Cephalus. *G. rufus*, capite magno, oculis cæruleis, pinna dorsali analique caudæ vicinis.

Rufous Gobiomore, with large, subdepressed head, blue eyes, and dorsal and anal fins placed near the tail.

Le Gobiésoce Testar. *Cepede* 2. p. 596.

NATIVE of the South-American rivers: head very large, rounded in front, and somewhat depressed at the hind part: eyes situated very near each other: lips double and extensile: abdomen very prominent: dorsal fin single, short, and situated near the tail: vent fin placed still nearer the tail, which is very long, and rounded at the tip: general colour bright rufous, deepest on the back: eyes bright blue. Described by *Cepede* from a drawing by *Plumier*. *Cepede* places it in a separate genus under the name of *Gobiesox*, the position of the dorsal and anal fin giving it in some degree the habit of a Pike.

COTTUS. BULLHEAD.

Generic Character.

Caput corpore latius, spinosum ; *Oculi* verticales, membrana nictitante instructi.

Membr: branch: radiis sex.

Corpus (plerisque) alepidotum, versus caudam attenuatum.

Pinnæ dorsales (plerisque) duæ.

Head broader than the body, spiny : *Eyes* vertical, and furnished with a nictitating membrane.

Gill-membrane six-rayed.

Body (in most species) without scales, attenuated towards the tail.

Dorsal fins (in most species) two.

RIVER BULLHEAD.

Cottus Gobio. C. lævis olivaceo-flavescens, pigro nariatus, subtus albicans, capite utrinque macronuto.

Smooth yellowish-olive Bullhead, variegated with black ; beneath whitish ; the head furnished with a spine on each side.

Cottus Gobio. C. lævis, capite spinis duabus. Lin. Syst. Nat. p. 452.

Cottus spinis curvatis duabus ad utrumque operculum. Bloch. t. 38.

Bullhead, or Miller's Thumb. *Will. itckth.*

River Bullhead. *Penn. Brit. Zool. 3.*

THIS species is an inhabitant of clear brooks and rivers, and is found in most parts of Europe. It generally remains at the bottom ; lying on the



WARRIOR BATTLESHIP



WARRIOR BATTLESHIP

gravel, or concealing itself beneath the stones, and preying on worms, water-insects, and very young fishes. It is said to deposit its spawn in a small hollow, which it forms in the gravel, and to quit its charge with great reluctance. Linnæus indeed, in the *Systema Naturæ*, appears to adopt the old idea of some naturalists; viz. that it will sooner suffer itself to be killed than desert its ova. Dr. Bloch however supposes the notion to be erroneous. In our own country this fish rarely exceeds the length of about three inches and a half; but in other parts of Europe it appears to arrive at a superior size, since the specimen figured by Baron Meidinger, in his descriptions of the fishes of Austria, measures five and a half; and it is even sometimes seen of the length of seven inches. Its general colour is yellowish olive, much deeper on the head and upper parts of the back; and the whole body is more or less clouded and speckled with small dusky variations: the fins are yellowish, speckled with brown: the pectoral fins large, round, and elegantly indented or scolloped at the edges: they commonly consist of thirteen or fourteen rays; the dorsal of seven; the ventral of only four; the anal of thirteen, and the tail of twelve: the head is large, broad, flat, and rather thin at its circumference, being well adapted, as Mr. Pennant observes, for insinuating itself under stones, &c. On the middle part of the gill-covers is a small but strong spine, turning inwards, near which on each side is a second of much smaller size: the eyes are small; the mouth wide, and furnished with pretty numerous,

minute teeth: the skin of the whole body is smooth and slippery. This fish occasionally swims with great strength and rapidity, when in pursuit of its prey, though its general habit is that of lying in an apparently inert state. Notwithstanding its disagreeable appearance, it is considered as an edible fish, and is even regarded as delicate: the flesh however turns of a red or salmon-colour on boiling, and the fish is admitted only at the tables of those of inferior rank. It usually spawns in the months of March and April.

GRUNTING BULLHEAD.

Cottus Grunniens. *C. fuscus albido-variatus, gula ramenti villosa, corpore nudo.*


Brown Bullhead, with whitish variegations, naked body, and throat shagged with cirri.

Cottus grunniens. *C. gula ramenti villosa, corpore nudo.*

Lin. Syst. Nat. p. 451.

Cottus glaber, maxilla inferiore multum cirrata. Bloch. t. 179.

NATIVE of the Indian and American seas: length six or eight inches: colour brown, varied with white on the sides: fins pale, spotted with brown; skin of the whole animal marked by numerous pores, exsuding a mucous fluid, as in most other species: head large; eyes small, mouth very wide; tongue and palate smooth; lower jaw longer than the upper, and furnished with numerous cirri, which appear also on each side the head: lips strong; teeth sharp, distinct, recurved, and placed in several



RED-POUNCE WATTLEHEAD.



LANTERN WATTLEHEAD.

Apogon niger

rows: gill-covers consist of a single piece, armed with three or four spines: branchial orifice very wide. This fish, like some of the Gurnards, &c. when first taken, exerts a kind of abrupt sound, resembling in some degree the grunting of a pig, and which is produced, as in the Gurnards, &c. by the sudden expulsion of air from the internal cavities, through the gill-covers and mouth. It is considered as an esculent fish, but the liver is said to be hurtful.

LASHER BULLHEAD.

Cottus Scorpius. C. fuscus albido-variatus, capite spinis pluribus, maxilla superiore paulo longiore.

Brown Bullhead, with whitish variegations, several spines on the head, and upper jaw rather longer than the lower.

Cottus Scorpius. C. capite spinis pluribus, maxilla superiore paulo longiore. Lin. Syst. Nat. p. 452.

Father-Lasher. *Will. ichth. Penn. Brit. Zool. 3.*

THE general length of this fish, in the Mediterranean sea, is about eight or ten inches, but it is said to arrive at a much larger size in the northern seas: the head is very large, and armed with extremely strong, sharp spines, the largest of which are disposed toward the hind part, and the smallest on the fore-part: the mouth is large, and the jaws are beset with rows of very small teeth: in the roof of the mouth is also a triangular patch of minute teeth: the body tapers pretty suddenly towards the tail, and is covered with a smooth, lubricous skin: the colour is brown, marbled with

white variegations, which are sometimes accompanied by a reddish tinge: the belly is entirely white: the fins and tail are transparent, slightly clouded with brown, and the rays barred regularly with deep-brown: the pectoral fins are very large, rounded, and denticulated or scolloped on the edges: the dorsal and anal of moderate size; the ventral small; the tail even at the end: the lateral line is strait, and formed by a train of small scaly points, covered by the common skin.

This fish is an inhabitant of the mediterranean and northern seas, and is said to be particularly plentiful about the coasts of Greenland, where it is much esteemed as a food. It is a strong fish, swimming with great rapidity, and preying on the smaller fishes, sea-insects, &c. It is said to be able to live a considerable time out of water; having the power of closing the gill-covers in such a manner as to exclude the effects of atmospheric air. When first caught, if held in the hand, it exerts a strong and peculiar sound by the expulsion of air through its mouth, &c. during this action the mouth is opened to the utmost width, the pectoral fins are strongly expanded, and the whole body is agitated by a vibrating or tremulous motion.

FOUR-HORNED BULLHEAD.

Cottus Quadricornis. *C. olivaceus, fusco-variatus, subtus albidus, capite verrucis quatuor osseis.*

Olivaceous Bullhead, variegated with brown, whitish beneath, with four bony tubercles on the head.

Cottus quadricornis. *C. capite verrucis quatuor osseis.* *Lin. Syst. Nat. p. 451.*

Cottus tuberculis quatuor in capite. *Bloch. t. 108.*

GREATLY allied in general appearance to the preceding species, but smaller, and marked on the head by four rough, bony tubercles, two of which are situated towards the middle of the head, and two at the back part : colour olivaceous, marbled with brown : beneath whitish : pectoral fins very large, and rounded. Native of the Mediterranean and Baltic, as well as of the northern seas ; and in manners similar to the preceding species.

ROUGH BULLHEAD.

Cottus Scaber. *C. subfuscus squamosus, fasciis transversis fuscis, capite striis longitudinalibus serratis, linea laterali aculeata.*

Brownish, scaly Bullhead, with transverse brown bands ; head marked with longitudinal serrated streaks ; and spiny lateral line.

Cottus scaber. *C. capite striis, corporisque squamis serratis linea laterali elevata.* *Lin. Syst. Nat. p. 451.*

Cottus linea laterali aculeata. *Bloch. t. 159.*

NATIVE of the Indian seas : habit rather slender : body compressed, of a silvery colour, tinged with

blue on the back, covered with small scales, and surrounded by six or seven moderately broad and distant rufous bands: length about twelve inches: head oblong, roughened in front by four reversed spines, and covered on the sides with small, rough scales: mouth very wide; lower jaw longer than the upper: lateral line marked by a row of spines: fins bluish, spotted on the rays, and variegated on the membranes with brown: pectoral fins rather small.

INSIDIOUS BULLHEAD.

Cottus Insidiator. C. griseus, fusco-variatus, capite supra lineis acutis, ad latus spinis duabus scabro.


Grey Bullhead, with brown variegations, head marked above by sharp lines and on each side by two spines.

Cottus Insidiator. C. capite supra lineis acutis, ad latus spinis duabus scabro. Lin. Syst. Nat. Gmel. p. 1213. Forsk. arab. p. 25.

ALLIED to the river bullhead; but a much larger species, exceeding the length of a foot. Native of the Arabian seas, where it frequents the coasts, concealing itself under the sand, and springing on such of the smaller fishes as happen to approach its haunts. It is of a depressed form, tapering towards the tail, and is of a greyish colour variegated with dusky clouds and spots: the abdomen is white; the tail whitish, and marked in the middle by a blind yellow spot accompanied by two unequal, linear, oblique black bars: on the head are spines similar to those of the river bullhead.



MADAGASCAR BUTLER-HEAD.



From the collection of the British Museum, London.

MARSEILLES BULLHEAD.

Cottus Massiliensis. C. capite polyacantho, pinnis dorsalibus coadunatis. Lin. Syst. Nat. Gmel. p. 1213.

Bullhead with many spines on the head, and united dorsal fins.

THIS species is much allied to the *Cottus Scorpio*, but differs in the particulars mentioned in the specific character. It is a native of the Mediterranean, and is said to occur about the coasts of Marseilles.

MADAGASCAR BULLHEAD.

Cottus Madagascariensis. C. squamosus, capite utrinque spinis duabus curvis sulcoque longitudinali medio.

Scaly Bullhead with two curved spines on each side the head, and a longitudinal middle furrow.

Le Cotte Madégaſſe. *Cepede* 3. p. 248. pl. 11.

DESCRIBED by *Cepede* from the manuscripts of *Commerson*: allied in some degree to the river Bullhead, but longer in proportion, and of larger size: it is also covered with pretty large scales, and the tail is divided by two indentations into three rounded lobes*: the head is depressed, and

* This circumstance appears rather doubtful; and perhaps the figure of this part, as given by *Commerson*, and engraved in *Cepede's* work, may have been meant rather to express two large dusky marks or spots, rather than indentations; especially as it is expressly said by *Commerson* to be *variegated*.

marked by a deep longitudinal furrow; and on each side the head are two sharp curved spines: the eyes are situated very near each other on the upper part of the head: the pectoral fins are of moderate size and of a rounded form: the colour of this fish is not mentioned. Native of the African seas, and observed by Commerson about the coasts of Madagascar.

BLACK BULLHEAD.

Cottus Niger. *C. niger, capite utrinque unispinoso, maxilla inferiore longiore, corpore squamis duris.*

Black Bullhead, with a single spine on each side the head, lower jaw longer than the upper, and body covered with hard scales.

THIS also is a native of the African seas, and is described by Cope from Commerson. Size and habit similar to those of the black Goby: general colour black or dark brown, the second dorsal fin as well as the vent fin being speckled and bordered with deep black; and the first dorsal fin clouded with yellow, and marked by two longitudinal blackish bands: head thick, and largest at the hind part, where it is armed on each side by a small spine, and appears as if swelled: mouth very wide: lower jaw longer than the upper, and both armed with a row of small, short, close-set teeth, resembling those on two bony protuberances near its throat: the whole body, as well as the tail itself, covered with very rough scales.

SOUTHERN BULLHEAD.

Cottus Australis. *C. albidus, capite aculeato, corpore fasciis transversis fuscis.*

Whitish Bullhead, with aculeated head, and body marked with transverse livid bands.

Southern Cottus. *White's Journ. Bot. Bay, p. 266.*

A DOUBTFUL species; described by myself in Mr. White's Voyage to Botany Bay. Length about three inches and a half: colour yellowish white, tinged with brown on the upper parts, and variegated by three or four transverse blackish bands: head armed with pretty numerous, strong spines both in front and on the gill-covers: eyes large: body covered with minute scales.

MAILED BULLHEAD.

Cottus Cataphractus. *C. fuscus subfus albidus, corpore loricato, rostro verrucis duobus bifidis, capite subfus cirroso.*

Brown Bullhead, whitish beneath, with the body mailed, the snout furnished with two bifid warts, and the head cirrated beneath.

Cottus cataphractus. *Lin. C. corpore octogono. Bloch. t. 39.*
 Armed Bullhead. *Penn. brit. zool. 3.*

THE general length of this species is about five inches; the head is large, bony, and very rugged: the end of the nose is armed with four upright, short spines, and on the throat are a number of short white beards: the teeth are very small: the

body is octagonal, covered with a number of bony crusts, and divided into several compartments, the ends of which project into a sharp point, and form several echinated lines along the back and the sides from head to tail: the first dorsal fin usually consists of six spiny rays; the second is placed immediately behind it, and consists of seven soft rays: the pectoral are broad and rounded, and are composed of fifteen rays. This fish is an inhabitant of the European seas, and is found in plenty on our own coasts, living on worms and water-insects, and spawning in the month of May. Its colour is brown above, and whitish beneath: and on the back are three or four blackish transverse bands.

JAPANESE BULLHEAD.

Cottus Japonicus. *C. fusco-flavescens, corpore loricato, cirris nullis.*

Yellowish brown Bullhead, with mailed body and without cirri.

Cottus Japonicus. *C. corpore octogono, squamis osscis aculeatis loricato, cirris nullis.* *Lin. Syst. Nat. Gmel. p. 1213.*

THIS, which is a native of the seas about the Kurile islands, was first described by Dr. Pallas, who informs us that it grows to the length of about twelve inches, and is of a yellowish white colour, dusky on the back: its general habit resembles that of the mailed Bullhead, being, like that species, covered with a bony armature, disposed into eight prominent lines along the body, but is destitute of the cirri or beards under the throat:

the head is elongated, compressed, and flattened on the upper part, which is also marked by a longitudinal channel: the snout is obtuse, and divided into two lobes: the jaws are armed with a great number of small teeth: the eyes are situated towards the end of the snout, and over each is a small protuberance: the gill-covers are denticulated at the edges: the fins are marked with dusky streaks, and have very rough rays: the pectoral fins are very large and of a rounded shape, as is likewise the tail.

SINGLE-FINNED BULLHEAD.

Cottus Monopterygius. C. corpore loricato, elongato, fasciis transversis fuscis, pinna dorsali unica, capite inermi.


Bullhead with elongated mailed body marked by transverse brown bands, single dorsal fin and unarmed head.


Cottus monopterygius. C. dors pinna unica, capite inermi. Lin. Syst. Nat. Gmel. p. 1213.

Cottus monopterygius. Bloch. t. 178.

NATIVE of the Indian seas, and found about the coasts of Tranquebar. Habit resembling that of the two preceding species, but the body more slender in proportion: it is covered by an octagonal bony mail, which towards the tail becomes hexagonal: the eyes are situated near the top of the head: the upper jaw is longer than the lower, and furnished with two recurved spines: the gill-covers

consist of a single lamina, and the branchial orifice is very large: the back is marked by a longitudinal furrow: the colour of this species is brown above, and pale or whitish towards the sides, which are marked by a few transverse reddish-brown bands, accompanied by several spots of similar colour: the fins are large, and of considerable length: the dorsal and anal rather short; the ventral narrow; and the tail rounded.





キハダニシ
ニホリイサナ

SCORPÆNA. SCORPÆNA.

Generic Character.

<i>Caput</i> magnum, aculeatum, cirrosum, obtusum, squamis nudum, subcompressum: <i>Oculi</i> vicini. <i>Dentes</i> maxillis, palato, faucibusque.	<i>Head</i> large, aculeated, cirrose, obtuse, without scales, subcompressed: <i>Eyes</i> placed near each other. <i>Teeth</i> in the jaws, palate, and throat.
<i>Membr. branch.</i> radiis septem.	<i>Gill-membrane</i> seven-rayed.
<i>Corpus</i> torosum. <i>Pinna dorsalis</i> unica, radiis anterioribus spinosis.	<i>Body</i> fleshy: <i>Dorsal fin</i> single, with the rays of the fore-part spiny.

PORCINE SCORPÆNA.

Scorpæna Porcus. S. fusca nigro variata, cirris ad oculos naresque.
Brown *Scorpæna* variegated with black, with cirri at the eyes and nose,

Scorpæna Porcus. S. cirris ad oculos naresque. Lin. Syst. Nat. p. 452.

Scorpæna squamis parvis, maxilla inferiore imberbi. Bloch. t. 181.

THE genus *Scorpæna* is distinguished by a peculiar uncouthness of appearance; the head, in some species, being abruptly truncated in front, of vast

size, and armed with various protuberances and spines. Among the most common of the European species is the *Scorpæna Porcus*, which is frequently seen in considerable numbers in various parts of the Mediterranean, where it chiefly frequents the shores, lying in ambush among stones, sea-weeds, &c. and preying on the smaller fishes and sea-insects: the head is large, the mouth wide, with many rows of small sharp teeth; the eyes large; the gill-covers armed with strong spines intermixed with cirri; the body covered with small rough scales, of a dusky colour, varied with black on the back, and beneath pale, with a reddish cast: the dorsal fin is furnished with very strong spiny rays, which the fish, when caught, erects, and thus wounds its adversary: its general length is about twelve or fifteen inches.

RUFIOUS SCORPÆNA.

Scorpæna Scrofa. *S. rufa fusco variata, linea laterali cirrosa.*

Rufous *Scorpæna*, with brown variegations and cirrose lateral line.

Scorpæna Scrofa. *S. cirris duobus ad labium inferius.* *Lin. Syst. Nat. p. 453.*

Scorpæna squamis magnis, linea laterali cirrosa. *Bloch. t. 182.*

In general appearance so nearly allied to the preceding, that at first view it might be mistaken for the same species; but differs in its superior size, as well as in its larger scales; and particularly

HORRID SCORPAENA.



ASTENNATED SCORPAENA.



in having several cirri or processes disposed along the lateral line: the colour also is rather rufous than brown as in the former kind. Of this species it is reported that it preys not only on the smaller fishes, but even occasionally seizes on such of the marine birds as happen to swim in its way. It grows to a very considerable size, and is said to have been seen of the length of four feet: it must consequently prove a very formidable enemy to the smaller marine animals. Notwithstanding its forbidding appearance, it is considered as an edible fish.

 HORRID SCORPÆNA.

Scorpæna Horrida. S. fusca alepidota, tuberculis callosis adpersa.
Brown Scorpæna, without scales, and scattered over with callous tubercles.

Scorpæna horrida. S. tuberculis callosis adpersa. Lin. Syst. Nat. p. 458.

Scorpæna horrida. Sc. alepidota. Bloch. t. 183.

OF all the species yet discovered the present exhibits the most uncouth and forbidding appearance; resembling rather some imaginary monster of deformity than any regular production of Nature. The head is very large, perfectly abrupt in front, and marked by numerous tubercles, depressions, and spines: on the top is a semilunar cavity: the mouth opens from the upper part, and is large and of a shape somewhat resembling a horse-shoe;

and when closed the lower jaw is in a perpendicular direction : both jaws are armed with numerous small teeth : and the upper is furnished with three cirri, viz. one on each side, and one in the middle : the tongue is large and smooth, but the throat is beset with a great number of small sharp teeth : the eyes are very small, and seated at the base of two large tubercles : the head, body, pectoral and dorsal fins, are beset with numerous abrupt cirri or beards : all the fins are very thick or fleshy, and furnished on the fore part with strong rays : the dorsal reaches nearly the whole length of the back ; the pectoral are large and ovate ; the ventral small ; and the tail rounded at the end. The general colour of this hideous fish is a ferruginous brown, deepest on the upper parts ; the abdomen being much paler. It is a native of the Indian seas, and measures twelve or fifteen inches in length.

PLUMIER'S SCORPÆNA.

Scorpæna Plumieri. *S. nigricans, capite lineaque laterali cirris plumatis.*

Blackish Scorpæna, with feathered cirri on the head and lateral line.

La Scorpene Plumier. *Cepede 3. p. 282.*

DESCRIBED by Cepède from the manuscripts and drawings of Plumier. Upper part, and sides of the head, together with the gill-covers, armed with



PLUVIOUS SCORPAENA.



BICAPITATED SCORPAENA.

triangular, flat, and sharp spines: between the eyes are situated four fringed cirri, and four others, of similar form but smaller, are placed on the upper lip; while several more are disposed along the lateral line: the body is covered with scales of moderate size: the first part of the dorsal fin, which is of a rounded form, is furnished with spiny rays, and the tail is also rounded: the general colour of the fish is a very deep or blackish brown, and on the pectoral fins are a few small brown spots: size not mentioned: native of the American seas.

COMMERSIONIAN SCORPÆNA.

Scorpæna Commersonii. S. corpore pinnisque fusco flavoque transversim fasciatis, superciliis serratis, maxilla superiore cirrosa.

Scorpæna with the body and fins transversely banded with brown and yellow, serrated eyebrows, and bearded upper lip.

La Scorpene Mahé. *Cepede 3. p. 279.*

DESCRIBED by Cepede from the manuscripts of Commerson: the back is arched and carinated; the lateral line curving downwards: body covered with small scales: head large, and armed with numerous spines: orbits of the eyes elevated, and forming two denticulated crests, with a furrow between them: lower jaw somewhat longer than the upper, which is extensile and furnished on each side with three or four beards: teeth very small and close: nostrils each marked by a small filament: gill-covers each composed of two plates, the first of

which has two cirri at the lower edge, and three spines at its posterior edge: the second plate is triangular, and greatly elongated behind: the dorsal fin is marked in different parts by large and unequal sinuations: the pectoral fins are so long as almost to reach as far as the extremity of the back fin, and the tail is of a rounded shape: the body is varied with brown and yellow transverse bands, and all the fins variegated with the above colours: this species was observed by Commerson about the isles of Mahe, in the year 1768.

CAPE SCORPÆNA.

Scorpæna Capensis. S. capite utrinque supra oculos quadridentato, cauda subequali. Lin. Syst. Nat. Gmel. p. 120. Gronov. zooph. p. 68.

Scorpæna with the head marked on each side over the eyes with four teeth or processes, and nearly even tail.

LENGTH near a foot: body covered with small, rough, imbricated scales: head large: eyes large, and situated on each side: mouth wide: jaws equal: gill-covers each composed of two very large scaly plates, the upper of which terminates in four denticulations; and the lower in a single point: back arched and carinated: pectoral fins large and ovate: native of the Cape of Good Hope: described by Gronovius.

BICAPILLATED SCORPÆNA.

Scorpæna Bicapillata. S. capite truncato, subdepresso, postice filamentis duobus longissimis instructo.

Scorpæna with abrupt subdepressed head, furnished at the back part with two extremely long filaments.

La Scorpene Double-Filament. *Cepede 3. p. 270.*

DESCRIBED by Cepede from Commerson : length about four inches : body rather thick, and covered with extremely minute scales ; head large, somewhat depressed, and marked with protuberances : lower lip much elevated, so as to make the head appear obliquely truncated in front : eyes very small, and much approximated : pectoral fins very large, and so long as to reach as far as the middle of the body : tail rounded : vent-fin rounded, as is likewise the opposite part of the dorsal fin, while the other part, consisting of spiny rays, has the membrane deeply scalloped : above the neck are seated two very slender filaments, of such a length as to reach beyond the tail : native of the Indian seas : colour not mentioned : dorsal fin has eighteen spiny rays, and seven soft ones : pectoral fins seventeen rays : anal seven, and the tail fourteen.

BRACHIATED SCORPÆNA.

Scorpæna Brachiata. *S. capite truncato subdepresso, pinnis ventralibus brachio adnatis.*

Scorpæna with abrupt depressed head, and ventral fins attached to a brachial process.

La Scorpene Brachion. *Cepede 3. p. 272.*

GREATLY allied to the preceding species, which it resembles in the shape of the head, the size and situation of the eyes, conformation of the jaws, smallness of the scales on the body, and rounded tail; but differs in being destitute of the two filaments, so remarkable in the other species, as well as in having the membrane of the dorsal fin much less deeply scalloped; while each of the pectoral fins forms as it were a band, which reaches from beneath the fore-part of the branchial aperture to the vent, and is attached by a fleshy longitudinal prolongation or root, as in some of the Gobies: described by Cepede from Commerson.

ACULEATED SCORPÆNA.

Scorpæna Aculeata. *S. aculeis quatuor supra oculos, laminis operculorum spinosis.*

Scorpæna with four spines above the eyes, and both plates of the gill-covers spiny.

La Scorpene Aiguillonnée. *Cepede 3. p. 268.*

DESCRIBED by Cepede from specimens in the French Museum: distinguished from the rest of

the genus by having four very strong recurved spines above the eyes: both plates of the gill-covers aculeated: dorsal and anal fins more elevated on one part than the other; the elevated part being of a rounded form, as are the pectoral fins and tail: colour not mentioned: native country unknown: seems allied to the *American Scorpæna*.

AMERICAN SCORPÆNA.

Scorpæna Americana. *S. onc cirroso, capite spinoso, pinnis pectoralibus suborbiculatis*.

Scorpæna with bearded mouth, spiny head, and suborbicular pectoral fins.

Scorpæna Americana. *S. pinnis pectoralibus suborbiculatis*. *Lin. Syst. Nat. Gmel. p. 1220.*

HEAD armed with protuberances and spines: two beards on the upper lip; and five or six on the lower: the last fifteen rays of the dorsal fin form a portion more elevated than the rest of the fin, and of a rounded figure: the anal fin, which is situated just beneath this part, is of similar shape: the pectoral fins and tail are also rounded: native of the American seas: size and colours not particularized by Cope, who describes this species from Duhamel: the dorsal fin has thirty-three rays; the pectoral thirteen; the vent fin sixteen, and the tail thirteen.

SPINY SCORPÆNA.

Scorpæna Spinosa. *S. pinna dorsali a capite ad caudam extensa, linea laterali spinosa.*

Scorpæna with the dorsal fin continued from head to tail, and spiny lateral line.

Scorpæna spinosa. *S. linea laterali spinosa. Lin. Syst. Nat. Gmel. p. 1220.*

BODY compressed : head spiny : lateral line marked by spiny points : dorsal fin longer than in most of the genus, reaching from between the eyes to the tail : an uncertain species, described in the index to the Leskian Museum : native country unknown.

BEARDED SCORPÆNA.

Scorpæna Barbata. *S. capite cavernoso, cirris geminis in maxilla inferiore. Gronov. nus. ichth. 1. p. 46.*

Scorpæna with head marked by protuberances and cavities, and two cirri at the lower lip.

THE head of this species is marked in some parts by several protuberances, and in others by pretty large depressions or cavities : on the upper lip are two beards, and the pectoral fins are united to each other by a small membrane : the tail is almost rectilinear : size not mentioned by Cope, who describes this species from Gronovius.

FLYING SCORPÆNA.

Scorpæna Volitans. S. fusco flavoque fasciata, pinnis pectoralibus maximis violaceis albo guttatis.

Scorpæna variegated with brown and yellow transverse bands, with very large violet-coloured pectoral fins spotted with white.

Gasterosteus volitans. Lin. Syst. Nat. p. 491.

Scorpæna volans. S. pinnis pectoralibus trunco longioribus. Bloch. t. 184.

A FISH of a highly singular appearance: general length ten or twelve inches: colour brownish yellow, variegated by very numerous, deep-brown, transverse stripes, three very narrow or linear ones being interposed between the broader ones: pectoral fins extremely large, and of a deep violet-colour, spotted with white: the ventral fins are of similar colours: the other fins hyaline, with the rays barred with brown; those of the dorsal fin projecting beyond the membrane: tail rounded: head sloping downwards, marked by various inequalities of surface, as well as by several spines and cirri, and striped like the body, which is covered with small scales: native of the rivers of Japan, Amboina, &c. and considered as an excellent food: probably uses its pectoral fins for the purposes of occasional flight, like the fishes of the genus *Exocoetus* and some of the *Triglæ*.

ANTENNATED SCORPÆNA.

Scorpæna Antennata. *S. fusco flavoque fasciata, pinnis pectoralibus magnis violaceis, fascia oculari fusca.*

Scorpæna variegated with brown and yellow, with large violet-coloured pectoral fins, and brown band across the eyes.

Scorpæna antennata. *S. fascia oculari.* *Lin. Syst. Nat. Gmel.* p. 1219. *Bloch. t.* 185.

GREATLY allied to the preceding species in habit or general appearance, but somewhat larger, and with a less sloping head: body variegated with numerous brown and yellow transverse stripes: eyes larger than those of the preceding fish: immediately over the eyes are seated two longish, cylindric tentacula or cirri, surrounded by several fibrous brown bars: upper jaw longer than the lower: pectoral fins smaller than in the flying *Scorpæna*, and of a deep violet-colour, with the rays projecting greatly beyond the membrane: dorsal, anal, and caudal fins hyaline, with the rays spotted with brown; those of the dorsal running far beyond the membrane: scales on the body larger than in the flying *Scorpæna*, and the head nearly plain or whitish in front: native of the rivers of Amboina.

DIDACTYLE SCORPÆNA.

Scorpæna Didactyla. *S. fusca, lituris guttisque flavis variata, digitis duobus distinctis ad pinnas pectorales.*

Brown Scorpæna, blotched and spotted with yellow, with two distinct fingers or processes at the pectoral fins.

Scorpæna didactyla. *S. digitis duobus distinctis ad pinnas pectorales.* *Lin. Syst. Nat. Gmel. p. 1219. Pall. sp. zool. 7. t. 4.*


GENERAL length about a foot : form extremely grotesque : general colour dusky brown, varied above by transverse yellow streaks, and beneath by roundish spots of the same colour : skin destitute of scales : head depressed, and furnished on different parts with several abrupt fleshy cirri : eyes large, and situated on two approximated protuberances : snout truncated, and marked on the sides by several angular wrinkles, as are also the lower parts of the gill-covers : lower jaw longer than the upper : tongue prominent, and marked with yellow granules and black streaks : on the lower mandible are strong fleshy beards : lateral line near the back : fins furnished with many far-projecting radii, as in the *volitans* and *antennata* : pectoral fins large, ovate, and barred with black ; tail of similar form and colours ; the dorsal fin reaches from head to tail, and the projecting rays are furnished here and there with several small lacerated appendages : the ventral fins, if they may be so called, consist of two long, distinct, naked processes, placed near each of the pectoral fins : this curious species

was first described by Dr. Pallas, and is a native of the Indian seas: it preys on the smaller fishes, sea-insects, &c. and, notwithstanding its forbidding appearance, is considered as an excellent fish for the table.






BRASILLIAN DORY.



INSTITUTIONS BOOKS



COMMON DORY.

ZEUS. DORY.

Generic Character.

<i>Caput</i> compressum, declive.	<i>Head</i> compressed, sloping down.
<i>Labium superius</i> membrana transversa fornicatum. <i>Lingua</i> (plerisque) subulata.	<i>Upper lip</i> arched by a transverse membrane. <i>Tongue</i> (in most species) subulate.
<i>Corpus</i> compressum, latum, subrhombeum, tenue, splendens.	<i>Body</i> compressed, broad, subrhomboid, thin, and of a bright colour.
<i>Membr. branch.</i> radiis septem perpendicularibus, infuso transverso.	<i>Gill-membrane</i> with seven perpendicular rays, the lowest transverse.
<i>Pinnæ dorsalis</i> radii (plerisque) filamentosi.	<i>Dorsal fin</i> (in most species) furnished with projecting, filiform rays,

BRASILIAN DORY.

Zeus Vomer. Z. argenteus, radio dorsali analique secundo longissimis.

Silvery Dory, with the second ray of the dorsal and anal fin very long.

Zeus Vomer. Z. cauda bifurca, &c. Lin. Syst. Nat. p. 454.

Zeus Vomer. Z. radio dorsali analique secundo longissimo. Bloch. t. 193. f. 2.

SHAPE rhomboidal : length six or eight inches, or more : body very thin, without scales, and of a

bright silver-colour, tinged with a blueish cast on the upper parts: head small: mouth furnished with very small teeth: second ray of the dorsal and anal fin reaching far beyond the membrane: tail forked: native of the American seas, and sometimes seen in those of the north of Europe: considered as an edible fish, but not much esteemed, on account of the extreme thinness of its body.

INDIAN DORY.

Zeus Gallus. Z. argenteus, radio dorsali decimo analique secundo corpore longioribus.

Silvery Dory, with the tenth ray of the dorsal and second of the anal fin longer than the body.

Zeus gallus. Lin. Syst. Nat. p. 454. Bloch. t. 192. f. 1.

SHAPE rhomboidal, like that of the preceding species: length the same: body very thin, silvery, and without scales: back tinged with a greenish hue: the tenth ray of the dorsal fin, and the second of the anal*, stretching far beyond the membrane: head large; mouth wide: native of the American and Indian seas, and numbered among esculent fishes: when first taken is said to make a grunting kind of noise, like some of the *Gurnaras*.

* It may perhaps be doubted whether the species of this genus are accurately distinguishable by this particular character.

CILIATED DORY.

Zeus Ciliaris. *Z. argenteus*, radius quibusdam in pinna dorsali analique longissimis. *Lin. Syst. Nat. Gmel. p. 1223. Bloch. t. 191.*

Silvery Dory, with some of the rays in the dorsal and anal fin excessively long.

BODY rhomboidal, thin, without scales, and of a bright silver-colour, with a blueish or greenish cast on the back: head small, and very sloping: lower jaw rather longer than the upper: teeth small and sharp: several of the last rays both of the dorsal and anal fin extend to a vast distance beyond the membrane, reaching farther than the tail itself: with these long and flexible filaments the Count de Cepede imagines that the smaller kind of fishes may be attracted, mistaking them for worms, while the animal lies concealed among sea-weeds, &c. waiting for its prey: he also conceives that they may serve to sustain the fish in this situation, by coiling round the stems of sea-plants, &c. This species is a native of the Indian seas, but is not esteemed as a food, the flesh being not only small in quantity but of a coarse nature: in size it nearly equals the preceding species.

INSIDIOUS DORY.

Zeus Insidiator. Z. argenteus, lateribus nigro punctatis, ore angusto extensili.

Silvery Dory, with the sides speckled with black, and narrow extensile mouth.

Zeus Insidiator. Z. ore angusto. Lin. Syst. Nat. Gmel. p. 1221. Bloch. t. 192. f. 2.

NATIVE of the rivers and fresh waters of India: shape rhomboidal, like the former: size smaller: colour bright silvery, blueish green above, and speckled with black points: body without scales: mouth of a much more lengthened form than in the preceding kinds, the lower lip being retractile, and the mouth so constituted as to form at pleasure a kind of tubular snout, through which the animal occasionally ejaculates a drop of water against such insects as happen to alight on, or fly about the aquatic plants near the shores of the waters it inhabits, and by this method (like the *Chaetodon rostratus*, and some few other fishes) obtains its prey.

COMMON DORY.

Zeus Faber. *Z. viridi-aureo fuliginosus, corpore utrinque macula media fusca.*

Gold-green, fuliginous Dory, with a dusky central spot on each side the body.

Faber. *Salvian* p. 203. *Will. ichth.* p. 294. *t. S.* 16.

Zeus Faber. *Z. cauda rotundata, lateribus mediis oculo fusco, pinnis analibus duabus.* *Lin. Syst. Nat.* p. 454.

Zeus Faber. *Z. macula fusca in lateribus.* *Bloch. t.* 41.

THE common Dory is a native of the Mediterranean, Northern, and Atlantic seas, and is distinguished by its large and long head, its dusky or smoky green colour, accompanied by a strong gilded tinge, and particularly by a large, oval, dusky spot on each side the body: it is of a less compressed form than others of its genus except the Opah, and is generally found of the length of twelve or fifteen inches, though it sometimes arrives at a far superior size, and to the weight of ten or twelve pounds: the mouth is wide; the lower jaw longer than the upper, the teeth small and sharp, and the eyes large: the whole body is covered with very small scales, and marked by a curved lateral line, which descending pretty suddenly from the gill-covers, passes on to the tail: the back is much arched, and furnished with a row of strong, small prickles, which are also continued along the curve of the abdomen: at the base of the pectoral fins are also two very strong and sharp spines: the first dorsal fin is large, and several of the first rays project to a considerable distance beyond the con-

necting membrane: the tail is of moderate size, and rounded at the end.

Though the reputation of this fish, at least in our own country, is now established as an excellent article of food, yet its introduction to the tables of the higher ranks is of no very distant date; Mr. Quin being generally considered as the first who ventured to recommend it to the attention of the polite world.

The Dory is of an extremely voracious nature, preying on the smaller fishes and their spawn, as well as on various kinds of sea-insects, the smaller shell-fish, &c. When first taken it is said to make the same kind of sound as the Gurnards and Scorpaenas, by violently forcing out the air from its gill-covers. It has been figured in the works of Salvian, Willughby, and others, but perhaps the most accurate representation may be found in that of Dr. Bloch.



QIAH POKY.

OPAH DORY.

~~Zeus~~ Opah. *Z. cauda sublunari, corpore nunc rubente, nunc viridi, nunc purpureo, albo-guttato.* *Vivarium Naturæ.* 4. t. 140.

Dory with somewhat lunated tail; the body generally either red, green, or purple, with oval white spots.

Zeus Luna. *Z. cauda sublunari, corpore rubente albo-guttato.* *Lin. Syst. Nat. Gmel. p.* 1225.

Opah. *Phil. Trans. abr.* 11. p. 879. t. 5. *Brit. Zool.* 3. t. 42. *Nat. Misc.* 4. pl. 140.

THIS superb species is sometimes found in the Mediterranean and Northern seas, into which it seems to wander as an occasional visitant; being probably a more regular inhabitant of the warmer regions. In size it exceeds every other species hitherto discovered, measuring between four and five feet in length: in colour it appears to vary, the ground being sometimes a brilliant silvery green, and sometimes a bright gold-colour; but in either case the body is variegated on the sides with pretty numerous, and moderately large, oval white spots, while the fins and tail are bright scarlet: the skin is apparently destitute of scales, and perfectly smooth.

Specimens of this fish have been occasionally thrown on the British coasts, instances of which are recorded in the British Zoology and in the Philosophical Transactions. The first regular description seems to have been drawn up by Dr. Mortimer, in the year 1750, from a specimen shewn to the Royal Society, and taken on the coast of Leith

in Scotland. Dr. Mortimer's description is inserted in the Philosophical Transactions for the same year, and is as follows.

“ This fish is smooth-skinned, has no scales, or teeth: it has one erect fin on its back, which arises below its neck, and runs within a little of its tail: on each side, about the middle, between its back and belly, a little forward of the vent, arise a pair of fins: from behind the vent runs one fin, within a little of the tail: the tail-fin is large and forked: the eyes are large; the irides are scarlet, encompassed with a circle of gold-colour verged with scarlet: its nostrils are placed above its eyes: the back and upper part of the body, quite to the tail, was of a dark blue or violet-colour; these and the sides of the body, which were of a bright green, were all speckled with oblong white spots: the chaps were of a pale red; the nose, gills, and belly, were of a silver-colour, and all the fins of a bright scarlet. It was three feet seven inches long, and three feet ten inches round, in the thickest part: it weighed eighty-two pounds: its mouth was small; its tongue thick, almost like the human tongue in shape, but rough, and thick set with beards or prickles, which pointed backwards; so that any thing might easily pass down, but could not easily slip back again: therefore these might serve instead of teeth for retaining its prey or food: its gills resembled those of the Salmon: its body grows very taper towards the tail; and from being compressed to ten inches thickness, becomes near the tail almost round, and about three inches thick:

the whole shape of this fish much resembles that of the sea-bream, but it differs in size, being much larger, and in not having teeth nor scales: the fin standing erect on the back has some aculei next the neck, and rises up to eight inches; but in the middle diminishes to one inch; and near the tail rises again to about three inches: the belly-fin, opposite to this, spreads three inches near the tail, and diminishes towards the vent: the tail-fin is forked, and spreads twelve inches: the gill-fins are nine inches long, and three wide at their basis: the two belly-fins were eleven inches long, and three wide at their basis: it seems to be a new species of fish, not yet described by any author."

Dr. Mortimer adds, that the Prince of *Anamaboe*, on the coast of Africa, being then in England, immediately knew it, and said it was common on that coast, and very good to eat: the natives call it *Opah*, and the English the King-Fish.

The cavity of the abdomen, in the above specimen, is said to have been extremely small, and the bones very strong, bearing some resemblance to those of quadrupeds, particularly the shoulder-blades (as they are termed), which were thought to resemble those of a sheep.

A second specimen was taken in the year 1769, near Newcastle, and is described in the British Zoology by Mr. Robert Harrison.

"*Sept.* 12th, 1769.

"On Saturday last was thrown upon the sands at *Blyth*, a very rare and beautiful fish, weighing between seventy and eighty pounds; shaped like

the sea-bream : the length was three feet and a half ; the breadth from back to belly almost two feet ; but the thickness from side to side not above six inches : the mouth small for the size of the fish, forming a square opening, and without any teeth in the jaws : the tongue was thick, resembling that of a man, but rough, and thick set with beards or prickles, so that any thing might easily pass down, but could not easily return back ; therefore these might serve instead of teeth to retain its prey : the eyes remarkably large, covered with a membrane, and shining with a glare of gold : the cover of the gills like the salmon : the body diminishes very small to the tail, which is forked, and expands twelve inches : the gill-fins are broad, about eight inches long, and play horizontally : a little behind their insertion the back fin takes its original, where it is about seven inches high, but slopes very suddenly, running down very near the tail, and at its termination becomes a little broader : the belly fins are very strong, and placed near the middle of the body : a narrow fin also runs from the anus to the tail : all the fins, and also the tail, are of a fine scarlet ; but the colours, and beauty of the rest of the body, which is smooth and covered with almost imperceptible scales, beggars all description : the upper part being of a bright green, variegated with whitish spots, and enriched with a shimmering golden hue, like the splendour of a peacock's feather : this by degrees vanishes in a bright silvery, and near the belly the gold again predominates in a lighter ground than on the back."

A third specimen was caught at *Brixham* in *Torbay* in 1772. It weighed an hundred and forty pounds, and measured in length four feet and a half, and in breadth two feet and a quarter: its greatest thickness was only four inches, and the general colour was a vivid transparent scarlet varnish over burnished gold, bespangled with oval silver spots of various sizes: the breast appeared like a hard bone, resembling the keel of a ship: the flesh is said to have resembled beef, both in colour and taste.

In the British Museum is a dried specimen of this fish, measuring in length about three feet five inches, from the tip of the snout to the tips of the tail, and in breadth about one foot seven inches in the broadest part.

The elegant figure of the Opah, given in the *Philosophical Transactions*, is repeated in the present work.

 SQUARE DORY.

Zeus Quadratus. *Z. caudæ æquali, corpore cinereo transversim nigroque fasciato.* *Lin. Syst. Nat. Gmel. p. 1225.*

Grey Dory with transverse dusky bars, and even tail.

Faber Glinus fere quadratus. *Sloane Jam. 2. p. 290. t. 251.*

“THIS fish (says Sir Hans Sloan in his *Natural History of Jamaica*) was almost square, with the fins: it was five inches long, and four broad where broadest in the middle, and decreased from thence

to the head and tail gradually: the mouth was little, and set with rows of small and sharp teeth: the tongue round and cartilaginous, the pupil large and black, in a white circle: it had seven fins, two *post branchias*, and two under them in the belly; one on the highest part of the back, with a black inch and half long *ligula*, and several prickles after it: it had one other after this on the back very large, and another on the belly, having each one ear: the tail was almost square; the whole body was coloured with grey or ash-coloured scales, bating three or four transverse and black lines: it had a very crooked line from head to tail."

RED DORY.

Zeus Aper. *Z. rubens, squamis asperis, cauda aquali.*

Reddish Dory, with rough scales and even tail.

Zeus Aper. *Z. cauda aquali, corpore rubente.* Lin. *Syst. Nat.*
p. 1225.

Aper. *Rondel. Aldrov. Will. &c.*

THIS is a small species, in habit much resembling the common Dory, but scarcely more than three inches in length, and of a reddish colour, like that of the sea-bream (*Erythrinus*): it is covered with rough, ciliated scales: the snout is somewhat produced, sharpish, and turns a little upwards, and the mouth is protruded in the act of opening, like that of the common Dory: there are no perceptible teeth, and the eyes are large, with white irides: the dorsal fins are two in number, the anterior

furnished with nine strong and sharp spines, the first of which is very low and scarcely observable, the second four times longer, and the third very long and thick: the second dorsal fin consists of twenty-three soft rays: the tail is even at the end, and bright red at the tip: the vent fin has twenty-six rays, of which the three first are strong, spiny, and thrice as short as the rest: the pectoral fins have about fourteen rays, and the ventral six, the first strong and spiny, the rest soft and branched: this fish generally resides at the bottom, and is only taken accidentally after great storms: from its small size it is not numbered among edible fishes, and is besides said to be coarse and of an unpleasant odor. It is a native of the Mediterranean, and seems to have been first described by Rondeletius.

PLEURONECTES. FLOUNDER.

Generic Character.

Oculi ambo in eodem latere
capitis.

Corpus compressum, latere
altero dorsum, altero ab-
domen referente.

Eyes both on the same side
of the head.

Body compressed, one side
representing the back,
and the other the abdo-
men.

THE singular structure of this genus is justly considered as one of the most curious deviations from the general uniformity or regularity observed by Nature in the external figure of animals, in which (except in a very few instances) both sides of the body are perfectly similar: but in the genus *Pleuronectes* the animal is so constituted that one side appears to represent the back, and the opposite side the abdomen. They swim laterally, and the eyes are always placed on one side. It is from this circumstance that the division of the species is conducted: viz. into those which have the eyes *dextrous*, or towards the right, when the fish is laid with its coloured side upwards with its abdomen towards the spectator; and *sinistrous* when the eyes are towards the left in the above situation of the fish. It is said however that instances have sometimes occurred in which this natural situation has been

reversed; but such instances must be considered as extremely rare*.

With eyes towards the right.

HOLIBUT.

Pleuronectes Hippoglossus. *P. corpore elongato glabro, supra fusco, subtus albido.*

Flounder with lengthened smooth body, brown above, and white beneath.

P. Hippoglossus. P. oculis dextris, corpore toto glabro. Linn. Syst. Nat. p. 456.

P. pinna caudali lunulata. Bloch. t. 47.

Holibut. *Will. p. 99. t. F. 6. Penn. brit. zool. 3.*

THIS species not only exceeds in size all the rest of the present genus, but may even be considered as one of the largest of fishes; having been sometimes found of the weight of three, and even, according to some accounts, four hundred pounds: it is a native of the Mediterranean and Northern seas, and appears to arrive at its greatest size in the latter; it is considered as the most voracious of its tribe; preying on a variety of other fishes as well as on different kinds of crabs, shell-fish, &c. The Holibut is of a longer or more slender form than most other flat-fish: its colour is deep brown above, and white beneath; the body being quite smooth, and covered with moderately small scales: as a food it is considered as very coarse in com-

* A deviation of this kind sometimes takes place in the uni-valve shells.


parison with many others of this genus : yet those which are young, or of moderate size, are in some degree of esteem, the part of the body nearest the fins being considered as far superior to the rest of the animal. In the London markets this fish is usually cut into large pieces when exposed to sale : the Greenlanders are said to cut it into thin slips, which they dry in the sun, and thus preserve for winter use.

SMALLER HOLIBUT.

Pleuronectes Cynoglossus. P. oculis dextris, corpore oblongo glabro, dentibus obtusis, cauda subrotunda. Lin. Syst. Nat. 156.

Flounder with smooth oblong body, obtuse teeth, and roundish tail.

EXTREMELY allied to the preceding, of which it may, perhaps, be rather considered as a variety than a truly distinct species : length somewhat more than two feet : lateral line oblique, but not bowed or curved : native of the northern seas, and found in considerable numbers in the deep bays about the coasts of Greenland : said to be superior to the common Holibut as an article of food.



PLAISE

PLAISE.

Pleuronectes Platessa. P. corpore subfusco glabro maculis rotundis fulvis, capite tuberculis sex.

Smooth brownish Flounder, with round fulvous spots, and six tubercles on the head.

Pleuronectes Platessa. P. oculis dextris, corpore glabro, tuberculis sex capitis. Lin. Syst. Nat. p. 456.

The Plaise. *Penn. brit. zool. 3. Will. ichth. p. 95. Bloch. t. 42.*

THIS species is, in general, easily distinguished at first sight from others of the genus by its shape and colours, being very broad and flat, and of a fine palish brown above, marked both on the body and fins by pretty numerous, but rather distant, round, and moderately large, orange-coloured spots: the under side is white: behind the left eye is a row of six tubercles, reaching as far as the commencement of the lateral line: the mouth is rather small, the lower jaw longer than the upper, and both furnished with a row of small, and rather blunt teeth.

The Plaise is an inhabitant of the Mediterranean, Baltic, and Northern seas, and is found in considerable plenty about our own coasts. Mr. Pennant observes that it is sometimes taken of the weight of fifteen pounds, but its more general weight is far short of this; one of eight or nine pounds being reckoned a large fish: the best are said to be taken off Rye on the coast of Sussex, and about the Dutch coasts: they spawn in the beginning of May: their general food consists of small fishes,

sea-insects, and the smaller kinds of shell-fish. The Plaise is in considerable esteem as a food, though far inferior to the Sole and Turbot: those are most esteemed which are of moderate size, the smaller ones being less firm than those of more advance 1 growth. It may be added that the figure of this fish in Willughby's Ichthyology conveys no very striking idea of the animal, the spots being very indistinctly expressed: the representation in Dr. Bloch's work is admirable.

DAB.

Pleuronectes Limanda. *P. subfuscus, squamis asperis, linea laterali supra pinnae pectorales arcuata.*

Brownish Flounder, with rough scales, and lateral line curving over the pectoral fins.

Pleuronectes Limanda. *P. oculis dextris, squamis ciliatis, spinulis ad radicem pinnarum dorsi anique, dentibus obtusis.* Lin. Syst. Nat. p. 457.

The Dab. Penn. brit. zool. 3. Will. ichth. p. 97. Bloch. t. 46.

THE Dab is of a very broad, ovate shape, of a yellowish brown colour above, and white beneath: it is covered with moderately large* rough scales: the head is small, and the eyes large: the mouth small, and the teeth are more numerous in the upper jaw than in the lower: the dorsal and anal fins are of moderate width, and the tail nearly even at the end: the lateral line curves downwards over

* Mr. Pennant calls them small, but they are certainly rather large in comparison with those of the major part of the genus.

the pectoral fins, and from thence runs strait to the tail.

This species is an inhabitant of the Mediterranean, the Baltic, and the Northern seas, but is less common than either the Plaise or Flounder, to both of which it is superior as a food, though inferior in its general size: it is in its highest season in the months of February, March, and April, after which it is observed to grow less firm: it spawns in May, or, if the spring prove cold, in June. Dr. Bloch observes that the small spines, mentioned by Linnæus as situated at the base of the dorsal and ventral fins, were not observable in the specimens which he examined.

SMEAR-DAB.

Pleuronectes lævis. *P. fusco-flavescens squamis lævibus, subtus albus, maculis quinque fuscis.*

Yellowish-brown smooth-scaled Flounder, beneath white with five dusky spots.

Rhombus lævis Cornubiensis maculis nigris? (a Kit, Mr. Jago). *Ran. syn. pisc.* 162. f. 1.

Smear-Dab. *Penn. brit. zool.* 3.

“ We found (says Mr. Pennant) one of this species at a fishmonger’s in London, where it is known by the name of the *Smear-Dab*. It was a foot and a half long, and eleven inches broad between fin and fin on the widest part: the head appeared very small, as the dorsal fin began very near its mouth, and extended very near to the tail: it consisted of seventy-nine rays: the eyes were

pretty near each other: the mouth full of small teeth: the lateral line was much incurvated for the first two inches from its origin, then continued strait to the tail: the back was covered with small smooth scales, was of a light brown colour, obscurely spotted with yellow: the belly white, and marked with five large dusky spots: it was a fish of goodness equal to the common Dab."

LONG DAB.

Pleuronectes Limandoides. P. fusco-flavescens, squamis magnis asperis, linea laterali lata, recta.

Yellowish-brown Flounder, with large rough scales, and strait, broad lateral line.

Pleuronectes Limandoides. P. corpore oblongo asperoque, linea laterali recta lataque. Lin. Syst. Nat. Gmel. p. 1232. Bloch. t. 186.

Of a much longer shape than the Dab, and more approaching to that of the sole: colour pale ferruginous brown above, white beneath: scales larger in proportion than those of the common Dab, and rough: lateral line strait and broad: eyes large; mouth wide, and beset with several rows of small teeth: dorsal and anal fins moderately broad: tail rounded at the end. It is an inhabitant of the northern seas, and like others of this genus, commonly lies at the bottom, where it preys on small crabs, &c. and is in considerable estimation as a food.

FLOUNDER.

Pleuronectes Flesus. *P. fuscus*, *linea laterali aspera, spinulis ad radices pinnarum.*

Brown Flounder, with rough lateral line, and a series of spines at the base of the fins.

Pleuronectes Flesus. *P. oculis dextris, linea laterali aspera, spinulis ad pinnas.* *Lin. Syst. Nat. p. 457.*

The Flounder. *Will. ichth. p. 98. Penn. brit. zool. 3. Block. t. 44.*

THE Flounder is allied to the Plaise in shape, but is generally of smaller size and of more obscure colours, the upper side being of a dull brown, marbled with paler and darker variegations, and the under side of a dull white, sometimes obscurely varied with brown: the body is covered with very small scales, and along the back, at the base of the dorsal fin runs a row of small sharp spines: a similar row runs along the base of the anal fin: the lateral line is marked by a third row, continued almost to the base of the tail, which is slightly rounded at the end: at the commencement of the anal fin is a pretty strong spine.

The Flounder is an inhabitant of the Northern, Baltic, and Mediterranean seas: about our own coasts it is extremely common, and even frequents our rivers, at a great distance from the salt waters: it is in considerable esteem as a food, though much inferior to some others of the genus.

The *Pleuronectes Passer*, of Linnæus, considered by Mr. Pennant and some others as a variety of the Flounder, having the eyes on the left side, is

at present allowed to constitute a distinct species, and will of course be described in its proper division.

ROSE-COLOURED FLOUNDER.

Pleuronectes Roseus. P. oculis dextris, corpore roseo.

Rose-coloured Flounder, with eyes towards the right. *Naturalist's Miscellany. Vol. 7. pl. 238.*

GENERAL proportions those of a Flounder: length about nine inches: colour most delicate rose, slightly tinged in some parts with yellowish, and in others with silvery white: lower surface paler, or very nearly white: fins and tail pale yellow-brown: skin apparently destitute of scales, though marked by very minute scale-like reticulations, and void of all asperity either on the side-line or at the origin of the fins. Taken in the Thames, and preserved in the Leverian Museum.

SOLE.


Pleuronectes Solea. P. oblongus fuscus, squamis asperis, maxilla superiore longiore.

Oblong brown Flounder, with rough scales, and upper jaw longer than the lower.

Pleuronectes Solea. P. oculis dextris, corpore aspero oblongo, maxilla superiore longiore. Lin. Syst. Nat. p. 456.

Buglossus. The Sole. *Will. ichth. p. 100. Penn. brit. zool. 3. Bloch. t. 45.*

THE Sole is an inhabitant of the Northern, Baltic, Mediterranean, and American seas, and



ROSE-COLOURED FLOUNDER.

and sold at London, England, by the London Fish Store.

grows to the length of more than two feet, and to the weight of eight pounds: its general size however is much smaller: its shape is that of a very long oval: its colour obscure brown above, and white beneath: it is covered with small rough scales of an oblong form, each terminated by numerous spines, and very strongly fastened to the skin: these scales, from the elegance of their structure, form a favourite microscopic object, and an erroneous idea sometimes prevails that the spiny end of the scale is that by which it was inserted into the skin: the pectoral fins, especially that on the upper side, are commonly tipped with black: the head is small; the eyes and mouth of moderate size; the upper jaw longer than the lower, and both fringed round the edges with a row of very small, white beards.

Next to the Turbot, this fish is considered as the most delicate of the genus, and is by many even preferred to the former; the flesh being remarkably firm, white, and well-flavoured: those of moderate size are, in general, most esteemed. The Sole delights in lying at the bottom of the coasts which it frequents, preying on small shell-fish, spawn, sea-insects, &c. and is generally taken by the trawl-net: the chief fishery, according to Mr. Pennant, is at *Brixham* in *Torbay*.

VENETIAN SOLE.


Pleuronectes Linguatula. *P. oblongus subfuscus, squamis asperis, ano sinistro, dentibus acutis.* *Lin. Syst. Nat. Gmel. p. 1233.*

Brownish oblong Flounder, with rough scales, vent on the left side, and sharp teeth.

Pleuronectes Linguatula. *P. oculis dextris, ano sinistro, dentibus acutis.* *Lin. Syst. Nat. p. 457.*

Linguatula. *Will. ichth. Bellon. &c. ?*

IF this be the *Linguatula* of Bellonius and Willughby, it is thus described. In figure it resembles the Sole, but is not above half as large, and is somewhat shorter in proportion: its colour is paler, and its scales much larger: the vent is placed on the under side, and not on the margin: the mouth is very wide; the tongue round and long; the teeth sharp: the eyes are placed near each other and of moderate size: the iris is gold-coloured: the pectoral and ventral fins have the same number of rays with those of the Sole: the dorsal has sixty-seven rays, and the anal forty-five. This species is said to be found about the coasts of Italy, and Willughby mentions his having seen it at Rome and at Venice.



LARGE-SCALED SOLE.



ZEBRA SOLE.



—Bull. Fish. & Fisheries, 1881, Vol. 1, No. 1, p. 1, Plate 1.

PALLASIAN SOLE.

Pleuronectes Glacialis. *P. lævissimus, supra fuscus, subtus albus, pinnæ dorsalis analisque radiis mediis spinis minimis hispidis.* *Lin. Syst. Nat. Gmel. p. 1234. Pall. it. 3. p. 706.*

Very smooth brown Flounder, white beneath, with the middle rays of the dorsal and anal fin roughened by minute spines.

INHABITS the frozen sea, frequenting sandy places: length about nine inches: the bony part of the head, behind the eyes, is prominent and rough: body perfectly smooth, of a brown colour above, and white beneath: observed by Dr. Pallas.

ZEBRA SOLE.

Pleuronectes Zebra. *P. oblongus albidus, fasciis pluribus transversis duplicatis fuscis.*

Whitish oblong Flounder, with several transverse double brown bands.

Pleuronectes Zebra. *P. fasciis transversis plurimis.* *Lin. Syst. Nat. Gmel. p. 1226. Bloch. t. 187.*

AN elegant species, and readily distinguishable by its colours, the upper part being white, with a brownish cast towards the back, and marked from head to tail by numerous, double, deep brown, transverse bands, passing across the fins themselves: the body is rather longer in proportion than in the common Sole, and the dorsal and anal fin are continued into the tail which is of a slightly pointed shape: the scales are rough and toothed.

at their edges, and the lateral line strait. Native of the Indian seas, and in considerable esteem as a food.

SLENDER-FINNED FLOUNDER.

Pleuronectes Trichodactylus. *P. griseus, fusco variegatus, pinnis pectoralibus filiformibus.* *Lin. Syst. Nat. Gmel. p. 1226.*
Grey Flounder, with dusky variegations, and filiform pectoral fins.

Pleuronectes asper canescens, pinnis lateralibus vix conspicuis.
Artedi gen. 18. syn. 33. spec. 60.

AN uncertain species, known only from the description given by Artedi, who says that it is rough, of a grey colour with dusky variegations; that the eyes are towards the right, and that the pectoral fins are so slender as to be scarce conspicuous, consisting only of four filiform rays on the coloured side of the body, and of one only on the under side. Native of the Indian seas, and found about Amboina.

CAROLINA SOLE.

Pleuronectes Plagiusa. *P. oblongus griseo-fuscus, pinna dorsali cauduli analique unitis.*

Grey-brown oblong Flounder, with the dorsal, caudal, and anal fins united.

Pleuronectes Plagiusa. *P. corpore oblongo scabriusculo, pinna dorsali analique caudæ unitis.* *Lin. Syst. Nat. Gmel. p. 1227.*

ALLIED in habit to the common Sole, but has the dorsal and anal fins united with the tail, as in

the Zebra Sole : colour greyish brown : scales roughish. Native of the American seas : observed about Carolina by Dr. Garden.

OCELLATED SOLE.

Pleuronectes Ocellatus. *P. corpore ocellis quatuor atris, iridibus albis. Lin. Syst. Nat. Gmel. p. 1227. Mus. Ad. Frid. 2. p. 68.*
Sole marked with four black spots with white irides.

NATIVE of Surinam : described by Linnæus in the second part of the Museum Adolphi Friderici : dorsal fin plaited : tail marked by a black bar.

RONDELETIAN SOLE.

Pleuronectes Rondeletii. *P. subfuscus, maculis magnis distantibus subnigris, pupillis pallidis.*

Brownish Sole, marked with large, distant, blackish spots with pale pupils.

Solea oculata Rondeletii. *Will. ichth. p. 100. t. F. 8. f. 4.*

SLIGHTLY described by Rondeletius, who represents it as by no means uncommon about Marseilles : shape like that of the common Sole : scales very closely fixed : body marked above by seven large dusky ocellated spots, of which that in the middle is largest : in the figure of this fish given by Rondeletius these spots appear perfectly well defined, round, and with pale or whitish centres.

PLATESSOID FLOUNDER.

Pleuronectes Platessoides. P. oblongus griseo-fuscus, squamis magnis asperis, corpore supra maculis brunneis notato.

Oblong grey-brown Flounder, with large rough scales, and body marked above by rufous-brown spots.

Pleuronectes platessoides. P. corpore supra maculis brunneis. Lin. Syst. Nat. Gmel. p. 1234.

NATIVE of the seas about Greenland, chiefly frequenting the sandy mouths of rivers, and is not a common fish: length about a foot: head smooth: eyes middle-sized, with silvery irides: lower jaw longer than the upper, and terminated by a tubercle: tongue slender, smooth, and long: pectoral and ventral fins small: tail large and roundish: general shape like that of the *P. Linguatula*; scales like those of the common Sole, and the colour like that of the Plaice, viz. greyish brown above, with fulvous spots, and white beneath. Observed by Otto Fabricius during his residence in Greenland.

SILVER SOLE.

Pleuronectes Argenteus. P. argenteus, subtus subfuscus.

Silvery Sole, brownish beneath.

Petiv. gazoph. 10. t. 26.

SLIGHTLY described by Petiver: shape like that of the common Sole: upper jaw longer than the lower: lateral line strait: upper side of the body of a silvery colour; under side pale brown: native of the Indian seas.

SMOOTH SOLE.

Pleuronectes Diaphanus. *P. candidus diaphanus, squamis tenuissimis deciduis.*

White, transparent Sole, with small, thin, deciduous scales.
 Arnoglossus, vel Solea lævis Rondeletii. *Will. ichth. p. 102.*
 Smooth Sole. *Penn. brit. zool. 3.*

DESCRIBED by Ray, from Rondeletius and others, as extremely thin, pellucid, and white, and covered with scales so minute, and those immediately deciduous, as to merit the appellation of *smooth sole*: it is said to be a rare species, and to be found about the coasts of Cornwall, where, from its transparency, it is called the *lantern-fish*.

 BEARDED SOLE.

Pleuronectes Barbatus. *P. oblongus lævis, maxillis utrisque latere albo cirrosis, pinnis pectoralibus nullis, corpore albo-maculoso. Gronov. zooph. 1. p. 75.*

Oblong, smooth Sole, with whitish spots, both jaws deeply bearded on the left side, and no pectoral fins.

DESCRIBED by Gronovius: shape similar to that of the common Sole, but rather wider in proportion: lips bearded on the left or uncoloured side, the series of beards reaching from the under jaw to the beginning of the ventral fins: colour, according to Gronovius, brown with whitish circular spots: native region unknown.

MARBLED SOLE.

Pleuronectes Marmoratus. P. fuscus, maculis lineisque flexuosis lacteis.

Brown Sole, with milk-white spots and flexuous streaks.

L'Achire marbré. *Cepede 4. p. 660.*

SLIGHTLY described by Cepede from Commerson: shape like that of the common Sole: colour, on the upper side brown, variegated or marbled with differently shaped milk-white spots and marks, some being circular, others long and confluent: fins varied with grey and blue, and speckled with black points: scales so small as to be difficultly seen: no pectoral fins: along the base of the dorsal and anal fins a row of pores, answering in number to the rays of the fins: from these, in the living animal, Commerson observed a whitish mucus to exsude on pressure: native of the Indian seas, and much esteemed as a food.

PAVONIAN SOLE.

Pleuronectes Pavoninus. P. fuscus, maculis irregularibus albidis, mediis fuscis.

Brown Sole, with irregular whitish spots with brown centres.

L'Achire pavonien. *Cepede 4. p. 660.*

OBSERVED by Cepede in the collection of the Prince of Orange: shape like that of the two preceding species; and equally distinguished by the want of pectoral fins: body brown on the upper

side, and marked by spots somewhat resembling in shape those on the tail of a peacock, being white with brown centres: probably a native of the Indian seas.

This species (if such it be) together with the two preceding, the lineated, the bilineated, and the ornamented Sole, is unnecessarily placed by Cope in a distinct genus, under the title of *Achirus*, from the circumstance of wanting the pectoral fins.

LINEATED FLOUNDER.

Pleuronectes Lineatus. *P. corpore fusco, scabro, nigro transversim fasciato, pinnis pectoralibus nullis.*

Brown, rough Flounder, with transverse black lines, and without pectoral fins.

Pleuronectes lineatus. *P. oculis sinistris, corpore scabro nigro fasciato, pinnis pectoralibus nullis.* *Lin. Syst. Nat. p. 458.*

Passer lineis transversis. *Sloane Jam. 2. p. 77.*

FORM that of a Turbot: length seven or eight inches: body covered with small rough scales, and of a deep brown above, marked by seven or eight transverse black streaks: the rays of all the fins and tail roughened by small shagreen-like tubercles: under part of the body white, marked with round, blackish spots. Native of the American seas.

With eyes towards the left.

TURBOT.

Pleuronectes Tuberculatus. *P. griseo-fuscus, subnigro variatus, corpore tuberculis muricato.*

Grey-brown Flounder, with blackish variegations, and body roughened with tubercles.

Pleuronectes maximus. *P. oculis sinistris, corpore aspero.* Lin. *Syst. Nat. p. 459.*

Pleuronectes tuberculis osseis scaber. Bloch. *It. 49.*

The Turbot or Bret. *Will. ichth. p. 94.* *Penn. brit. zool. 9.*

THE Turbot, generally considered as superior to every other species as an article of food, is an inhabitant of the Mediterranean and Northern seas, where it often arrives at a very large size: it is however far inferior in this respect to the Holibut, and is therefore not very happily distinguished by Linnæus under the name of *Pleuronectes maximus*. It is of a broader and squarer form than any other of the genus except the Pearl, and is of a dark brown above, marbled with blackish spots of different sizes; and white beneath: the scales are so small as to be scarcely observable, but the skin is of a wrinkled appearance, and covered with pretty numerous and moderately large pointed tubercles or abrupt spines, those on the upper or coloured side being far larger than those on the under side: the lateral line forms an arch over the pectoral fins, and from thence runs strait to the tail.

Like the rest of this genus, the Turbot generally lies in deep water, preying on worms, shell-fish,

and marine insects, as well as on various kinds of small fishes: it is taken in great quantities about the northern coasts of England, as well as on those of France, Holland, &c. and is baited for with pieces of herring, haddock, &c. but more particularly with the smaller or river lamprey, vast quantities of which are said to be purchased of our fishermen by the Dutch, to the annual amount of not less than seven hundred pounds: they are chiefly taken about Mortlake, and sold to the Dutch as bait for the cod-fishery, but that people are said to have the art of preserving them till the commencement of the turbot-fishery.

The general manner in which the Turbot-fishery is practised at *Scarborough* is thus detailed by Mr. Pennant in the *British Zoology*, from the communications of Mr. Travis of that place.

“ When they go out to fish, each man is provided with three lines: each man’s lines are fairly coiled upon a flat, oblong piece of wicker-work; the hooks being baited, and placed very regularly in the centre of the coil: each line is furnished with fourteen score of hooks, at the distance of six feet two inches from each other: the hooks are fastened to the lines upon snoods of twisted horse-hair seventeen inches in length: when fishing, there are always three men in each coble, and consequently nine of these lines are fastened together, and used as one line extending in length near three miles and furnished with 2520 hooks: an anchor and a buoy are fixed at the first end of the line, and one more at each end of each man’s lines, in all four

anchors, which are commonly perforated stones, and four buoys, made of leather and cork: the line is always laid across the current: the tides of flood and ebb continue an equal time upon our coast, and when undisturbed by winds, run each way about six hours: they are so rapid that the fishermen can only shoot and haul their lines at each turn of the tide; and therefore the lines always remain on the ground about six hours: the same rapidity of the tide prevents their using hand-lines, and therefore two of the people commonly wrap themselves in the sail and sleep, while the other keeps a strict look-out, for fear of being run down by ships, and to observe the weather; for storms often rise so suddenly, that it is with extreme difficulty they can escape to shore, leaving their lines behind. The coble is twenty feet six inches long, and five feet extreme breadth: it is about one ton burthen, rowed with three pair of oars, and admirably constructed for the purposes of encountering a mountainous sea: they hoist sail when the wind suits."

PEARL.

Pleuronectes Rhombus. P. fuscus, subflavo maculatus, corpore glabro.

Brown Flounder, with yellowish spots, and smooth body.

Pleuronectes Rhombus. P. oculis sinistris, corpore glabro.
Lin. Syst. Nat. p. 452.

Pleuronectes corpore lato et glabro. Bloch. t. 36.

The Pearl *Will. ichth. p. 95. Penn. brit. zool. 3.*

THE Pearl is distinguished from the Turbot by the perfect smoothness of its skin, which is covered with scales of moderate size, and is of a pale brown colour above, marked by scattered yellowish or rufous spots, and white beneath: the lateral line, as in the Turbot, is first arched over the pectoral fins, and from thence runs strait to the tail: in its general form this fish resembles the Turbot, but is usually inferior in size.

The Pearl is by most modern authors supposed, perhaps erroneously, to have been the Rhombus of the ancient Romans, of which a specimen of enormous size is said to have been taken in the reign of Domitian*, who ludicrously ordained a *Senatusconsultum* relative to the mode of bringing it to table.

* *Juvén. sat. 4.*

WHIFF.

Pleuronectes Punctatus. *P. fuscus, maculis nigricantibus, punctis rubris, squamis asperis.*

Brown Flounder, with blackish spots, red freckles, and rough scales.

Pleuronectes punctatus. *P. corpore lato asperoque.* *Lin. Syst. Nat. Gmel. Bloch. t. 189.*

The Whiff. *Penn. brit. zool. 3. ?*

THIS, says Mr. Pennant, bears some resemblance to the Holibut: the length of a specimen examined by that author was eighteen inches; the greatest breadth not seven, exclusive of the fins: the mouth was extremely large; the teeth very small; the under jaw hooked over the upper; the eyes were large; the scales large and rough; the side-line tuberculated and uncommonly incurvated at the beginning, making a sharp angle, and then going straight to the tail, which is rounded: the colour of the body was cinereous brown, clouded in parts, and obscurely spotted: the under side white, tinged with red.

The figure of this species in the work of Dr. Bloch is represented as of a dusky brown above, with numerous, rather large, roundish black spots of unequal sizes, between which are interspersed several very small red spots. In these particulars it seems to agree much better with the description contained in Ray's synonym than the *Smear-Lad*, which Mr. Pennant supposes to be the *Kit* of the Cornish fishermen.

BILINEATED SOLE.

Pleuronectes Bilineatus. *P. fuscus*, pinna dorsali, anali, caudaliq^{ue} unitis, lineis lateralibus duabus.

Brown Sole, with dorsal, anal, and caudal fin united, and two lateral lines.

Pleuronectes bilineatus. *P. lineis lateralibus duabus.* Bloch, t. 188.

SHAPE resembling that of the Zebra Sole: colour above pale brown, beneath rufous white: scales small, round, and denticulated: head larger than in the rest of the Soles: mouth small: dorsal and anal fin united, as in the Zebra Sole, with the tail, which is of a slightly pointed shape: native of China: particularly distinguished by having two lateral lines, one running strait along the middle, the other near the back.

 ORNAMENTED SOLE.

Pleuronectes Ornatus. *P. subfuscus*, maculis ocellatis albis medio fuscis.

Brownish Sole, with white ocellated spots with brown centres.

Achirus ornatus. *Cepede* 4. p. 663.

SHAPE like that of the preceding: colour pale brown above, with eight or nine deep-brown transverse bands: scales small, round, and denticulated: lateral line running in the direction of the back: observed by Cepede in the collection of the Prince of Orange.

LARGE-SCALED SOLE.

Pleuronectes Macrolepidotus. *P. fuscus, squamis magnis, pinnis pectoralibus lanceolatis.*

Brown Sole, with large scales, and lanceolate pectoral fins.

Pleuronectes macrolepidotus. *P. squamis magnis.* Bloch. t. 190.

SHAPE like that of a Sole, but somewhat broader in proportion: colour yellowish brown above, whitish beneath: scales very large, each marked at the base by a dusky spot: head without scales and spotted with brown: mouth very wide: teeth rather long: pectoral fins rather long and pointed: ventral very small: dorsal, anal, and caudal like those of the common Sole. Native of the South-American seas.


SINISTROUS FLOUNDER.

Pleuronectes Passer. *P. fuscus, oculis sinistris, linea laterali aspera, spinulis ad radices pinnarum.*


Brown Flounder, with eyes toward the left, rough lateral line, and a series of spines at the base of the fins.

Pleuronectes Passer. *P. oculis sinistris, linea laterali sinistra aculeata.* Lin. Syst. Nat. p. 459. Bloch. t. 50.

So nearly allied in general appearance to the common Flounder that it has been considered by Pennant and some others as a variety of that species differing merely in having the eyes towards the left: Artedi, Linnæus, and Bloch however concur in establishing it as a distinct species: the



ARGUS FLOUNDER.



TURBOT.

lateral line on the upper side is rough or spiny, as in the Flounder, as also the bases of the dorsal and anal fin, but the spines on those parts are all strait, whereas in the former species they are intermixed with curved ones: in colour it resembles the Flounder.

 ARGUS FLOUNDER.

Pleuronectes Argus. *P. albidus, maculis rotundis flavis caruleo marginatis.*

Whitish Flounder, with round yellow spots margined with blue.

Pleuronectes Argus. *P. corpore vario, pinna cauda rotundata.*

Bloch. t. 48. Linn. Syst. Nat. Gmel. p. 1239.

Passer oculatus. Plumier, Mss.

THIS elegant species is of the same general form with the Turbot, and is of a pale yellow, or rather a yellowish white colour on the upper side marked by numerous eye-shaped spots, consisting of bright-blue circles with yellow centres: the whole skin is also marked both on the body and fins with small blue and brown specks, and is covered with small scales: the larger spots first mentioned vary somewhat in different individuals, as well as on different parts of the same fish, being sometimes, formed by complete circles of blue, but more commonly by two or three broken crescent-shaped marks of that colour: the under side is of a whitish or pale grey colour: the lateral line is arched over the pectoral fins, and is thence continued strait to the tail,

which is rounded at the tip. This beautiful fish is a native of the American seas, and some of its varieties seem to have been considered by authors as distinct species.

VAR. ?

Pleuronectes lunatus. *P. corpore ocellis dimidiatis sparsis caeruleis*. *Lin. Syst. Nat. Gmel. p. 1239. Solea lunata et punctata. Catesb. Car. 2. t. 27.*

OBSERVED by Catesby, who in his Natural History of Carolina represents it as of a pale brown colour, with numerous blue crescent-shaped spots, and lunated tail.

VAR. ?

Pleuronectes Mancus. *P. capite tuberculato, pinnis pectoralibus inaequalibus*. *Lin. Syst. Nat. Gmel. p. 1238. Brouss. ichth. dec. 1. t. 3. 4.*

BODY of an oblong elliptic form, covered with rather large scales: right side and fins cinereous, with blackish specks and unequal, blueish, lunated and circular variegations: left side greyish white, with round brown spots: pectoral fin on the left side falcate, on the right oblong-ovate. Native of the American and Indian seas.

STELLATED FLOUNDER.

Pleuronectes Stellatus. *Pl. fuscus, tuberculis numerosissimis albidis spinoso-radiatis.*

Brown Flounder, with very numerous whitish tubercles radiated with spines.

Pleuronectes stellatus. *Pall. nov. act. Petrop. 1787.*

THIS elegant species is frequently found, according to Dr. Pallas, about the mouths of rivers in the Curile islands. Its usual length is supposed to be somewhat less than a foot and half; or nearly that of a Plaise, which it also resembles in shape: the eyes are on the left side, which is of a brown colour, and very thickly mucronated all over with small tubercles of a whitish cast, smooth in the centre but radiated round the edge with small spines: the under side is white, and the fins are pale with dusky or blackish streaks here and there in the direction of the rays.

CAROLINA SOLE.

Pleuronectes Dentatus. *P. subfuscus, dentibus acutis exsertis.*

Brownish Sole, with sharp, exserted teeth.

Pleuronectes dentatus. *P. corpore oblongo glabro, dentibus exsertis.* *Lin. Syst. Nat. Gmel. p. 1236.*

OBSERVED about Carolina by Dr. Garden: teeth sharp; tail rounded and scaly: in the dorsal fin are eighty-seven rays, in the pectoral twelve, in the ventral sixty-six, and in the caudal seventeen.

JAPANESE FLOUNDER.

Pleuronectes Japonicus. P. lingua aspera. Lin. Syst. Nat. Gmel. p. 1240. Houttuyn. act. haarl. 20. 2. p. 317.
Flounder with rough tongue.

LENGTH about six inches; colour brownish above, whitish beneath: rays of the dorsal and anal fin so extremely numerous as to be with difficulty counted: tongue remarkably rough: native of the Japanese seas.

SCHÆTODON. CHÆTODON.

Generic Character.

Caput parvum : *Os* exiguum : *Dentes* conferti, flexiles, setacei.

Membr. branch. : radii tribus ad sex.

Corpus latum, compressum, pluribus fasciatum ; pinna dorsi anique basi crassa, carnosae, squamosae.

Head small : *Mouth* small : *Teeth* close-set, flexible, setaceous.

Gill-Membrane three, four, five, or six-rayed.

Body broad, compressed, and, generally, fasciated : *dorsal* and *anal* fin, thick, fleshy, and scaly at the base.

THE numerous genus *Chætodon* is, in a general view, distinguished by the great depth and highly compressed form of the body, which, in most species, is beautifully variegated by transverse, oblique, or longitudinal bands, and covered with strong scales, finely denticulated on the margins : the dorsal and anal fin are remarkably broad, scaly at the base or lower part, and in many species, of an unusual length, and furnished, as in some of the *Dories*, with rays projecting beyond the membrane : the teeth are, in general, small, very slender, and flexible. The species are very numerous, and are mostly natives of the Indian and American seas, being very rarely, if ever, taken in those of Europe.

With single dorsal fin, and even or rounded tail, or at most, but very slightly inclining to crescent-shaped in some few species.

IMPERIAL CHÆTODON.

Chætodon Imperialis. *C. flavus, lineis numerosissimis longitudinalibus obliquis cæruleis.*

Yellow Chætodon, with very numerous oblique longitudinal blue lines.

Chætodon Imperator. *Lin. Syst. Nat. Gmcl. p. 1255. Bloch. t. 194.*

A MOST magnificent species, growing to the length of a foot or more : body oval ; dorsal and anal fin broad, and scaled to a considerable distance from the base : gill-covers furnished on each side with a very strong spine : ground-colour of the animal golden yellow, longitudinally, but somewhat obliquely striped with very numerous bright-blue parallel rays. Native of Japan, where it is held in high esteem as an article of food, and is said to be superior to the Salmon in flavour. .

17

IMPERIAL, CHARTERED.

1802, July 1. London. Purchased by W. Brewster. First sold.

Efficient, on-line

PARU CHÆTODON.

Chætodon Paru. *C. nigricans, squamis aureo marginatis.*

Blackish Chætodon, with the scales bordered with a golden gloss.

Chætodon Paru. *Lin. Syst. Nat. Gmel. p. 1256. Bloch. t. 197.*

A HIGHLY elegant species, though inferior in beauty to the preceding: grows to about the same size, and is of an oval shape, the dorsal and anal fins extending at their tips far beyond the thicker or scaly part of the fin: on the gill-covers a strong spine: colour of the whole animal black, accompanied by a strong golden tinge, most conspicuous toward the edges of the scales, which are large and rounded: native of the South-American seas.

 ANGEL CHÆTODON.

Chætodon Catesbeii. *C. viridi-aureus, squamis squamulosis.*

Gold-green Chætodon, with the scales covered by smaller ones.

Angel-Fish. *Catesb. carol. 2. t. 31.*

DESCRIBED and figured by Catesby in his Natural History of Carolina; who names it the Angel-Fish, and represents it of a golden-green colour, with the pectoral, ventral fins, and tail of a bright orange-colour; the dorsal and anal violet-blue at the base, and bright crimson towards the tips: these fins extend to a great distance, reaching far beyond the tail: the body is very broad, terminating very

abruptly on the hind part, from the middle of which the base of the tail suddenly emerges: the gill-covers are armed with several strong spines, and the whole body is covered with moderately large rounded scales, which are themselves covered with much smaller ones. Native of the American seas, and according to Catesby, common off Carolina and the Bahama isles, where it is much esteemed for its delicacy.

CILIATED CHÆTODON.

Chætodon Ciliaris. *C. suburgenteus, squamis ciliatis.*

Silvery-grey *Chætodon*, with ciliated scales.

Chætodon ciliaris. *Lin. Gmel. p. 1252. Bloch. t. 214.*

LENGTH about a foot: body oval; dorsal and anal fin extended into sharp semilunar extremities, projecting beyond the tail: colour of the whole animal silvery-grey, with a more dusky tinge on the back, and an annular black spot placed immediately before the origin of the dorsal fin: scales of moderate size, and all ciliated round their edges with numerous hair-like processes, which take their rise from the middle of each scale: gill-covers furnished at the middle with a strong spine, accompanied with a few smaller ones. Native of the Indian seas.

FASCIATED CHÆTODON.

Chætodon Fasciatus. *C. corpore oblongo-ovato albido, fasciis pluribus transversis cæruleis.*

Chætodon with oblong ovate body, marked by several transverse blue bands.

Chætodon fasciatus. *Bloch. t. 195.*

Chætodon Dux. *Lin. Gmel. p. 1255.*

A BEAUTIFUL species: shape oblong-ovate; colour white with a slight silvery cast, and a dusky tinge on the upper parts: body most elegantly banded by several broad, transverse, deep blue bands with brown edges: dorsal and anal fin moderately broad, but obtuse at their extremities, and not extending to the end of the tail: at the gill-covers a very strong spine. Native of the Indian seas.

SURAT CHÆTODON.

Chætodon Suratensis. *C. corpore subrotundo-ovato subargenteo, fasciis transversis fuscis, maculaque utrinque pectorali nigra.*

Silvery-grey Chætodon with roundish-ovate body banded with brown, and marked on each side by a black pectoral spot.

Chætodon Suratensis. *Bloch. t. 217.*

SHAPE roundish-oval: colour silvery grey, darker on the back and fins: body marked by six or seven moderately broad, transverse, brown bands, and at the base of the pectoral fins by a semi-lunar black spot: dorsal and anal fin broadish at the extremities, but obtuse, and not reaching to

the tip of the tail: scales moderately large, and each marked in the middle by a bright spot. Native of the Indian seas, growing to the length of ten or twelve inches.

KLEINIAN CHÆTODON.

Chætodon Kleinii. *C. flavescens, corpore rotundato-ovato, capite fascia transversa nigra.*

Yellowish Chætodon, with round-ovate body, and head crossed by a black band.

Chætodon Kleinii. *Bloch. t, 218. f. 2.*

SIZE rather small: shape round-ovate: colour pale yellow, with a dusky tinge on the back, and a silvery cast on the sides: from the origin of the dorsal fin to that of the pectoral fins, runs a slightly curved, blackish, or deep-brown stripe, passing through the eye in its course: dorsal and anal fin rather broad at the ends, but not projecting so far as the end of the tail: scales large and rounded. Native of the Indian seas.

EIGHT-BANDED CHÆTODON.

Chætodon Octofasciatus. *C. subargenteus, corpore rotundato-ovato, fasciis octo angustis transversis nigris.*

Silvery-grey *Chætodon*, with round-ovate body crossed by eight narrowish black bands.

Chætodon octofasciatus. *Lin. Gmel. p. 1262. Bloch. t. 215. f. 1.*

Size rather small: shape, round-ovate: colour whitish or silvery-grey, brownish on the back: body marked by eight rather narrow, blackish or deep-brown, transverse bands disposed in pairs, viz. one pair across the head and gills, the first line passing through the eye: two pair across the middle of the body, and the remaining pair across the end of the body and tail: lateral line curving in the direction of the back, and formed by a row of round scales: dorsal fin almost completely scaled, and furnished at the fore-part with extremely strong spines; rounded at the hind-part and slightly stretched towards the tail, but not reaching farther than the middle of that part: anal fin of similar shape; and furnished with three very strong short spines at its origin. Native of the Indian seas. This species was arranged by Linnæus, in the twelfth edition of the *Systema Naturæ*, in the genus *Perca*, under the name of *Perca nobilis*.

ANNULAR CHÆTODON.

Chætodon Annularis. *C. luteus*, corpore rotundo-ovato, fasciis sex obliquis longitudinalibus curvatis annuloque postoculæri cæruleis.


Deep yellow *Chætodon*, with round-ovate body, marked by six longitudinal curved blue stripes and a blue ring behind the eye.

Chætodon annularis. *Lin. Gmcl. p. 1262. Bloch. t. 215.*

SHAPE round-oval: colour yellow or pale fulvous, with six oblique, longitudinal, curved, bright-blue lines or narrow stripes, of which the four superior are continued into the dorsal fin, at the hind part of which they meet in an angle: the two lower stripes run along the bottom of the abdomen: the dorsal fin is broad at its hind-part, and projects above the tail into an angular form: the anal fin is rounded at the hind-part: on each side the body, beyond the eye, is a bright-blue circle, and at the bottom of the gill-covers is a strong spine. Native of the Indian seas: length about a foot.



WANDERING CHAETODON .



ROSTRATED CHAETODON .



ANGULAR CHAETODON .

COLLARED CHÆTODON.

Chætodon Collaris. *C. corpore rotundo-ovato albido, capite fusco fasciis duabus lacteis, fascia caudali nigra.*

Whitish Chætodon, with round-ovate body, brown head marked by two milk-white bars, and tail crossed by a brown band.

Chætodon collare. *Lin. Gmel. p. 1263. Bloch. t. 216.*

SHAPE round-ovate; colour blueish or greyish white, tinged with brown on the back: head deep-brown, with two milk-white perpendicular stripes, the former immediately before the eyes, the latter behind: the mouth is also white: the dorsal and anal fin are shaped nearly as in the *C. octofasciatus*, and the tail is marked across the middle by a brown bar: the body is covered with large scales. Native of Japan; length of the specimens described by Seba, Bloch, &c. about six inches.

BLACK-AND-WHITE CHÆTODON.

Chætodon Mesoleucos. *C. corpore subrotundo-ovato nigricante, antèrius albo, fasciis capitis nigra.*

Roundish-ovate blackish Chætodon, with the fore-parts white, and a brown bar across the head.

Chætodon mesomelas. *Lin. Gmel. p. 1263.*

Chætodon mesoleucos. *Bloch. t. 116.*

SHAPE roundish ovate, but less so than the preceding species: colour of the body, dorsal, and anal fin deep-brown, but the fore-part of the body

gradually softens into white: down each side of the head, through the eyes, runs a deep-brown band: the pectoral, ventral fins, and tail are white: the body is covered with moderately small scales, and at the lower part of the gill-covers is a very strong spine: the shape of the fins resembles those of the major part of this division. Native of Japan.

ARGUS CHÆTODON.

Chatodon Argus. *C. subargenteus, maculis numerosis rotundis fuscis, corpore subquadrato-ovati.*

Silvery-grey Chatodon with squarish-ovate body, marked with numerous round brown spots.

Chatodon Argus. *Lin. Gmel. Bloch. t. 204.*

GENERAL shape ovate, or rather squarish-ovate: colour silvery-grey, with a brownish cast on the upper parts, and the whole body, as well as part of the fins, marked by numerous round brown spots: scales very small: first rays of the dorsal fin moderately long, those on the middle part much shorter: terminal part both of dorsal and anal fin rounded and slightly dilated: tail even at the end, but not rounded: gill-covers perfectly smooth: native of the fresh waters of India.

BIMACULATED CHÆTODON.

Chætodon Bimaculatus. *C. albidus*, corpore subrotundo-ovato, capite elongato, fascia transversa nigra, pinna dorsali macula nigra alteraque dimidiata.

Whitish Chætodon, with roundish-ovate body, elongated head crossed by a black bar, and dorsal fin marked by a black spot and a second dimidiated one.

Chætodon bimaculatus. *Bloch. t. 219. f. 1.*

SHAPE of the body roundish-ovate, but the head considerably lengthened, so as to form a kind of snout in front: colour white, with a slight cast of brown on the upper part and fins: scales very large: head marked on each side by a black or deep-brown stripe, beginning at the origin of the dorsal fin, and passing through the eyes downwards, under the throat: pectoral and ventral fins pale red: dorsal marked at the base of the hind-part, by a large, round black spot, bordered with white, and at its extreme edge by a much smaller half-spot of similar colours: native of the seas about Japan.

STRIPED CHÆTODON.

Chætodon Striatus. *C. albidus, corpore subrotundo-orato, lineis longitudinalibus subfuscis, fasciisque quatuor transversis fuscis.*

Whitish Chætodon, with roundish body, marked by several longitudinal dusky lines, and by four transverse dusky bars.

Chætodon striatus. *Lin. Gmel. Bloch. t. 1249.*

SHAPE roundish ovate: colour whitish, tinged with yellowish brown on the upper parts, and marked by four distant brown bands, the narrowest of which runs on each side the head, passing through the eyes to the breast; the two next, which are much broader, pass, at nearly equal distances, across the body, and the third crosses the setting on of the tail: the pectoral and ventral fins are of a deep brown, and the dorsal, anal fin, and tail, are deeply edged with the same colour: the body is covered with scales of moderate size, and is marked by several narrow longitudinal dusky streaks between the rows of scales. Native of the Indian and American seas.

DIVIDED CHÆTODON.

Chætodon Bicolor. *C. corpore oblongo-orato, dimidio antice lacteo, postico fusco.*

Chætodon with oblong-ovate body, the anterior half milk-white, the posterior brown.

Chætodon bicolor. *Lin. Bloch. t. 205.*

A SPECIES of moderate, or rather smallish size, and easily distinguishable by the remarkable distribution

of its colours, the fore-part of the body being milk-white; the hind-part reddish brown, passing through the dorsal and anal fins, which are both edged with white: the tail, pectoral, and ventral fins are also white: the gill-covers are furnished with a very strong spine: the dorsal and anal fins are nearly covered with scales, and are of moderate breadth, extending into an acuminate form above and below the tail, but not reaching to the tip: the body is of an oblong-ovate shape. Native of the Indian and American seas.

SINGLE-SPOTTED CHÆTODON.

Chætodon Unimaculatus. *C. argenteus, supra cærulescens, corpore subrotundo-ovato macula utrinque versus dorsum nigra.*

Silvery Chætodon with roundish-ovate body marked on each side towards the back by a black spot.

Chætodon unimaculatus. *Lin. Gmel. Bloch. t. 201.*

SHAPE roundish-ovate: colour silvery white, with a few obliquely transverse yellow streaks, and tinged on the upper parts with blue-grey: body covered with large scales, and marked on each side, towards the back, with a large oval black or deep-brown spot: the head is marked, as in some others of this tribe, by a deep-brown band, running across the eyes: the base of the tail, and the edges of the dorsal and anal fin are also brown: the tail is even at the end, or, at most, so very slightly lunated as to be scarce perceptibly so. Native of the Indian seas.

WANDERING CHÆTODON.

Chætodon Vagabundus. *C. subluteus, corpore subrotundo-ovato, longitudinaliter striato, postice fascia transversa nigra.*

Yellowish Chætodon, with roundish-ovate body longitudinally striated, and marked at the hind-part by a transverse black bar.

Chætodon vagabundus. *Lin. Bloch. t. 204.*

SHAPE roundish-ovate: head lengthened a little in front: body covered with large scales, and of a pale orange-colour, marked with numerous oblique narrow streaks running in the interstices of the scales: over the shoulders are several similar streaks in an opposite direction, or crossing the former: head marked by a black band running across the eyes, as in many others: end of the body, at the setting on of the tail, marked by a similar band, which runs both into the dorsal and anal fin: tail crossed in the middle by a black or deep-brown crescent, and bordered at the tip with the same colour. Native of the Indian seas, and said to be in considerable estimation as a food.

OCCELLATED CHÆTODON.

Chætodon Ocellatus. *C. corpore ovato argenteo, capite fascia nigra, pinna dorsali macula rotunda nigra albo marginata.*

Chætodon with silvery-white ovate body, head crossed by a black bar, and dorsal fin marked by a round black spot edged with white.

Chætodon ocellatus. *Bloch. t. 211. f. 2.*

A SMALL species: length about four inches: shape ovate: colour silvery white, with a few oblique yellow streaks: dorsal and anal fin broad, rounded at the hind part, and very scaly at the base, dorsal marked with a large, round deep-black spot, edged with white, head marked by a transverse black band, running across the eyes. Native of the Indian seas.

ROSTRATED CHÆTODON.

Chætodon Rostratus. *C. corpore subrotundo-ovato albido, fuscis transversis fuscis, rostro longo, pinna dorsali macula nigra.*

Whitish Chætodon, with roundish-ovate body crossed by brown bands, long snout, and dorsal fin marked by a black spot.

Chætodon rostratus. *Lin. Bloch. t. 202. f. 1.*

Chætodon Enceladus. *Nat. Hist. 2. pl. 67. Jaculator. Phil. Trans. 1765.*

SHAPE roundish-ovate: length about six or eight inches: snout lengthened into a tubular form: colour of the body whitish, with a dusky tinge on the back, and marked by five transverse, and nearly equidistant, brown bands, with milk-white edges: the first band, which is narrower than the rest, passes across the head, through the eyes, the three

next across the body, and the last across the base of the tail: the dorsal and anal fin are very broad at the hind part, and the dorsal is marked in that part by a large black spot, bordered with white: the gill-covers are scaly, but destitute of spines.

This fish is a native of the fresh waters of India, and is celebrated for the extraordinary manner in which it takes its prey, which chiefly consists of the smaller kind of flying insects: when it observes one of these, either hovering over the water, or seated on some aquatic plant, it shoots against it from its tubular snout a drop of water, with so sure an aim as generally to lay it dead, or at least stupefied, on the surface. In shooting at a sitting insect, it is commonly observed to approach within the distance of from six to four feet, before it explodes the water. When kept in a state of confinement in a large vessel of water, it is said to afford high entertainment by its dexterity in this exercise, since if a fly or other insect be fastened to the edge of the vessel, the fish immediately perceives it, and continues to shoot at it with such admirable skill as very rarely to miss the mark.

It may not be improper to observe, that this faculty is not entirely confined to the present fish, but takes place in some few others belonging to very different genera.

ORB CHÆTODON.

Chætodon Orbis. *C. corpore orbiculari, radiis anterioribus pinnæ dorsalis elongatis.*

Chætodon with orbicular body, and the first rays of the dorsal fin elongated.

Chætodon Orbis. *Block. t. 202. f. 2.*

A SMALL species: length about four inches: shape of the body remarkably orbicular: colour blueish-grey above, yellowish-white beneath: body covered with large scales: ventral fins narrow and falciform; first rays of the dorsal running beyond the membrane, as in some of the Dories: side-line curving irregularly, with a kind of double descent before it reaches the tail. Native of the Indian seas.

BUTTERFLY CHÆTODON.

Chætodon Capistratus. *C. corpore ovato fulvo, lineis obliquis oppositis notato, postice macula nigra, capite fascia fusca.*

Chætodon with ovate fulvous body, marked by oblique opposite lines, a black spot on the hind-part, and head crossed by a brown bar.

Chætodon capistratus. *Lin. Block. t. 205. f. 2.*

A VERY small species, length about three inches and a half: shape ovate: dorsal and anal fins gradually dilated and rounded at their extremities: ground-colour of the whole animal pale orange, the body being covered with scales of moderate size, and marked on each side the lateral line by

numerous oblique dusky lines, running towards the tail, and forming so many acute angles along the sides of the body : the head is crossed by a brown band passing through the eyes: the base of the tail is crossed by a smaller bar ; the dorsal and anal fin are edged on their broadest part by a brown margin, and on each side the body, a little above the tail, is a large, round, black spot, with a white border. Native of the American seas, and found about the coasts of Jamaica, &c.

CHALYBEAN CHÆTODON.

Chætodon Faber. *C. subargenteus, caeruleo tinctus, pinnis fasciisque corporis sex transversis caeruleis.*

Subargenteous Chætodon, tinged with blue, with blue fins, and six transverse blue bands.

Chætodon Faber. *Bloch. t. 212.*

SHAPE roundish ovate : length about twelve inches : ground-colour silvery white, but with a strong tinge of dusky blue on the upper parts, and marked by six unequal, dark-blue, transverse bands, of which the first, which is narrow, passes across the head, through the eyes ; the next, which is much broader, through the body ; the third passes only half way down from the back ; the fourth completely through the body ; the fifth, which is narrower, across the end of the body ; and the sixth, which is the narrowest of all, across the base of the tail : all the fins are of the same deep-blue colour with the bands, and the hind-parts of

the dorsal and anal fin extend in a somewhat falcated manner, as far as the end of the tail: the body is covered with scales of moderate size, and the gill-covers are destitute of spines. Native of the American seas.

ARCUATED CHÆTODON.

Chætodon Arcuatus. *C. nigricans, fasciis angustis, transversis, arcuatis, niveis.*

Blackish Chætodon, with narrow, arcuated, transverse snow-white bands.

Chætodon arcuatus. *Lin. Bloch. t. 201. f. 2.*

SHAPE of the body ovate; dorsal and anal fin very broad, and terminating in sharpened or slightly falciform extremities, but not extending so far as the tail: colour of the whole fish very deep black-brown, varied by arcuated milk-white transverse bands; which are five in number; the first passing across the mouth, and the next across the thorax, the two next across the body, dorsal, and anal fin, and the last across the tail, which is also tipped with a milk-white margin: gill-covers tipped with a strong spine. Native of the South-American seas: general length from three to six inches.

BIFASCIATED CHÆTODON.

Chætodon Bifasciatus. *C. albo-cærulescens*, corpore suborbiculari, fasciis duabus fuscis, radio quarto pinnae dorsalis longissimo. Blueish-white Chætodon, with suborbicular body marked by two brown bars, and the fourth ray of the dorsal fin extremely long.

Chætodon macrolepidotus. *Linn. Bloch. t. 200. f. 1.*

SHAPE of the body round-oval; or even orbicular: colour blueish grey above, silvery-white beneath: dorsal and anal fin of moderate breadth, and not extending to the end of the tail; but the fourth ray of the dorsal fin, which is very thick and strong, is extended to a vast distance beyond the rest, reaching considerably farther than the tip of the tail, over which it passes in a curved direction: across the body, are two very broad blackish-brown bands, viz: one from the origin of the dorsal fin to the ventral fins; the other from the middle of the dorsal fin to the bottom of the anal; passing through both: this species is the *C. macrolepidotus* of Linnæus; but as the scales are not so large as in some other species since discovered, I have given it a different trivial name: it is a native of the Indian seas, where it is said to arrive at a very considerable size, sometimes weighing upwards of twenty pounds, and to be much esteemed as a food, resembling the Sole in delicacy.

HORNED CHÆTODON.

Chætodon Cornutus. *C. albo-cærulescens, corpore elevato-orbiculari, antice fasciis duabus, postice unica nigris, radio tertio pinnae dorsalis longissimo.*

Blueish-white Chætodon, with high-orbicular body, crossed on the fore-part by two, on the hind-part by one black band, and the third ray of the dorsal fin extremely long.

Chætodon cornutus. Lin. Bloch. t. 200. f. 2.

SHAPE of the body nearly orbicular, but deeper than long, or measuring more from the back to the abdomen than from head to tail: dorsal and anal fins produced in a falciform manner, so as to reach beyond the tail, more especially the dorsal fin, several rays of which, from about the third to the eighth, are gradually extended beyond the length of the rest, though still preserving the outline of the fin; but the third or longest ray curves over the tail to a great distance downwards: colour of the body silvery; with a blueish cast on the upper parts, and marked by three broadish black zones, two passing across the thorax, so as to represent a double band, and the other passing across the hind part of the body, dorsal, and anal fin: tail black, arched by a narrow white crescent near the end: scales on the body rather small: snout rather produced, but in a much less degree than in the *C. rostratus*: over each eye is generally a small pointed or horn-shaped process, which in the younger specimens is scarce perceptible; native of the Indian seas: general length from four to six inches.

BAT CHÆTODON.

Chætodon Vespertilio. *C. albidus, corpore elevato-orbiculari, pinna dorsali analique maximis, falcatis.*

Whitish Chætodon, with high-orbicular body, and extremely large falcated dorsal and anal fin.

Chætodon Vespertilio. *Lin. Bloch. t. 199. f. 2.*

A VERY singular species, and the most remarkable of all others for the vast extent as well as breadth of the dorsal and anal fin, both which nearly equal the body itself in size, and are of a somewhat triangular shape, the points extending, both above and below, to about the same distance with the tip of the tail: the body in shape bears a considerable resemblance to some of the silvery Dories, the head sloping suddenly down and the general form being similar: the colour silvery, with rather small scales: the base of the tail marked by a black bar: native of Japan: length of a specimen figured by Dr. Bloch, about four inches from head to the end of the tail, but the depth, measured from the opposite tips of the dorsal and anal fins, about six inches and a half.



LONG-FINNED CHAETODON.



BORDERED CHAETODON.

Rees's Fishes

LONG-FINNED CHÆTODON.

— **Chætodon Teira.** *C. albus, corpore elevato-orbiculari, pinna dorsali analique magnis, falcatis, longissimis.*

White Chætodon, with high-orbicular body, and large, extremely long, falcated, dorsal and anal fin.

Chætodon Teira. *Lin. Bloch. t. 199. f. 1.*

GENERAL shape similar to that of the preceding species, and the dorsal and anal fin even considerably longer in proportion, but not so broad: colour of the body silvery-white, with rather small scales, and variegated by three broad, transverse black bands, the first of which runs across the head and eyes, the second across the fore-part of the body and dorsal fin, and the third across the hind part of the body, dorsal, and anal fin; the whole forming an appearance singularly elegant: the ventral fins, which are of a lanceolate shape, are also black. This curious fish is a native of the Indian and Arabian seas, and is said to arrive at a considerable size, though the specimens usually seen are but small.

RED-STRIPED CHÆTODON.

Chætodon Setifer. *C. flavescens rubro-striatus, ocello setoquæ ad pinnam dorsi.*

Yellowish Chætodon, with numerous red stripes on the body, and an eye-shaped spot and bristle on the dorsal fin.

Chætodon setifer. *Bloch. t. 446. f. 1.*

A HIGHLY beautiful species: shape of the body round-ovate, but with the head slightly produced: colour pale yellow, with a strong tinge of crimson towards the back, and beautifully marked in different directions, with numerous crimson stripes, some running obliquely across the sides, and others in an opposite direction across the fore part of the back: scales very large: across the head a broad, black band, passing through the eyes: on the hinder or rounded part of the dorsal fin a large round black spot, bordered with white; over which passes a long setiform single ray, reaching beyond the tip of the tail, which is terminated by a black edge, as are likewise the rounded part of the dorsal and anal fin. Native of the Indian seas: length about six inches.

SICKLE CHÆTODON.

— Chætodon Falcula. *C. albo-flavescens, capite fascia curvata nigra, corpore duabus brevibus falciformibus.*

Yellowish-white Chætodon, with a curved black band across the head, and two short sickle-shaped bands across the body.

Chætodon Falcula. *Bloch. t. 426. f. 2.*

In shape and general appearance very nearly allied to the *C. setifer*, but differs in colours, and in the want of the lengthened ray at the hind part of the dorsal fin: colour white, with a tinge of yellow: scales very large, a black bar across the head, as in the preceding: a second, of a curved form crossing the middle of the body, from the dorsal fin downwards, but vanishing before it reaches the abdomen, and a third still more curved, or of a shape somewhat resembling a sickle, crossing the hind-part of the body, from the middle of the dorsal fin, and like the former vanishing before it reaches the abdomen: the base of the tail is crossed by a black bar, and tipped with black. Native of the Indian seas.

CHARIOTEER CHÆTODON.

Chætodon Auriga. *C. albidus, oblique fasciatus, radio pinnæ dorsalis quinto in filum producto.*

White Chætodon, obliquely striped, with the fifth ray of the dorsal fin lengthened into a filament.

Chætodon Auriga. *Forsk. F. Arab. p. 60. Lin. Gmel. p. 1266.*

Body subrhomboidal, compressed, and about five inches long: teeth in several rows: general colour blueish white, with several oblique, parallel, dusky streaks: the first six dorsal ones descend forwards; the remaining ten pass perpendicularly backwards: hind part of the body, and tail fulvous: head flattish above, and marked by four transverse fulvous bands: dorsal fin edged with black at the back part: anal fin striped with black and pale yellow: tail marked with a fulvous crescent directed backwards, and tipped with white. Native of the Arabian seas: described by Forskal.

ARUAN CHÆTODON.

Chætodon Aruanus. *C. argenteus, corpore ovato-oblongo, fasciis tribus nigris.*

Silvery Chætodon, with ovate-oblong body crossed by three black bars.

Chætodon Aruanus. *Bloch. t. 198. f. 2.*

Body ovate-oblong: colour white, with a brownish cast on the upper parts; and marked by three

broad transverse black bars, the first of which crosses the head, the second the middle, and the third the hind-part: both these latter bars pass through the dorsal and anal fin as well as the body: the ventral fins are black: the scales on the body are rather large than small, and the gill-covers are destitute of spines. Native of the Indian and Arabian seas.

In the Museum Adolphi Friderici this species is named *arcuatus*, but as the author had before described a different species under that name, it is to be presumed, according to the opinion of Dr. Bloch, that *aruanus* is the true reading. Perhaps from the Aroo islands among the Moluccas.

CHINESE CHÆTODON.

Chætodon Chinensis. *C. albo-flavescens, corpore oblongo, fasciis pluribus fuscis, macula operculorum rotunda.*

Yellowish-white Chætodon, with oblong body crossed by several brown stripes, and a round spot on the gill-covers.

Chætodon Chinensis. *Bloch. t. 218. f. 1.*

Body oblong, and the whole habit more resembling that of the generality of fishes than of the major part of the Chætodons: colour dull yellowish white, brownish towards the back, and transversely barred from head to tail with several deep-brown stripes, some of which appear double: on the gill-cover a large round spot, and behind the eyes two small longitudinal streaks of the same colour: scales moderately large: dorsal and anal

fin resembling those of the generality of fishes.
Native of China.

SPOTTED CHÆTODON.

Chætodon Maculatus. *C. flavus, corpore oblongo-ovato longitudinaliter fusco maculato, fasciisque pallidioribus fuscis.*

Yellow Chætodon, with oblong-ovate body, spotted longitudinally with brown, and crossed with bands of paler brown.

Chætodon maculatus. *Bloch. t. 427. f. 2.*

SHAPE oblong-ovate, and in habit resembling some of the Carp tribe : colour yellow, marked by a longitudinal row of five or six large round brown spots, some of those on the fore and hind part of the body being partly obscured by a transverse bar or cloud of a similar colour passing over each : the middle spot is larger than the rest, and has no accompanying bar : the gill covers are smooth and scaly, the fins simple and of a bright yellow : the scales on the body are of moderate size : the tail rather large, and nearly even at the end, though perhaps, a very little inclining to a lunated shape. Native of India and South America, inhabiting fresh waters.

BIACULEATED CHÆTODON.

Chætodon Biaculeatus. *C. argenteo-cærulescens, corpore oblongo, fascus tribus distantibus, angustis, ferrugineis.*

Blueish-silvery Chætodon, with oblong body, crossed by three distant, narrow, ferruginous bands.

Chætodon biaculeatus. Bloch. t. 219. f. 2.

SHAPE oblong, with the back slightly sinking towards the middle: colour silvery, with a cast of blue, and crossed by three narrow, distant, ferruginous bands, the first across the gill-covers, the second across the middle, and the third across the end of the body: scales rather small: on the gill-covers, behind the eyes, a strong spine: dorsal and anal fins but slightly dilated at the hind part. Native of the Indian seas.

GOLDEN-SIDE CHÆTODON.

Chætodon Maculosus. *C. cinereus, antèrius maculis transversis cæruleis, posteriùs magna transversa aurea.*

Grey Chætodon, marked with transverse blue lines on the fore-part, and with a large transverse gold-coloured spot on the hind-part.

Chætodon maculosus. Forsk. F. Arab. p. 62. Lin. Gmel. p. 1267.

SHAPE ovate-oblong: ground-colour bright glossy grey: towards the beginning of the back several transverse blue streaks: on the gill-covers several blue spots: towards the hind part of the body a large transverse gold-coloured patch or bar, some-

what broken into the ground-colour: tail spotted with yellow: gill-covers furnished with a strong spine. Native of the Arabian seas: described by Forskal.

LUNULATED CHÆTODON.

Chætodon Lunulatus. C. fusco-virescens, lunula superciliari nigra, pectorali duplicata.

Brownish-green Chætodon, with black crescent over the eyes, and a double one near the breast.

Pomacentre Croissant. *Cepede. 4. p. 511.*

DESCRIBED by Cepede from the MSS. of Commerçon: back elevated: snout projecting, with the upper lip extensile, and shorter than the lower: general colour-dusky yellowish green: above each eye a black crescent: a double crescent of the same colour passing in an oblique direction from the pectoral fins to the middle of the back: a black spot on each side the tail, which is crossed by a black bar: a black streak at the extremity of the dorsal and anal fin, and a few longitudinal purple streaks on the abdomen. Native of the Indian seas. Shape of the tail not mentioned.

GOLDEN CHÆTODON.

Chætodon Aureus. *C. luteus, corpore ovato, pinna dorsali analique magnis, falcato-elongatis, apice viridibus.*

Gold-yellow *Chætodon*, with ovate body, and large, elongated, and falcated dorsal and anal fin green towards the extremities.

Chætodon aureus. *Bloch, t. 193. f. 1.*

CONSIDERABLY allied in point of habit to the Angel *Chætodon*, having a great breadth of body, with far-extended, falciform dorsal and anal fin, reaching considerably beyond the tail: colour of the body bright gold-yellow, shaded with orange on the upper parts: fins and tail yellow, at the base, gradually changing into green towards the extremities: at the bottom of the gill-covers a very strong spine: body and bases of the fins covered with middle-sized scales: lateral line curving in the direction of the back. Native of the American seas, growing to the length of a foot or more.

 ASFUR CHÆTODON.

Chætodon Asfur. *C. niger, fascia transversa lunari-cuneata flava.* *Forsk. F. Arab. p. 1267.*

Black *Chætodon*, with transverse, lunar-cuneated, yellow band.

Chætodon Asfur. *Lin. Gmel. p. 61.*

OBSERVED, and slightly described by Forskal: trunk oval; teeth numerous, filiform, and flexile: anterior opercula furnished with a strong spine:

general colour black, with a transverse sublunar yellow band in the middle of the body, the horns pointing backward: dorsal and anal fin falcated: tail fulvous, rounded, with a black margin: lateral line near the back: scales rhomboid, and finely toothed. Native of the Arabian seas.

VAR.

Of a blueish colour, with violet streaks and bands, disposed in an oblique direction. Observed by Forskal about the Arabian coasts, among corals, &c. the flesh is said to be bitter.

GEOMETRICAL CHÆTODON.

Chætodon Geometricus. *C. fuscus, albo variatus, lateribus circulis pluribus concentricis.*

Dusky Chætodon, with white variegations, disposed on the sides into several concentric circles.

L'Holocanthe géométrique. *Cepede. 4. p. 537*

Douwing formose. *Renard. 1. pl. 5. f. 34.*

SLIGHTLY described by Cepede, from the suspicious authority of Renard: it is named from the geometrical regularity of its pattern, which on a dusky ground exhibits on each side of the body seven or eight concentric white circles, of which the three or four interior are complete, the rest more or less broken: the head is streaked with various lines of the same colour. Native of the Indian seas.

CHÆTODON TRIFASCIATUS.

Chætodon Trifasciatus. *C. longitudinaliter striatus, fuscis tribus capitis nigris.* Park. *Lin. Trans.* vol. 3. p. 34.

Longitudinally striated Chætodon, with three black bands on the head.

LENGTH three inches: colour pale or brownish: body marked with sixteen longitudinal dusky stripes, and covered with rather large, ciliated scales: on the dorsal fin a black band, edged with yellow; another at the base of the anal fin, and a third through the middle of the tail; on the head three black bands. Observed by Mr. Mungo Park about the coast of Sumatra.

 GREAT-FINNED CHÆTODON.

Chætodon Pinnatus. *C. griseus, fascia frontali apiceque caudæ albis, pinna dorsali analique maximis falcatis.*

Grey Chætodon, with the front band and tip of the tail white, and extremely large falcated dorsal and anal fin.

Chætodon pinnatus. *Lin. Gmel.*

In habit greatly resembling the *Bat Chætodon*, but with the points of the fins more extended: colour grey, with a white band across the head, and the tail tipped with white: scales rather large: described by Linnæus in the *Museum Adolphi Friderici*. Native of the American and Indian seas.

CHILI CHÆTODON.

Chætodon Chilensis. C. aureus, fasciis quinque discoloribus, cauda integra, spinis dorsalibus undecim. Molin. Chil. p. 200. Lin. Gmel. p. 1264.

Golden Chætodon, with five coloured bands, even tail, and eleven dorsal spines.

NATIVE of the American seas. Observed by Molina about the coasts of Chili. Length twelve inches: shape oval: snout lengthened: scales small: head crossed by a black band; body by two brown ones, and tail by one cinereous and one black band: dorsal fin large and yellow: tail silvery.

PLEAT-NOSE CHÆTODON.

Chætodon Plectorhinchus. C. nigricans, maculis albis nigro-maculatis, naso plicato.

Blackish Chætodon, with white patches spotted with black, and pleated snout.

Le Plectorhinque Chetodonoide. Cope. 3. p. 135.

DESCRIBED by Cope from a specimen in the museum of the Prince of Orange, and, rather unnecessarily, ranked under a distinct genus, by the name of *Plectorhinchus*. Length, according to the plate, about five inches: habit that of the major part of the oblong-bodied Chætodons: colour blackish, or very dark, variegated with several large roundish, white or light-coloured patches, which



Chaetodon

PLEANT-NOSE (CHAETODON).

and other London Publishers by G. Francis, New York.

have generally one or more round black spots towards the middle: gill-covers without spine: mouth small, and concealed as it were among several membranaceous pleats into which the lips or edges are disposed: eyes rather large: scales small: tail even, or rounded: native region unknown.

With single dorsal fin, and forked or lunated tail

GLAUCOUS CHÆTODON.

Chætodon Glaucus. C. corpore rhombeo-ovato, supra cæruleo fasciis transversis nigris, subtus argenteo.

Blue Chætodon, with rhombic-ovate body; the upper parts crossed by black bands; the sides and abdomen silvery.

Chætodon glaucus. Lin. Gmel. Bloch. t. 210.

A highly elegant species: body rhombic-oblong: colour on the upper parts fine blue, on the lower silvery white with a tinge of rose-colour: on each side the upper part of the body are five or six narrow black bands, descending only half way down: dorsal and anal fin blue, very long, slightly falcated, and reaching as far as the tips of the tail, which is very widely forked: at the origin of the dorsal fin are five pretty strong spines, but none on the gill-covers: the scales are of moderate size. This species is a native of the American seas, and grows to a considerable size, often exceeding a foot or eighteen inches in length, and is in considerable esteem as a food.

RHOMBOID CHÆTODON.

Chætodon Rhomboides. *C. corpore rhombico-ovato, supra caeruleo-virescente, subtus flavescente, aculeis analibus duobus, dorsalibus quinque.*

Chætodon with rhombic-ovate body, greenish-blue above, yellowish beneath, two spines before the anal fin, and five before the dorsal.

Chætodon rhomboides. *Bloch. t. 209.*

OF a broader shape in proportion to its length than the former, and with a much larger and blunter head : colour on the upper parts grey or greenish blue, on the sides white, and on the abdomen pale yellow : the blue of the upper parts breaks into the white of the sides, in two or three acuminate bars or fasciæ reaching almost to the abdomen : the dorsal and anal fin are shaped as in the preceding species, but are much shorter, the points not reaching beyond the base of the tail, which is very widely forked : both fins and tail are blue, tinged with yellow at the base : pectoral and ventral fins yellow : scales moderately large : a few strong spines at the origin of the dorsal fin, but none on the gill-covers. Native of the American seas, where it grows to a very considerable size.

BORDERED CHÆTODON.

— *Chætodon Marginatus.* *C. flavus, fasciis transversis fuscis, pinnis fusco-marginatis.*

Yellow Chætodon, with transverse brown bands, and fins bordered with brown.

Chætodon marginatus. *Bloch. t. 207.*

A SPECIES of considerable beauty: body ovate: colour on the upper parts yellow, on the lower dull rose-colour: back of a dusky cast: body crossed by seven or eight equidistant deep-brown bars, the last crossing the base of the tail: all the fins are yellow, bordered with deep brown: dorsal fin broad and falcated at its extremity, but not reaching to the end of the tail: anal fin much smaller than the dorsal and situated more backward, being placed beneath the dilated part of the dorsal: tail very much forked, the points turning inwards in a falcated manner: scales moderately large: none at the base of the fins: no spines, either at the origin of the dorsal fin or at the gill-covers. Native of the American seas: grows to the length of a foot, and is numbered among the edible fishes.

TRICOLOR CHÆTODON.

Chætodon Tricolor. *C. antice flavus, postice niger, cauda marginibusque pinnarum rubris.*

Yellow Chætodon with the hind-part black, the tail and fins bordered with red.

Chætodon tricolor. *Bloch. t. 425.*

A HIGHLY superb species, and remarkable for the singular disposition of its colours: general shape resembling that of the *golden, angel, and ciliated* Chætodons, but longer in proportion: colour of the fore-part golden-yellow, of the hind-part, except the tail, jet-black, the two colours abruptly separated, in an oblique direction: edges of the gill-covers and of all the fins bright red: tail yellow, growing red towards the end, and terminating in pretty long tips, but not deeply divided in the middle: scales rather large: falcatures of the dorsal and anal fin not reaching to the tips of the tail: at the gill-covers a very large, long spine: mouth surrounded by a black zone. Native of the Indian and American seas: length about a foot.

SPECKLED CHÆTODON.

— *Chætodon Guttatus.* *C. griseo-albidus, corpore oblongo, maculis creberrimis rotundatis rufis.*

Whitish-grey *Chætodon*, with oblong body sprinkled with very numerous round rufous spots.

Chætodon guttatus. *Bloch. t. 196.*

SHAPE oblong, or resembling that of the generality of fishes: length about ten or twelve inches: colour grey, with a dusky tinge on the upper parts, and every where speckled, except on the head, pectoral, ventral, dorsal, and anal fins, with numerous small, round, rufous spots: scales small: none at the bases of the fins: gill-covers smooth: dorsal and anal fin of moderate breadth: tail slightly inclining to a forked or rather lunated shape, and speckled like the body: native of Java.

This species is supposed by Dr. Bloch and Cope to be the *Teuthis jacus* of Linnæus (the genus *Teuthis* being now very properly absorbed in those of *Chætodon* and *Acanthurus*). The description however given by Linnæus of his *Teuthis jacus* does not agree in the disposition of its colours with those of the present fish; being expressly said to be marked with longitudinal streaks: in this state it is also figured in the work of Gronovius. The Gronovian variety (if such it be) occurs in the British Museum, and perfectly corresponds with that author's figure, being marked by numerous abrupt or short longitudinal streaks.

PEACOCK CHÆTODON.

Chætodon Pavo. *C. fulvus, lateribus glaucis fusco transversim undulatis, capite caeruleo variegato.*

Fulvous Chætodon, with glaucous sides transversely undulated with brown, and head variegated with blue.

Chætodon Pavo. *Bloch. t. 198.*

IN shape somewhat resembling the preceding, but the body not tapering so much on the hind part: length about seven inches: colour dull orange, the sides glaucous, and marked by numerous, slightly undulated brown lines passing over the scales, which are rather large: head and fore parts sprinkled with numerous pale-blue spots: from each eye to the mouth three pale-blue streaks: dorsal and anal fin of moderate width and pointed at the ends: tail rather large and pretty deeply forked: native of the Indian seas.

 CURASSAO CHÆTODON.

Chætodon Curacao. *C. corpore rotundato-ovato subargenteo, fasciis tribus obscuris semidecurrentibus, squamis magnis.*

Subargenteous Chætodon with round-ovate body, marked by three obscure semi-decurrent bands, and covered with large scales.

Chætodon Curassao. *Bloch. t. 212.*

SHAPE roundish-ovate: scales very large: colour blueish silvery white, tinged with brown on the upper parts, which colour descends on the sides in

the form of three obscure bands or clouds: fins and tail, pale brown: dorsal and anal of moderate breadth, and scaled at the base: gill-covers smooth, or without spines: tail moderately forked or divided: native of the American seas, where it often grows to a considerable size, and is esteemed as an edible fish.

BENGAL CHÆTODON.

Chætodon Bengalensis. C. corpore subrotundato-ovato griseo, fasciis transversis fusco-flavescentibus.

Grey Chætodon, with roundish-ovate body marked by yellowish-brown transverse bands.

Chætodon Bengalensis. Bloch. t.

In shape resembling the preceding, but not quite so broad or deep: colour grey, with five or six yellowish brown bars across the body: scales rather large, like those of a carp: dorsal and anal fin of moderate breadth, scaled at the base, and pointed at the tips: gill-covers smooth: tail of moderate size, and rather deeply, but not sharply forked: native of the Indian seas: length about six inches.

ROCK CHÆTODON.

Chætodon Saxatilis. *C. griseus, corpore ovato, fasciis quatuor pinnis caudaque nigricantibus.*

Grey Chætodon, with ovate body, and with the fins, tail, and four transverse bands blackish.

Chætodon saxatilis. *Bloch, t. 206. f. 2.*

ALLIED to the preceding species, but of a more oblong shape: colour grey, darkest on the back: body crossed by four equidistant blackish bars: fins and tail blackish: gill-covers smooth: tail deeply forked, with subfalcated lobes: scales rather large, and resembling those of a carp. Native of the South-American and Indian seas, seldom growing to a greater length than six or eight inches, and not much esteemed as a food.

MAURITIAN CHÆTODON.

Chætodon Mauritii. *C. supra cœruleus, fasciis sex angustis transversis decurrentibus nigris, subtus argenteus.*

Chætodon with the upper parts blue crossed by six narrow decurrent bands, the lower parts silvery white.

Chætodon Mauritii. *Bloch. t. 213.*

SHAPE similar to that of a carp: scales of moderate size: body dusky blue above, silvery white beneath: across the upper parts six narrow blackish bands, reaching almost to the abdomen: fins and tail brown: gill-covers smooth. Native of the Brazilian seas, growing to the length of two feet, and esteemed as a food. Described by Dr. Bloch

from the drawings and manuscripts of the celebrated Prince *John Maurice of Nassau-Sigen*, under whose conduct the Dutch, in the years 1637 and 1638, became possessed of the richest part of Brasil, where he afterwards resided some years in the character of Governor. During his residence he amused his hours of leisure in making drawings and composing descriptions of the most remarkable animals of the country: these drawings and manuscripts are preserved in the royal library at Berlin.

PUNCTATED CHÆTODON.

Chætodon Punctatus. C. argenteus, fusco-punctatus, spinis pinnae dorsalis octo, pinnis pectoralibus falcatis.

Silvery Chætodon, speckled with dusky points, with eight rays in the dorsal fin, and falcated pectoral fins.

Chætodon punctatus. C. spinis pinnae dorsalis octo, pinnis pectoralibus falcatis. Lin. Gmel. p. 1243.

SHAPE resembling that of a Carp: colour white or silvery, scattered over with dusky points: eyes red: lateral line curving upwards: the three first rays of the anal fin distant: in the dorsal fin are eight rays, in the caudal seventeen: native of the Indian seas?

ROUND CHÆTODON.

Chætodon Rotundus. *C. rotundatus cinereus, fasciis quinque pallidis.* Lin. Syst. Nat. p. 466.

Orbicular grey Chætodon, with five pale bands.

THIS seems a species very little known: from the name applied to it by Linnæus it should seem to have the body more orbicular in its outline than any other species.

SINGLE-SPOTTED CHÆTODON.

Chætodon Unimaculatus. *C. corpore rotundato-ovato griseo, macula utrinque prope dorsum ovata nigra, capite fascia transversa nigra.*

Grey Chætodon, with round-ovate body, marked on each side near the back with an oval black spot, and with a transverse black bar across the head.

Chætodon unimaculatus. Bloch. t. 201.

Body round-ovate: colour grey, with a moderately large oval black spot on each side, near the back; across the head a black bar, passing through the eye: dorsal and anal fin rounded and edged with black: scales rather large: gill-covers smooth. Native of India.

ORBICULAR CHÆTODON.

Chætodon Orbicularis. *C. corpore rotundato griseo punctis fuscis aspero, abdomine albido.*

Chætodon with orbicular body freckled with dusky specks, and whitish abdomen.

Chætodon orbicularis. *Forsk. F Arab. p. 59. Lin. Gmel. p. 1265.*

LENGTH about a foot: colour dusky grey; variegated with dusky specks, especially towards the back and tail: abdomen whitish, tinged behind with yellow: scales of moderate size: tail slightly divided. Native of the Arabian seas: observed by Forskal, who adds that it has something of the habit of a *Pleuronectes*.

SILVER CHÆTODON.

Chætodon Argenteus. *C. cauda bifida, spinis pinnae dorsalis octo, pinnarum ventralium loco spinis duabus.* *Lin. Syst. Nat. Gmel. p. 1242. Ann. Acad. 4. p. 240.*

Silvery *Chætodon*, with bifid tail, eight spines in the dorsal fin, and two spines in the place of ventral fins.

THIS, which appears to be an anomalous species, is said by Linnæus to differ from the rest of the genus in the defect of the pectoral fins, in place of which are two small spines: the first spines of the dorsal fin are so small as to be scarce perceptible. Native of the Indian seas.

LONG-SNOURED CHÆTODON.

Chætodon Longirostris. *C. corpore rhombo-oblongo subflavo, rostro elongato, pinna anali macula nigra.*

Chætodon with rhombic-oblong yellowish body, lengthened snout, and black spot on the anal fin.

Chætodon longirostris. *Lin. Gmel. Brouss. dec. 1.*

SHAPE oblong-rhomboid: head sloping, and produced into a cylindric snout, as in the *rostratus*, to which this species is allied: head brown above, and of a silvery flesh-colour beneath: body yellowish or citron-colour, covered with unequal, obliquely imbricated scales: lateral line strait: dorsal and anal fins citron-colour with a black streak on the hind-part, edged with white; and towards the tip of the anal fin a black ocellated spot: pectoral fins and tail glaucous, the former tinged with yellow at the base. Native of the Pacific ocean.

BODDAERTIAN CHÆTODON.

Chætodon Boddaerti. *C. fasciis fuscis et caeruleo-viridibus, spinis pinnarum ventralium duabus.* *Lin. Syst. Nat. Gmel p. 1243.*

Chætodon variegated with brown and blueish bands, and with two spines at the ventral fins.

A SPECIES hitherto but very little known: native of the Indian? seas.

SARGOID CHÆTODON.

Chætodon Sargoides. *C. luteus, cupite fasciisque sex transversis violaceis.*

Gold-yellow Chætodon, with the head and six transverse bands violet-colour.

Le Chetodon Sargoide. *Cepede. 4. p. 473.*

SLIGHTLY described by Cepede from the drawings and manuscripts of Plumier. Allied, as it should seem by the title, to the Sparus Sargus in habit: general colour gold yellow; head and six transverse bands, together with the edges of the dorsal, anal, and caudal fin of a fine violet-colour: in the dorsal fin are thirteen spiny rays, and one in each pectoral fin: a depression before the eyes; the opening of the mouth small, and the upper lip thick: gill-covers rounded. Native of the American seas.

CANESCENT CHÆTODON.

Chætodon Canescens. *C. cauda bifida, spinis dorsalibus duabus, radio tertio longissima, ore bidentato. Lin. Gmel. p. 1240.*

Greyish Chætodon, with bifid tail, two dorsal spines, the third ray very long, and bidentated snout.

Chætodon canescens. *Seb. mus. 3. t. 25. f. 7.*

SHAPE resembling that of the *C. cornutus* and *bifasciatus*: colour greyish: scales very small: on each side the mouth a sort of spine or process: third ray of the dorsal fin extremely long, as in the two species just mentioned. Native of the American and Indian seas.

SCALELESS CHÆTODON.

Chætodon Alepidotus. *C. cauda bifida, spinis dorsalibus tribus, pinnis ventralibus nullis.* Lin. Syst. Nat. Gmel. p. 1240.

Chætodon with bifid tail, three dorsal spines, and no ventral fins.

SHAPE rhomboid : colour blueish above, whitish beneath : scales either none or so small as to be difficultly seen : teeth in a single row : lateral line parallel with the back : tail bifid : no ventral fins : dorsal and anal falcated : native of the American seas : observed by Dr. Garden.

 SORDID CHÆTODON.

Chætodon Sordidus. *C. ex fusco cinereus, ovatus, fasciis transversis obsoletis quatuor.* Lin. Gmel. p. 1267. Forsk. Arab. p. 62.

Dusky-grey Chætodon with four obscure transverse bands.

LENGTH about a span : body ovate : colour dull grey, with four paler transverse bands : scales large and membranaceous : native of the Arabian seas, frequenting the shores among corals, &c. observed by Forskal, who informs us that it is much esteemed as an edible fish.

LAMARK'S CHÆTODON.

Chætodon Lamarkii. *C. luteus, striis tribus longitudinalibus fuscis.*

Gold-yellow Chætodon, with three longitudinal dusky stripes.

Chætodon Lamarck. *Cepede. 4. p. 526.*

GENERAL colour gold-yellow, with three longitudinal dusky stripes on each side the body: lower jaw longer than the upper: scales rounded, striated, and denticulated: fifteen spiny rays and sixteen soft ones in the dorsal fin: three spiny and twenty soft ones in the anal: gill-covers armed with a very long spine. Described by Cepede from a specimen in the Prince of Orange's museum.

With two dorsal fins.

PLUMIER'S CHÆTODON.

Chætodon Plumieri. *C. subflavus, subtus subargenteus, corpore fastiis sex virentibus, pinna dorsali analique falcatis.*

Yellowish Chætodon, subargenteous beneath, with six greenish bands across the body, and dorsal and anal fin falcated.

Chætodon Plumieri. *Block. t. 211.*

SHAPE rhombic-ovate, in habit resembling the *angel, rhomboid*, and *ciliated* Chætodons. Colour dull yellow above, silvery beneath: across the body six greenish stripes: first spines of the back connected by a membrane, so as to form a small separate fin: second dorsal and caudal, large, long, and falcated, so as to extend beyond the tail, which is

of moderate size, and slightly lunated : gill-covers without spine. Native of the American seas, where it arrives at a considerable size.

CONSTRICTED CHÆTODON.

Chætodon Constrictus. *C. griseo-flavescens, nigro transversim fasciatus, corpore medio constricto.*

Yellowish-grey Chætodon, with transverse black bands and body constricted in the middle.

Chætodon constrictus. *New Holl. zool. p. 27. t. 6.*

LENGTH about eight inches : shape of the body inclining to an oblong square, and remarkably contracting in diameter about the middle, so as to appear constricted in that part : scales of moderate size : colour yellowish grey, tinged on the back and part of the fins with blue : across the body eight black bands, that in the middle narrowest : on the back two very distinct fins, the rays of the first being all spiny : tail inclining, though very slightly, to a lunated form. Native of the Indian seas, and observed about the coasts of New Holland.

ARMED CHÆTODON.

Chætodon Armatus. *C. argenteus, fasciis septem transversis nigris, capite elongato, pinnis dorsalibus duabus.*

Silvery Chætodon, with seven transverse black bands, lengthened head, and two dorsal fins.

Chætodon albescens, corpore fasciis septem nigris, spinis pinnæ dorsalis sex, tertia longissima. *White journ. New South Wales, p. 254. pl. 39. Naturalist's Miscellany, vol. 2. pl. 57.*

LENGTH of the specimen described about four inches: colour silvery white, darker and with a blueish tinge on the back: head of a somewhat lengthened form: across the body seven black bands: on the back two distinct dorsal fins, of which the first consists of very strong spiny rays, the third exceeding the rest in length; tail very slightly inclining to a lunated form at the extremity. Native of the Indian seas, and observed about the coast of New Holland.

ACANTHURUS. ACANTHURUS.

Generic Character.

<i>Dentes</i> parvi, plerisque lobati.	<i>Teeth</i> small, in most species lobated.
<i>Cauda</i> utrinque aculeata.	<i>Tail</i> aculeated on each side.
<i>Habitus</i> generis præcedentis.	<i>Habit</i> or general appearance as in the preceding genus.

THE genus *Acanthurus* consists of such species of the Linnæan genus *Chatodon* as, in contradiction to the principal character of that genus, have, in general, moderately broad and strong teeth, rather than slender and setaceous ones: they are also furnished on each side the tail with a strong spine.

UNICORN ACANTHURUS.

Acanthurus Unicornis. *A. griseo-fuscus*, cornu frontali antrosum porrecto, cauda utrinque biaculeata.

Grey-brown *Acanthurus*, with a frontal horn stretching forwards over the snout, and two spines on each side the tail.

Monoceros minor. *Will. ichth. p.* 216. *t. O.* 4.

Le Nason Licornet. *Cepedc.* 3. *p.* 103.

THIS and the following are by far the largest of the genus, growing to the length of three feet or



THALASSOMA ACANTHURUS

Surge wrasse. Thalassoma acanthurus. J. S. G. 1854.

more: the present species is of the same general shape with the major part of the *Chaetodons*, being of a moderately broad or deep form, tapering pretty suddenly at each extremity: from the front proceeds a strong, conical, horn-shaped process, extending forwards nearly as far as the tip of the snout, and terminating somewhat obtusely: the mouth is rather small, and the teeth moderately strong: the skin is somewhat rough, of a greyish brown colour, and destitute of apparent scales: the dorsal fin, which is rather shallow than broad, commences almost immediately behind the head, and is continued very nearly to the tail: at its origin are two or three very strong spines: the anal fin commences immediately beyond the breast, and like the dorsal, is continued nearly to the tail, which is moderately broad and of a lunated shape: on each side the tail are situated two strong, but short spines, rising from a very broad base: these, as well as the horn in front, are much less conspicuous in the young than in the more advanced specimens.

It is a native of the Indian and Arabian seas, in the latter of which it is generally seen in large shoals of two or four hundred each, swimming with great strength, and feeding principally on the different kinds of sea-weed. Forskal, in his *Fauna Arabica*, tells us he was assured by a fisherman that he saw an eagle (probably an Osprey) dart among a shoal of these fishes, which swim pretty close together, and happening to seize two at once in his claws, was, in consequence of their sudden

and violent separation in different directions, torn asunder in the conflict and destroyed.

This fish appears to have been first described by Grew in his Museum of the Royal Society, under the name of the *lesser unicorn fish*; he observes that the head, when viewed in profile, bears some resemblance to that of a baboon: the caudal spines seem to have been wanting in this specimen; otherwise it is scarcely to be imagined that so accurate an observer as Grew would have omitted to particularize them. Willughby repeats Grew's description, and his figure is evidently engraved from the same specimen. It is singular that so remarkable a fish should have been entirely overlooked by Linnæus, even in the twelfth edition of the *Systema Naturæ*. In the British and Leverian museums it occurs in fine preservation, and in the former are specimens in a young state, shewing the comparatively inconspicuous appearance both of the frontal process and caudal spines at that period of its growth.


ROUND-FRONTED ACANTHURUS.

Acanthurus Nasus. *A. griseus, nigro irroratus, gibbere frontali rotundato, cauda utrinque biaculeata.*

Grey Acanthurus, freckled with black, with a rounded frontal tubercle and two spines on each side the tail.

Le Nason Loupe. *Cope*. 3. p. 111.

In size this species equals, or even exceeds, the preceding: the body is of a more oblong shape,



Round snout

ROUND-FRONTED ACANTHURUS.

From the London Illustrated by a Quarterly Press Series.

and is of a greyish colour, speckled on the upper parts, dorsal fin, and tail, with black: the head is rather large, and is formed in front into a rounded protuberance immediately above the mouth which is rather small and furnished in each jaw with a row of numerous small teeth: the dorsal fin commences at a small distance from the head, and is continued nearly to the tail: at its commencement is a strong spine, but the rest of the rays are soft: the anal fin is placed in an opposite situation beneath, and extends, like the former, almost to the tail, which is moderately broad, and lunated: on each side the base, at a small distance from the commencement of the finny part, are situated two very strong, slightly curved, sharp-pointed spines, in a reversed direction, or with the points towards the head, as in most others of the present genus. This species is a native of the Indian seas, and according to Cope, was first described by Commerson.

 BLUE ACANTHURUS.

Acanthurus Teuthis. *A. cæruleus, corpore medio pallidiorè, cauda utrinque aculeata.*

Blue Acanthurus, with the middle of the body paler, and a spine on each side the tail.

Chætodon cærulescens, &c. *Seb.* 3. t. 33. f. 3.


Teuthis Hepatus. *Lin. Syst. Nat.* p. 507.

LINNÆUS, in the twelfth edition of the *Systema Naturæ*, thought it necessary to arrange this fish under a distinct genus among the *pisces abdominales*, under the name of *Teuthis*. It is however a genuine

Acanthurus, and the genus Teuthis is, in consequence, annulled. The blue Acanthurus is a native of the Indian and American seas, and is said to arrive at the length of ten or twelve inches, or even more, though the specimens usually seen are considerably smaller : in shape it resembles the major part of the Chaetodons, the head sloping in front from the origin of the dorsal fin : the colour, in the recent fish, is a deep or blackish blue, which in specimens preserved in spirits of wine, usually fades into a ferruginous brown : on each side the body is a very large, oblong-ovate whitish patch or spot, surrounded by a border of a deeper cast : the skin is roughened by very small scales : the tail is slightly lunated, dusky on the upper and lower part, and marked towards the base by a whitish ovate spot : the teeth are crenated, and on each side the base of the tail, is a very strong spine, which, as in most others of the present genus, is erigible at the pleasure of the animal, in an horizontal direction, from the channel in which it lies.

The fish figured by Catesby in his *Natural History of Carolina*, under the name of the *Tang**, is generally considered as representing the present species : yet, if Catesby's description be accurate, the Tang is destitute of teeth ; and, from the figure accompanying the description, appears to want the ventral fins, is covered with middle-sized scales, and is entirely of a deep blue colour without any variegation. As Catesby however was not remark-

* *Turdus rhomboidalis.* *Catesb. Car. 2. t. 10.*



LANCET ACANTHURUS.

able for minute accuracy either in his descriptions or figures, it is probable that it may have been intended for the present species. Catesby informs us that it grows to the length of a foot and half, and is in request as an edible fish.

LANCET ACANTHURUS.

Acanthurus Chirurgus. *A. luteus, corpore postice striis transversis fuscis, cauda utrinque aculeata.*

Orange-yellow *Acanthurus*, with the body crossed on the hind-part by transverse brown stripes, and a spine on each side the tail.

Chætodon Chirurgus. *Bloch. t. 208.*

LENGTH about nine or ten inches : shape ovate : colour orange-yellow : scales rather small : on each side the hind part of the body five or six narrow, transverse, dusky stripes : gill-covers smooth or unarmed : on each side the base of the tail a very strong, curved spine, erigible at pleasure from the furrow in which it lies : tail lunated : dorsal fin pale violet, without scales at the base, and marked by several oblique, narrow, dusky streaks : pectoral and ventral fins tinged with crimson towards the base : native of the American seas, where it is generally known by the name of the Surgeon or Lancet-Fish, a title however which seems not entirely confined to the present species.

BLACKISH ACANTHURUS.

Acanthurus Nigricans. *A. nigricans, subtus subargenteus, cauda utrinque aculeata.*

Blackish Acanthurus, subargenteous beneath, with a spine on each side the tail.

Chætodon nigricans. *Lin. Bloch. t. 203.*

Of nearly the same size with the preceding, but with the front sloping more suddenly downwards from the origin of the dorsal fin: colour dusky, or blackish brown, with the under parts inclining to a subargenteous cast: scales rather small: dorsal and anal fin scaled at the base: gill-covers unarmed: on each side the tail a strong and sharp spine, as in the former species: tail nearly entire, or but slightly lunated: teeth broad and crenulated or lobed on the edges: native of the Indian, American, and Arabian seas, and said to grow to the length of two feet, and to be esteemed for the table.

MILITARY ACANTHURUS.

Acanthurus Militaris. *A. corpore rhombico-ovato fusco, cauda utrinque aculeata.*

Brown Acanthurus, with rhombic-ovate body, and strong spine on each side the tail.

LENGTH about five inches: shape rhombic-ovate: head sloping suddenly downwards: colour brown: scales extremely small: teeth denticulated or cre-

nated: a strong spine at the base of the pectoral fins. Tail slightly lunated: spine on each side very strong and sharp: native of the Indian and American seas: in the British and Leverian Museums.

TRIRADIATED ACANTHURUS.

Acanthurus Triostegus. *A. fusco-virescens, fasciis quatuor transversis fuscis, cauda utrinque aculeata.*

Greenish-brown Acanthurus, with four transverse dusky bands, and a spine on each side the tail.

Chaetodon triostegus. *Lin. Chaetodon, &c. Seb. 3. t. 25. f. 4.*

LENGTH about four inches: shape rhombic-ovate: head suddenly sloping: general colour pale greenish brown, marked with four narrow transverse brown stripes or bands: scales very small: gill-covers unarmed: gill-membrane three-rayed: in the throat three small rough bones: teeth crenated: tail lunated: spine on each side very strong. Native of the Indian seas.

GUARDED ACANTHURUS.

Acanthurus Harpurus. *A. corpore rhombeo-ovato fusco, squamis minutissimis, cauda utrinque biaculeata.*

Rhombic-ovate, brown Acanthurus, with extremely minute scales, and two spines on each side the tail.

SHAPE rhombic-ovate: length about six inches: colour dusky brown, without any variegation:

scales most extremely small, so as to be scarce visible : gill-covers unarmed : teeth crenated : tail lunated, and furnished on each side the base with two very strong spines. Native of the Indian seas. In the British Museum.

VIOLET-STRIPED ACANTHURUS.

Acanthurus Sohal. *A. fuscus striis longitudinalibus violaceis, cauda utrinque aculeis duobus in fossula rubra.*

Dusky Acanthurus, with longitudinal violet streaks, and two spines on each side the tail imbedded in a red depression.

Chætodon Sohal. *Forsk. Arab. p. 63.*

LENGTH about three spans : shape ovate : colour dusky, with longitudinal violet streaks : abdomen whitish : fins violet-coloured : pectoral fins marked by a yellow spot : teeth crenated : tail lunated : on each side the base a long, red carina, imbedded in a depression of the same colour, at each * extremity of which is a strong spine. Native of the Red sea : not esteemed as an edible fish.

* Spina pone et ante hanc carinam. *Forsk. arab. p. 64.*

DUSKY ACANTHURUS.

Acanthurus Nigro-fuscus. A. nigro-fuscus, corpore ovato, cauda utrinque aculeata.

Dusky Acanthurus, with ovate body, and spine on each side the tail.

Acanthurus nigro-fuscus. Forsk. Arab. p. 64.

Much allied, according to Forskal, to the *A. Sohal*, but differs in being entirely of a blackish-brown colour, as well as in having the lobes of the tail much shorter, and a single spine on each side.

A variety of the above, entirely of a black colour, is mentioned by Forskal, under the title of *Gahm*. Both are natives of the Red sea.

ACHILLEAN ACANTHURUS.

Acanthurus Achilles. A. corpore ovato fusco, posterius macula ovata nuda rubra, medio aculeata.

Brown, ovate Acanthurus, with a bare ovate red spot on the hind-part, aculeated in the middle.

LENGTH about five or six inches: shape ovate: colour brown: scales small: gill-covers unarmed: teeth broad and crenated: on each side the hind-part of the body or base of the tail a large, ovate, bare, red patch or spot, in the middle of which is a strong spine imbedded in a furrow: tail lunated and tipped by three concentric crescents, the first of which is white, the next black, and the third ferruginous, edged with ash-colour. An elegant

specimen of this fish occurs in the Leverian Museum. In some particulars it seems allied to the *A. Sohal*, but, if Forskal's description of that fish be accurate, must be a very distinct species.

LINEATED ACANTHURUS.

Acanthurus Lineatus. *A. ovatus fuscus, lineis numerosis longitudinalibus albis, cauda utrinque aculeata.*

Ovate brown *Acanthurus*, with numerous longitudinal white stripes, and spine on each side the tail.

Chaetodon lineatus. *Lin.*


LENGTH about six inches: shape ovate: colour brown, marked by pretty numerous, longitudinal, narrow, white stripes, running from head to tail: each stripe edged with a darker colour than that of the ground: scales small: gill-covers unarmed: teeth crenated: tail lunated: tail-spines very strong, of moderate length, and sharp-pointed. Native of the Indian and American seas: in the Leverian Museum.

UMBRE ACANTHURUS.

Acanthurus Umbratus. *A. corpore ovato fusco, squamis minutissimis, cauda utrinque aculeata.*

Brown ovate *Acanthurus*, with extremely minute scales, and a spine on each side the tail.

LENGTH about four inches: shape ovate: colour brown: scales extremely minute: gill-covers un-



Scorpaenopsis diabolus (Lacépède) (Pisces, Scorpaenidae)

armed: tail slightly lunated: tail spines very sharp and strong. Native of the Indian seas. In the British Museum.

SPECKLED ACANTHURUS.

Acanthurus Meleagris. *A. fusco-nigricans, maculis creberrimis rotundis albis, cauda utrinque aculeata.*

Blackish-brown *Acanthurus*, thickly marked with round white spots, and spine on each side the tail.

LENGTH about four inches: shape roundish-ovate: colour deep brown, thickly marked on all parts with small, round, white spots: teeth crenated: gill-covers unarmed: scales extremely small: tail lunated: spines moderately strong. Native of the Indian and American seas. In the British Museum.

BROAD-FINNED ACANTHURUS.

Acanthurus Velifer. *A. corpore rotundato-ovato albido, fasciis plurimis transversis fuscis, cauda utrinque aculeata.*

Whitish *Acanthurus*, with roundish-ovate body, marked by numerous brown transverse bands, and spine on each side the tail.

Acanthurus Velifer. *Block. t. 427. f. 1.*

LENGTH about four inches and a half: body roundish-ovate: ground-colour whitish, with pretty numerous narrow transverse stripes of yellowish-brown: across the head a broader black stripe, passing through the eyes: teeth crenated, and

pointed at the tips: dorsal and anal fin very broad, and thickly marked with round blue spots: tail slightly lunated, and furnished on each side the base with a spine. Native of the American seas.





AMERICAN KNIFE-FISH.

1848. *Illustration Published by G. & J. S. Brown.*

EQUES. KNIGHT-FISH.

Generic Character.

<i>Dentium</i> ordines plurimi.	<i>Teeth</i> in several rows.
<i>Corpus</i> fasciatum.	<i>Body</i> banded.

AMERICAN KNIGHT-FISH.

Eques corpore oblongo flavescente, fasciis tribus nigris, oculari, pectorali, longitudinalique.

Yellowish oblong-bodied Knight-Fish, with three black bands, the first across the eyes, the second across the thorax, and the third along the body.

Chætodon lanceolatus. Lin. Eques Americanus. Bloch. t. 347.

THIS fish, which by Linnæus was associated with the Chætodons, has been formed into a distinct genus by Dr. Bloch on account of its teeth, which, instead of being arranged in a single row, as in those animals, are disposed into several concentric ranges. It is a native of the American seas, and is found of the length of near twelve inches.

TRICHOPUS. TRICHIOPUS.

Generic Character.

<i>Corpus compressum.</i>	<i>Body compressed.</i>
<i>Pinnæ ventrales filamento longissimo.</i>	<i>Ventral fins with an extremely long filament.</i>

THE fishes belonging to this new genus are described by Cope from the manuscripts and drawings of Commerson. Cope however, very unnecessarily, divides them into two distinct genera, to one of which he assigns the title of *Osphronemus* (*Osphroneme*), and to the other of *Tricopodus* (*Trichopode*). These fishes are natives of the Indian seas or rivers, and are distinguished, except in one species, by the remarkable length of one ray of the ventral fins, which extends the whole length of the body, or even beyond it.


GORAMY TRICHOPUS.

Trichopus Goramy. T. subargenteo-rufescens, radio secundo pinnarum ventralium longissimo.


Rufescent *Trichopus*, with a silvery cast on the sides, and the second ray of the ventral fins extremely long.

Osphronemus Goramy. Cope. 3. p. 116.

NATIVE of the fresh waters of China: length two metres: depth very great in proportion: body much compressed, and covered with large scales,



GORAMY TRICHOPTUS .



SATYR TRICHOPTUS .

the hind part of the back, and the lower part of the abdomen sharpening into a carina: the bases of the dorsal and anal fin are also scaly, as in the genus *Chaetodon*: mouth somewhat extensile, each jaw being furnished with a double row of teeth, those of the outer row short and slightly curved, those of the inner row smaller and more closely set: in the palate is a callous protuberance: the tongue is situated deep in the fauces, and in the inside of the mouth, above the gills, is a kind of ethmoid bone, the use of which does not appear to be well ascertained: the dorsal fin is of moderate width, and reaches from the back of the head almost to the tail, near which it is very much rounded: the vent is situated towards the fore-part of the body, and the anal fin reaches from it in the manner of the dorsal, but is much wider: the pectoral fins are of moderate size: the ventral consist each of about six rays, of which the first is strong and sharp, and the second terminated by an extremely long filament; reaching as far as the end of the tail which is of a rounded shape: the lateral line is strait, and seated somewhat nearer the back than the abdomen: the colour of this species is reddish brown, somewhat paler on the under parts: the sides are tinged with a silvery hue, and the scales are bordered with brown: it is greatly esteemed in China as an article of food, and is known by the name of *Goramy*: it is a prolific species, and is easily introduced into new situations; having been imported into Java and many other of the eastern islands.

ARABIAN TRICHOPUS.

Trichopus Arabicus. *T. virescens violaceo caruleoque striatus, pinnarum ventralium radio secundo longissimo.*

Greenish Trichopus, with violet and blue stripes, and second ray of the ventral fins extremely long.

Osphroneme Gal. *Cepede.* 3. p. 122.

Labrus Gallus. *Lin. Syst. Nat. Gmel.*

Scarus Gallus. *Forsk. arab.*

OBSERVED by Forskal in the Arabian seas, and referred by him to the genus *Scarus*: considered by the Arabians as a poisonous fish, owing, probably, to its feeding on some noxious kinds of Mollusca, and sea-insects: size not particularized; colour dull green, with violet stripes: on the abdomen two blue stripes, with a green one in the middle: scales lax, membranaceous at the edges, striated, and marked by a transverse purple stripe: teeth in a single row, those in front larger than the rest, and standing separate from each other: lateral line slightly branched: pectoral fins ovate violet-blue: ventral blue, with the second ray extremely long: dorsal and anal greenish, edged with blue: tail yellow in the middle, violet-coloured on the sides, and bordered with blue.

SATYR TRICHOPUS.

Trichopus Satyrus. *T. fronte descendente, mento prominente, pinnis ventralibus uniradiatis longissimis.*

Trichopus with sinking forehead, projecting chin, and extremely long, single-rayed ventral fins.

Trichopode Mentonier. *Cepede*. 3. p. 126.

DESCRIBED by Cepede from the manuscripts of Commerson: size not particularized: body very deep, much compressed, and covered with moderately large scales: dorsal and anal fin widish, like those of the first species; pectoral short, large, and rounded: ventral consisting each of a single filament, thickish at the base, and gradually lessening in its progress, reaching beyond the tail: head of an extremely singular aspect, being so formed as to bear a rude general resemblance to the human face, or rather to that of a monkey, the chin projecting, the forehead somewhat sinking in, and the gill-covers on each side representing the cheeks: colour and size not mentioned: native of the Indian seas. This fish, like the first species, is known in some countries by the name of *Goramy* or *Gouramy*.

PALLASIAN TRICHOPUS.

Trichopus Pallasii. *T. fuscus albido-undulatus, corpore candoque macula utrinque nigra, pinnis ventralibus uniradiatis longis.*

Brown *Trichopus*, with pale undulations, a black spot on each side the body and tail, and long single-rayed ventral fins.

Labrus pinnis ventralibus uniradiatis. *Lin. Syst. Nat. Gmel.* p. 1286. *Pall. spic. zool.* 6. p. 45.

Sparus, &c. Kolreuter nov. comm. petrop. 9. p. 452. t. 10. f. 1.

DESCRIBED by Dr. Pallas as a species of *Labrus*, and by Kolreuter, in the Petersburg Transactions, as a *Sparus*: length about four inches: head rather small, and covered with scales: mouth small, ascending; jaws extensile, and rough on the edges: colour brownish, with paler undulations, each side of the body and tail being marked with a black ocellate spot: lateral line somewhat broken or interrupted towards the tail: pectoral fins thin and pale; the rest brown: anal and caudal roundish-bilobate, and spotted with white: ventral consisting each of a long single ray: native of the Indian seas.

MONODACTYLE TRICHOPUS.

Trichopus Monodactylus. *T. argenteus, dorso sulfusco, pinnis ventralibus uniradiatis, rigidis, brevibus.*

Silvery *Trichopus*, with brownish back, and short, single-rayed, rigid ventral fins.

Monodactyle Falciforme. *Cepede.* 3. p. 132.

DESCRIBED by Cepede in a distinct genus, under the name of *Monodactylus*. Differs from the rest

in having two short spines only, by way of ventral fins: is a small fish; measuring only about twenty-six centimetres in length: body much compressed, and covered with small, rounded, smooth scales; as are also the bases of the dorsal and anal fin: colour silvery, tinged with brown on the upper parts: eyes large: mouth semicircular: teeth short, sharp, and close-set: dorsal and anal fin of nearly equal size and of a somewhat falciform shape, extending nearly to the tail, which is crescent-shaped: lateral line curved in the direction of the back, and placed pretty near to it: vent almost immediately behind the pectoral fins, which are of a pointed shape: gill-covers composed of two laminæ, of which the hindmost is irregularly scalloped on the edges: native of the Indian seas: described from the manuscripts of Commerson.

SCARUS. SCARUS.

Generic Character.

<i>Maxillæ</i> osseæ, medio di- visæ, margine crenatæ, dentibus connato-glomer- atis.	<i>Jaws</i> bony, divided in the middle, crenated on the edge; the teeth connate and conglomerate.
--	---


CRETAN SCARUS.

Scarus Cretensis. *S. viridis, subtus flavescent, squamis per-
magnis, linea laterali ramosa, cauda sublunata.*

Green Scarus, yellowish beneath, with very large scales, rami-
fied lateral line, and sublunated tail.

Scarus Cretensis. *Bloch. t. 220.*

GENERAL length about twelve inches: body broad, sloping, scales extremely large, lateral line ramified on every scale over which it passes. Na-
tive of the Mediterranean, and particularly about
the coasts of Crete, but is also found in the Indian
seas.



Surge wrasse

GREEN SCARTS.

Wash. Ship, London Published by G. Benceley, Fleet Street

GREEN SCARUS.

Scarus Viridis. *S. viridi-flavescens, squamis magnis viridi marginatis, linea laterali versus caudam interrupta.*

Yellowish-green Scarus, with large scales edged with green, and lateral line interrupted towards the tail.

Scarus viridis. *Bloch. t. 222.*

LENGTH ten or twelve inches : head strait : scales large, and each bordered with a deeper green than the ground-colour : tail slightly subnated, Native of the Japanese seas.

RIVULATED SCARUS.

Scarus Rivulatus. *S. caeruleus nigro-maculatus, rivulis longitudinalibus flavis.*

Blueish Scarus, spotted with black, and marked by longitudinal yellow undulations.

Scarus rivulatus. *Forsk. arab. p. 25.*

NATIVE of the Red sea : observed by Forskal : said to arrive at a great size ; scales very small : dorsal and anal fin occasionally recumbent in a channel : tail forked : supposed to feed principally on the different kinds of fuci, and considered as an edible fish ; but said to be sometimes productive of disagreeable symptoms from the wounds inflicted by the sharp rays of its dorsal fin.

STELLATED SCARUS.

Scarus Stellatus. *S. ocalis nigricans, annulis subhexagonis pallidis variatus.*

Oval blackish Scarus, variegated with subhexagonal pale rings.

Scarus stellatus. *Forsk. arab. p. 26.*

LENGTH about half a foot: colour dusky or blackish, variegated on all parts by small, contiguous, pale, subhexagonal rings: native of the Arabian seas: observed by Forskal: an edible species.


RED SCARUS.

Scarus Croicensis. *S. roscus, abdomine argenteo.*

Rose-red Scarus, with silvery abdomen.

Scarus Croicensis. *Blach. t. 221.*

LENGTH about ten inches: shape like that of a Carp: colour strong rose-red, with silvery abdomen; and in some specimens two longitudinal silvery stripes on the body: scales large: fins pale: tail slightly lunated. Native of the Indian seas.



RED SCORPION.

PARROT SCARUS.

Scarus Psittacus. *S. virescens, lineis flavicantibus, pinnarum margine, vitta abdominali, caracteribusque capitis caeruleis.*
Forsk. arab. p. 29.

Greenish Scarus, marked with yellowish lines, and with the edges of the fins, abdominal band, and variegations of the head blue.

Scarus Psittacus. *Forsk. arab. p. 29.*

Jaws blue: fins purple, edged with blue: back straiter in its outline than the abdomen: tail rising, spotted with blue, and subfurcated: lateral line double and ramified: eyes small. Native of the Arabian seas.

PURPLED SCARUS.

Scarus Purpuratus. *S. obscure viridis, vittis utrinque tribus purpureis longitudinalibus serratis, ventre caeruleo.*

Dull-green Scarus, with three longitudinal serrated purple bands on each side, and blue abdomen.

Scarus purpureus. *Forsk. arab. p. 27.*

An elegant species; in habit allied to the *Labri*: body abruptly lanceolate; the purple stripes on the body serrated at their upper edges: pectoral fins green, and marked at the tip by a large, lunated, marginal, black spot: dorsal and anal marked towards the base by a purple stripe: ventral fins blue: tail marked with longitudinal purple spots, and on each side by a purple stripe: shape slightly rounded: lateral line ramified: scales lax, as in the Mullet. Native of the Arabian seas: observed by Forskal.

BLACK SCARUS.

Scarus Niger. *S. ovato-oblongus, nigro-fuscus, labiis rudris, pinnarum marginibus viridi-cæruleis.*

Ovate-oblong, blackish-brown Scarus, with red lips, and the margins of the fins greenish-blue.

Scarus niger. *Forsk. arab. p. 28.*

NATIVE of the Arabian seas : lateral line double and ramified : tail greenish, and obtusely forked.

SCHLOSSERIAN SCARUS.

Scarus Schlosseri. *S. aureus, maculis utrinque quinque fuscis, dorso fusciscente, cauda subæquali.* *Lin. Syst. Nat. Gmel. p. 1282. Pall. spic. zool. 8. p. 41.*

Gold-coloured Scarus, with five dusky spots on each side, brownish back, and nearly even tail.

SIZE of a roach : body compressed and broad : scales large : eyes large : irides fulvous : lower mandible rather longer than the upper : ventral fins triangular, and connected to the abdomen by the intervention of a perpendicular membrane. Native of Java.

BLUE-STRIPED SCARUS.

Scarus Ghobban. *S. albidus*, squamis transversim subcaeruleo fasciatis, linea laterali duplici.

Whitish Scarus, with the scales marked by transverse blueish bands, and double lateral line.

Scarus Ghobban. *Forsk. arab. p. 28.*

SHAPE ovate: head marked by greenish-blue streaks: fins edged with a similar colour: tail entire: lateral line ramified. Native of the Arabian seas: observed by Forskal.

 FERRUGINOUS SCARUS.

Scarus Ferrugineus. *S. fusco-ferrugineus*, maxillis marginibusque pinnarum viridibus, caudâ aequali.

Brown-ferruginous Scarus, with the jaws and margins of the fins green, and even tail.

Scarus ferrugineus. *Forsk. p. 29.*

SHAPE ovate-oblong: jaws green: body ferruginous-brown: pectoral fins of the same colour: ventral and anal violet: dorsal and tail yellowish: at the base of the tail several long lanceolate scales: tail itself broad and rounded. Native of the Arabian seas.

SORDID SCARUS.

Scarus Sordidus. *S. fusco-ferrugineus elongatus, pinnis obscurioribus, cauda surgente aequali.*

Brown-ferruginous Scarus, with darker coloured fins, and rising, even tail.

Scarus sordidus. *Forsk. arab. p. 30.*

SHAPE slender: colour brown: dorsal fin of a deeper colour: pectoral fins and tail yellowish: ventral violet: lateral line double and ramified: native of the Arabian seas.

SCALY-TAILED SCARUS,

Scarus Harid. *S. squamis magnis, abdomine violaceo, cauda furcata, media basi squamosa.*

Scarus Harid. *Forsk. arab. p. 30.*

NATIVE of the Arabian seas: lateral line double, strait, ramified: dorsal, anal, and caudal fin violet-coloured: pectoral fins yellowish.

TRILOBATE SCARUS.

Scarus Trilobatus. *S. varius, cauda triloba.*

Variegated Scarus, with trilobate tail.

Le Scare trilobé. *Cepede. 4. p. 21.*

SLIGHTLY described by Cepede from the drawings and MSS. of Plumier: its principal character seems to be its very distinctly trilobate tail: the upper

jaw is longer than the lower : colours of the body various : two spiny rays and sixteen soft on the dorsal fin. Native of the American seas : size not particularized.

ROSTRATED SCARUS.

Scarus Rostratus. *S. maxillis productis.*

Scarus with elongated jaws.

L'Ostorhinque Fleurieu. *Cepede. 4. p. 24.*

SLIGHTLY described by Cepede from the MSS. of Commerson : differs, according to Cepede, from the Scari in no other respect except in having the jaws much elongated : the lower jaw is rather longer than the upper : the eyes large : the head without scales : the dorsal and anal fin rather short : the tail very large, lunated : and near it is a transverse band of a deep colour : no lateral line could be perceived : on the dorsal fin are eight spiny rays. No particular description is given either of the size or colour of this fish, which Cepede names *Fleurieu* in honour of Monsr. Fleurieu, of the national institute, well known by his hydrographical and other publications.

SPARUS. SPARUS.

Generic Character.

<i>Dentes</i> validi: <i>Primores</i> in aliis ordine simplici, in aliis, duplici, triplici, vel quadruplici dispositi.	<i>Teeth</i> strong: <i>front-teeth</i> , in some species disposed in a single row, in others in a double, treble, or quadruple row.
<i>Molares</i> (plerisque) validi, convexi, læves, in series dispositi, et quasi pavementum in ore efformantes.	<i>Grinders</i> (in most species) convex, smooth, and disposed in ranges, forming a kind of pavement in the mouth.
<i>Labia</i> crassa: <i>Opercula</i> multica, squamosa.	<i>Lips</i> thick: <i>Gill-covers</i> unarmed, smooth, scaly.

THE genus *Sparus* is extremely numerous, and as the greater number are exotic, very little is known of their history: a general survey is therefore all that can be expected: it may be observed that they are much allied to the *Labri*, and that the distinction between these two genera is not, in all cases, so clear as might be wished: in the *Systema Naturæ* of Linnæus an evident confusion takes place with respect to the characters of both.