

C02983

A
RUDIMENTARY TREATISE
ON
ANALYTICAL GEOMETRY
AND
CONIC SECTIONS.

BY
JAMES HANN,
MATHEMATICAL MASTER OF KING'S COLLEGE SCHOOL, LONDON;
HONORARY MEMBER OF THE PHILOSOPHICAL SOCIETY OF NEWCASTLE-UPON-TYNE,
JOINT AUTHOR OF MECHANICS FOR PRACTICAL MEN,
AUTHOR OF THEORY OF BRIDGES, JOINT AUTHOR OF NAUTICAL TABLES,
AUTHOR OF A TREATISE ON THE STEAM ENGINE,
AUTHOR OF THEORETICAL AND PRACTICAL MECHANICS,
AUTHOR OF PLANE AND SPHERICAL TRIGONOMETRY, AND AUTHOR OF
EXAMPLES ON THE INTEGRAL CALCULUS.

LONDON:
JOHN WEALE, 59, HIGH HOLBORN.

1850.

LONDON:
GEORGE WOODFALL AND SON,
ANGEL COURT, SKINNER STREET.

PREFACE.

IN compiling this work the Author has read nearly all the best writers on the subject,—BOURDON, HYMERS, O'BRIEN, SALMON, YOUNG, and WOOLHOUSE, all containing valuable information.

In an elementary treatise like the present, it is impossible to go into the subject in such a manner as O'BRIEN and SALMON have done in their able and extensive works; yet it is to be hoped that the student will find this work an useful introduction to the more elaborate writings of those two eminent mathematicians, as well as to contain almost, if not all, the properties of the Conic Sections most in use.

Mr. GOMPERTZ, one of the most eminent of our English mathematicians, has just published a valuable tract on Porisms, which he calls "Hints on Porisms, in a Letter to T. S. DAVIES, Esq., F.R.S., F.S.A.," and in which he solves the following beautiful problem-connected with the subject before us:—"A circle and a parabola cutting each other being given in position and magnitude, to find the points of intersection."

It is announced that Professor DAVIES is about to publish a treatise on this subject himself. This will be a work which those interested in Porisms will look forward to with the greatest interest, as Mr. DAVIES is undoubtedly inferior to no living geometer on these subjects.


The student will find many examples fully worked out in this treatise, to some of which there are no figures given, but so clearly indicated as to give little trouble in forming them.

ANALYTICAL GEOMETRY.

SECTION I.—STRAIGHT LINE.

(1.) THE position of a point in a plane surface is determined by referring it to what are termed its axes of co-ordinates, as follows:—

Let there be taken in a plane two indefinite right lines AY , AX , inclined to one another at a given angle YAX . Now, if a point P be taken in the plane, it is evident that if PN , a line parallel to AY , be drawn, the position of the point P will be *determined* if AN and PN be given, since no other than *one*




point can fulfil the conditions required by the relation of AN and PN . Now, AN and NP are termed the co-ordinates of the point P , AY and AX are called respectively the axis of ordinates and the axis of abscissæ, and from their relation to one another co-ordinatal axes. And since in using the co-ordinates of a point the abscissa is usually denoted by x , and the ordinate by y , AY and AX are respectively termed the axis of y and the axis of x . The point A is termed the *origin*. In general the co-ordinate axes are taken perpendicular to one another; they are then called rectangular axes, and the co-ordinates are termed rectangular co-ordinates.

(2.) We have said that a point P was *determined* by the co-ordinates AN and NP . This is not strictly true, for the

conditions imposed by the *magnitude* of the co-ordinates would be fulfilled also by another point, as shall be shewn.

(3.) Produce the axes of AY and AX , as in the figure, so that the plane surface may be portioned out into four quarters. Now, suppose it is required to *locate* the point whose co-ordinates are a and b . Let $AN = a$, $NP = b$; then P x_1 is one point fulfilling the required conditions. But AN_1 may be also taken $= a$, and $N_1P_1 = b$, $N_1P_2 = b$, $NP_3 = b$, and we shall therefore have



four points equally satisfying the original condition, viz., that the co-ordinates should be a and b . But it is evident that if such ambiguity attend our determination of the position of a point, there will often arise inconvenience. This defect algebraic geometry supplies in the following manner:— A being the origin, AN is equal to $+a$, $NP = +b$. But the abscissa of a point situated *on* the axis of y , is $= 0$; so that at the termination of the first quarter of space the abscissa is $= 0$. And, similarly, a line measured in the direction AN_1 , which is equal in magnitude to AN , must be analytically expressed by $-(a)$, to show that we have passed the limit of the *first* quarter, and come into the *second* quarter of space. In a similar manner, it will be easy to conclude that, if the ordinates be measured *below* the axis of x , they must likewise be affected with a different sign. The student should compare this with the change of sign in $\sin \theta$, $\cos \theta$, $\tan \theta$, &c., in passing through the four quadrants.

(4.) It is evident that, conformable to these remarks, we shall have the following results:—

$$\begin{array}{ll}
 \left. \begin{array}{l} +a \\ +b \end{array} \right\} & \text{determine the point } P \\
 \left. \begin{array}{l} -a \\ +b \end{array} \right\} & \dots\dots\dots P_1 \\
 \left. \begin{array}{l} -a \\ -b \end{array} \right\} & \dots\dots\dots P_2 \\
 \left. \begin{array}{l} +a \\ -b \end{array} \right\} & \dots\dots\dots P_3.
 \end{array}$$

We thus see that the position of a point is determined by means of its co-ordinates.

(5.) To complete the theory of points, we shall now proceed to determine an analytical expression for the distance between two given points.


To find the distance between two given points referred to rectangular axes.

Let the co-ordinates of M and $M' = (x'y')$, $(x''y'')$, and let $M'M = D$ be the distance; draw $M'R$ parallel to the axis of x , then in the right-angled triangle $M'RM$,

$$\text{we have } MM'^2 = MR^2 + M'R^2,$$

$$\text{but } MR = MP - RP = y' - y'', \quad M'R = PP' = x' - x'',$$

$$\therefore MM'^2 = D^2 = (y' - y'')^2 + (x' - x'')^2,$$



$$\therefore D = \sqrt{(y' - y'')^2 + (x' - x'')^2}^*.$$


This formula is general, and holds equally in the case where the points are situated in an opposite direction; with respect to one of the axes, it is only necessary to change the sign which corresponds to the change of position.

Thus, for example, to obtain the distance of two points, one of which, M , is in the angle YAX , and the other, M' , is in the angle YAX' , we must change the sign of x'' , which gives

* If the angle $MM'R$, or the angle which the right line MM makes with the axis of $x = \alpha$, we have $\frac{MR}{M'R} = \tan \alpha$; but $MR = y' - y''$, and $M'R = x' - x''$,

$$\therefore \tan \alpha = \frac{y' - y''}{x' - x''}.$$

$$D = \sqrt{(y' - y'')^2 + (x' + x'')^2},$$



for $M'M^2 = MR^2 + M'R^2$, and $MR = y' - y''$, as before,


$$\text{but } M'R = AP + AP' = x' + x'',$$

$$\therefore D = \sqrt{(y' - y'')^2 + (x' + x'')^2}.$$

If one of the points M be at the origin, then $x' = 0$, $y' = 0$, and the above becomes

$$D = \sqrt{y^2 + x^2}.$$

When the axes are oblique, the triangle $MM'R$ is oblique; then, by Hann's Trigonometry, page 56,



$$M'M^2 = MR^2 + M'R^2 - 2MR \cdot M'R \cos MRM';$$

but $MR = y' - y''$; $M'R = x' - x''$; and $\cos MRM' = -\cos MRK = -\cos \beta$ (β being the angle MRK , which is the same as the angle that the axes make with each other),

$$\therefore D^2 = (y' - y'')^2 + (x' - x'')^2 + 2(y' - y'')(x' - x'') \cos \beta,$$

$$\text{or } D = \sqrt{(y' - y'')^2 + (x' - x'')^2 + 2(y' - y'')(x' - x'') \cos \beta}.$$

SECTION II.

(6.) Curves are divided into orders or degrees corresponding to the sum of the exponents of x and y , and the term which contains the highest sum of exponents shows the order of the curve. Thus, $y = x + a$ is an equation of the first degree; $y^2 + xy + b^2 = 0$, $y^2 - x^2 + xy - c^2 = 0$, are equations of the second degree; and $y^3 + x^2 + x^2y + cxy^2 + bxy + cx + fy + d$ is an equation of the third degree.

Equation of the first degree.

(7.) The general form of this equation is $Ax + By + C = 0$, the constants A , B , C , being either positive or negative.

This equation may be put under the form $y = -\frac{A}{B}x - \frac{C}{B}$,

$$\text{or } y = mx + h \text{ if } m = -\frac{A}{B}, \quad h = -\frac{C}{B},$$


m and h being unrestricted as to sign.

So that we have two forms of the equation of the first degree,

$$Ax + By + C = 0 \dots (1), \quad \text{or } y = mx + h \dots (2),$$

both equally general.

(8.) To find the equation to a straight line.



Referring the line to the rectangular axes AX and AY ,

$$\text{let } AN = x, \quad PN = y, \quad AB = h.$$

By similar triangles, $\frac{PD}{BD} = \frac{P'D'}{BD'} = \text{a constant ratio,}$

$$\text{or } \frac{PN - DN}{BD} = \frac{PN - AB}{AN} = \text{a constant ratio,}$$

$$\therefore \frac{y - h}{x} = \text{a constant quantity ;}$$


but $\frac{PD}{BD} = \frac{y - h}{x} = \tan PBD = \tan PCX$, which call m ,

$$\text{and we have } \frac{y - h}{x} = m,$$

$y - h = mx$, or $y = mx + h$, the equation to the straight line.

(9.) For oblique co-ordinates, the ratio $\frac{PD}{BD}$ is constant, as before, but in this case we have

$$\frac{PD}{BD} = \frac{\sin PBD}{\sin BPD} = \frac{\sin PCX}{\sin PBY}.$$



If YAX , the inclination of the axes, $= \beta$, and PCX , the angle which the line PC makes with the axis of x , $= \alpha$, the above becomes

$$\frac{PD}{BD} = m = \frac{\sin \alpha}{\sin (\beta - \alpha)}.$$

(10.) To determine the equation to a right line in terms of the intercepts on the axes*.

The general equation to a right line is $Ax + By + C = 0$,

let $AG = a$ and $AH = b$;

when $x = 0$, $By + C = 0$,

$$\therefore y \text{ or } AH = -\frac{C}{B};$$

when $y = 0$, $Ax + C = 0$,

$$\therefore x \text{ or } AG = -\frac{C}{A}.$$


Divide the general equation by C ,

$$\frac{A}{C}x + \frac{B}{C}y + 1 = 0,$$

$$\text{but } \frac{A}{C} = -\frac{1}{AG} = -\frac{1}{a} \text{ and } \frac{B}{C} = -\frac{1}{AH} = -\frac{1}{b},$$

$$\therefore -\frac{1}{a}x - \frac{1}{b}y + 1 = 0,$$

or $\frac{x}{a} + \frac{y}{b} = 1$; this is the equation to HG , which cuts off the intercepts AH and AG , which are positive;



* The equation to the line $G'H$ is evidently

$$y = +mx + h.$$

The equation to the line $H'G$, cutting the axis of Y below the origin, is

$$y = mx - h.$$

The equation to HG , having m negative, is

$$y = -mx + h.$$

The equation to the line $G'H'$, which cuts the axis of x to the left of the origin, and the axis of y below it, having both m and h negative, is

$$y = -mx - h.$$

$\frac{x}{a} - \frac{y}{b} = 1$ is the equation to the line GH' , which cuts off a positive intercept on the axis of x , and a negative intercept on the axis of y ;

$\frac{y}{b} - \frac{x}{a}$ is the equation to the line HG' , which cuts off a negative intercept on the axis of x , and a positive intercept on the axis of y ;

$$\frac{x}{a} + \frac{y}{b} = -1 \text{ is the equation } G'H'.$$


(11.) Let the angle

$$PAG = \alpha, \quad AG = a,$$

$$AH = b,$$

the intercepts of the axes, and
let p = perpendicular AP ,

$$a \cos \alpha = p, \quad \therefore a = \frac{p}{\cos \alpha},$$



$$b \cos (90 - \alpha) = p, \quad \therefore b = \frac{p}{\cos (90 - \alpha)} = \frac{p}{\sin \alpha};$$

these substituted in the equation $\frac{x}{a} + \frac{y}{b} = 1$,

$$\text{give } x \cos \alpha + y \sin \alpha = p.$$

(12.) Find the equation to a straight line passing through a given point.

Let $x'y'$ be the co-ordinates of the given point; then, as the general equation for every point in the line is

$$y = mx + h \dots\dots\dots (1);$$

therefore, at the given point,

$$y' = mx' + h,$$

$$\therefore h = y' - mx' \dots\dots\dots (2).$$

Substitute this value of h in (1), and we have

$$y - y' = m(x - x'), \text{ or } y = m(x - x') + y',$$

$$\text{or } y = mx + y' - mx';$$

we have here determined h , but having no other given condition we cannot determine m , which, therefore, remains indeterminate, as it ought, for any number of lines may be drawn through one point.

(13.) To find the equation to a line passing through two given points.

Let M and M' be the two given points whose co-ordinates are x', y' , and x'', y'' .

The equation will be of the form,

$$y = mx + h \dots\dots\dots (1),$$

and since each of the points M and M' is in the line,

$$y' = mx' + h \dots\dots\dots (2),$$

$$y'' = mx'' + h \dots\dots\dots (3).$$

Subtract (3) from (2), and we have,

$$y' - y'' = m(x' - x''), \quad \therefore m = \frac{y' - y''}{x' - x''}.$$

Substituting this value in equation (2),

$$h = y' - \frac{y' - y''}{x' - x''} \cdot x' = \frac{x'y'' - y'x''}{x' - x''};$$

these values of m and h , substituted in equation (1) give,

$$y = \frac{y' - y''}{x' - x''} \cdot x + \frac{x'y'' - y'x''}{x' - x''} \dots\dots (4),$$

for the required equation.

Or thus :—subtract (2) from (1), and we have,

$$y - y' = m(x - x'),$$

an equation which still contains the unknown quantity m .

Subtracting (3) from (2) we obtain,

$$y' - y'' = m(x' - x''),$$

$$\therefore m = \frac{y' - y''}{x' - x''}.$$

The value of m substituted in the preceding equation gives

$$y - y' = \frac{y' - y''}{x' - x''} (x - x') \dots\dots\dots (5),$$

an equation which contains only given quantities.

The identity of the equations (4) and (5) may be easily shewn: we find, from equation (5),

$$y = y' + \frac{y' - y''}{x' - x''} \cdot x - \frac{y' - y''}{x' - x''} \cdot x';$$

which by reduction becomes,

$$y = \frac{y' - y''}{x' - x''} \cdot x + \frac{x'y'' - y'x''}{x' - x''}.$$

The second method is evidently much more simple and elegant than the first.

(14.) To draw through a given point a right line parallel to a line given in position.

Let x', y' be the co-ordinates of the point M , through which we can draw a line DH parallel to a given right line BL .

The equation of the first right line being

$$y = mx + h \dots\dots\dots (1),$$

that of the required line will be of the form

$$y = m'x + h' \dots\dots\dots (2),$$

m' and h' being the two constants which we must determine.

Since DH passes through the point M , the equation is

$$y' = m'x' + h' \dots\dots\dots (3).$$

Subtracting (2) from (3), we have $y - y' = m'(x - x')$; and because the lines are parallel, $m' = m$,

$$\therefore y - y' = m(x - x'),$$

which is the equation of the required line, and it differs only from the above equation by the ordinate at the origin, which is in this case $y' - mx'$.

(15.) Two right lines being given to find their point of intersection and also their angle of intersection.

Let $y = mx + h$ }
 $y = m'x + h'$ } be the equations to the two lines.

At the point of intersection the co-ordinates are the same for both lines; hence, at this point,

$$mx + h = m'x + h'$$

$$m'x - mx = h - h'$$

$$x = \frac{h - h'}{m' - m};$$

this substituted in the first equation gives

$$y = m \cdot \frac{h - h'}{m' - m} + h, \text{ or } y = \frac{hm' - mh'}{m' - m},$$

therefore $x = \frac{h - h'}{m' - m}$, $y = \frac{hm' - mh'}{m' - m}$, are the co-ordinates of the point of intersection.

If $m' = m$, we have $x = \frac{h - h'}{0}$, $y = \frac{m(h - h')}{0}$; that is, these values are infinite, as they ought to be, as the lines are parallel.

To find the angle of intersection. See figure, page 13.

Taking the axes rectangular and putting V for the angle of intersection; in the triangle EMG,

$$MGX = EMG + MEG,$$

$$\therefore EMG, \text{ or } V = MGX - MEG = \alpha' - \alpha,$$

α' and α being the angles that the two right lines form with the axis of x . Then by Hann's Trigonometry, p. 31, we have

$$\tan(\alpha' - \alpha), \text{ or } \tan V = \frac{\tan \alpha' - \tan \alpha}{1 + \tan \alpha' \tan \alpha} \dots\dots\dots (1),$$

and since the axes are rectangular, $\tan \alpha' = m'$, and $\tan \alpha = m$,

$$\therefore \tan V = \frac{m' - m}{1 + m'm} \dots\dots\dots (2).$$

When the axes are oblique, we have

$$\frac{\sin \alpha'}{\sin (\beta - \alpha')} = m', \quad \frac{\sin \alpha}{\sin (\beta - \alpha)} = m.$$

From the second relation,

$$\sin \alpha = m \sin (\beta - \alpha), \text{ or}$$

$$\sin \alpha = m \sin \beta \cos \alpha - m \sin \alpha \cos \beta.$$

Dividing by $\cos \alpha$,

$$\tan \alpha = m \sin \beta - m \tan \alpha \cos \beta,$$

$$\therefore \tan \alpha = \frac{m \sin \beta}{1 + m \cos \beta}.$$

In the same manner,

$$\tan \alpha' = \frac{m' \sin \beta}{1 + m' \cos \beta};$$

substituting these in (1) we have,

$$\tan V = \frac{\frac{m' \sin \beta}{1 + m' \cos \beta} - \frac{m \sin \beta}{1 + m \cos \beta}}{1 + \frac{m m' \sin^2 \beta}{(1 + m' \cos \beta)(1 + m \cos \beta)}},$$

reducing and observing that $\sin^2 \beta + \cos^2 \beta = 1$,

$$\tan V = \frac{(m' - m) \sin \beta}{1 + m m' + (m + m') \cos \beta} \dots\dots\dots (3).$$

If we suppose $\beta = 90$, $\sin \beta = 1$, and $\cos \beta = 0$, we have

$$\tan V = \frac{m' - m}{1 + m m'}, \text{ as before;}$$

$$\text{when } V = 90, 1 + m m' = 0, \therefore m' = -\frac{1}{m};$$

when the axes are oblique $1 + m \cdot m' + (m + m') \cos \beta = 0$,

$$\therefore m' = -\frac{1 + m \cos \beta}{m + \cos \beta}.$$

(16.) The relation $1 + m.m' = 0$ can be shewn from the figure. The triangle EMG is right-angled at M ; the two angles, MEG and MGE , are complements of each other,

$$\begin{aligned}\therefore \tan MGE &= \cot MEG \\ &= \frac{1}{\tan MEG},\end{aligned}$$

but $\tan MGX$, or $m' = -\tan MGE$, $\tan MEG = m$,

$$\therefore m' = -\frac{1}{m}, \text{ or } mm' + 1 = 0.$$



(17.) From a given point to draw a perpendicular to a given line, and find the length of this perpendicular.

Let BL be the given right line, and MG the perpendicular on BL , let x' and y' be the co-ordinates of the point M , and let the equation to the right line BL be

$$y = mx + h \dots\dots\dots (1),$$

since the right line MG passes through the point x' , y' , its equation (12) will be of the form

$$y - y' = m'(x - x');$$



but since these lines are to be perpendicular to each other, we have $mm' + 1 = 0$, or

$$m' = -\frac{1}{m};$$

therefore, the preceding equation becomes

$$y - y' = -\frac{1}{m}(x - x')^* \dots\dots\dots (2).$$

This is the equation to the perpendicular MG. Now, to find an expression for the length of the perpendicular MH, we have x', y' , the co-ordinates of the point M; and if we can find those of the point H, we have only to substitute these four co-ordinates in the expression for the distance between the given points (art. 5), and we shall have the distance MH.

As the point H is the intersection of BL and MG, we must eliminate x, y by the equations (1) and (2); thus, we must find the values of $x - x', y - y'$, considered as unknown; in order to do this we must put equation (1) in the form

$$y - y' = m(x - x') - y' + mx' + h \dots\dots\dots (3),$$

$$y - y' = -\frac{1}{m}(x - x') \dots\dots\dots (2).$$

Subtract, and we have

$$0 = \left(m + \frac{1}{m}\right)(x - x') - y' + mx' + h,$$

* If we take the general equation $Ax + By + C = 0$, then the equation to a right line perpendicular to it is $A(y - y') = B(x - x')$, for

$$y = -\frac{A}{B}x - \frac{C}{B},$$

$$\therefore y = mx + h, \text{ as } m = -\frac{A}{B}, \text{ and } h = -\frac{C}{B};$$

substituting this value for m , $y - y' = -\frac{1}{m}(x - x')$ becomes

$$y - y' = \frac{B}{A}(x - x'), \text{ or } A(y - y') = B(x - x').$$

$$\therefore x - x' = \frac{y' - mx' - h}{m + \frac{1}{m}} = \frac{m(y' - mx' - h)}{m^2 + 1};$$

this value being substituted in (2), gives

$$y - y' = -\frac{1}{m} \cdot \frac{m(y' - mx' - h)}{m^2 + 1} = -\frac{(y' - mx' - h)}{m^2 + 1}.$$

Substituting these values of $x - x'$, $y - y'$, in the expression for D, we have, putting P for the perpendicular,

$$\begin{aligned} P &= \sqrt{\frac{m^2(y' - mx' - h)^2 + (y' - mx' - h)^2}{(m^2 + 1)^2}} \\ &= \frac{\sqrt{(m^2 + 1)(y' - mx' - h)^2}}{(m^2 + 1)} = \frac{(y' - mx' - h)\sqrt{m^2 + 1}}{m^2 + 1}, \\ \therefore P &= \frac{\pm (y' - mx' - h)}{\sqrt{m^2 + 1}} \end{aligned}$$

for the length MH.

(18.) The double sign may be thus explained: y' represents the ordinate MP, and $mx' + h$ represents the ordinate NP of BL, corresponding to x' or AP; hence,

$$MN = MP - PN = y' - mx' - h.$$

Now, this distance may be either positive or negative, that is, it may be either greater than nothing or less than nothing, according as the point M is above or below BL; for example, if the point was in M', we should have

$$M'N' = N'P' - M'P' = mx' + h - y'.$$

When the point M is above the right line BL, in which case $y - mx' - h > 0$, we have

$$P = \frac{y' - mx' - h}{\sqrt{m^2 + 1}};$$

and if the point M is placed below the right line BL, then we have the condition $y' - mx' - h < 0$,

$$\therefore P = \frac{mx' + h - y'}{\sqrt{m^2 + 1}}.$$

We can obtain both these results geometrically; in fact, MP and MH being respectively perpendicular to AP and BL, it follows that the angle NMH equals the angle BLX = α . Now, in the right-angled triangle NMH, we have

$$MH = MN \cos \alpha = \frac{MN}{\sec \alpha} = \frac{MN}{\sqrt{1 + \tan^2 \alpha}},$$

but $MN = MP - NP = y' - mx' - h$, and $\tan \alpha = m$,

$$\therefore MH = P = \frac{y' - mx' - h}{\sqrt{m^2 + 1}}.*$$

If the point M was below BL, we should have

$$\begin{aligned} M'N' &= mx' + h - y', \\ \therefore P &= \frac{mx' + h - y'}{\sqrt{m^2 + 1}}. \end{aligned}$$

(19.) If the point from which we wish to let fall the perpendiculars be at the origin, we have in this case $x' = 0$, $y' = 0$, then

$$P = \mp \frac{h}{\sqrt{m^2 + 1}};$$

this result is positive or negative, according as the point B is above or below the origin.

(20.) Through a given point to draw a straight line which shall make a given angle with a given straight line.

Let the given line be represented by the equation $y = mx + h$, then, because the required line passes through the given point x', y' , its equation will be of the form $y - y' = m'(x - x')$, and since these right lines ought to form an angle whose tangent is t , we have (art. 15.)

* This is sometimes given $P = \frac{Ax' + By' + C}{\sqrt{A^2 + B^2}}$, which is found by putting $-\frac{A}{B}$ for m , and $-\frac{C}{B}$ for h .

$$\frac{m' - m}{1 + mm'} = t, \text{ or } \frac{m - m'}{1 + mm'} = t;$$

either of these equations expresses the tangent of the given angle.

We can comprehend these two expressions in one, thus

$$\frac{m' - m}{1 + mm'} = \pm t, \quad \therefore m' = \frac{m \pm t}{1 \mp mt},$$

this value substituted in the above equation for m' , gives the equation of the required line.

$$y - y' = \frac{m \pm t}{1 \mp mt} (x - x').$$

Examples on the Straight Line.

1. The equation $(2y + x)(3y - x) = 0$ represents two straight lines inclined to one another at an angle of 135° .

The equation is at once resolved into the two

$$\left. \begin{array}{l} 2y + x = 0 \\ \text{and } 3y - x = 0 \end{array} \right\},$$

$$\left. \begin{array}{l} \text{or (1) } \dots\dots\dots y = -\frac{1}{2}x \\ \text{and (2) } \dots\dots\dots y = \frac{1}{3}x \end{array} \right\},$$

each of which represents a straight line passing through the origin, since $x = 0$, gives $y = 0$.

Now, the coefficient of x is the tangent of the angle which the line makes with the axis of x .

Hence, if θ_1 be the angle which (1) makes with axis, and θ_2 be the angle which (2) makes with the same,

$$\left. \begin{array}{l} \tan \theta_1 = -\frac{1}{2} \\ \text{and } \tan \theta_2 = \frac{1}{3} \end{array} \right\}.$$

Now, if ϕ be the angle between the lines (1) and (2),
 $\phi = \theta_1 - \theta_2$,

$$\therefore \tan \phi = \frac{\tan \theta_1 - \tan \theta_2}{1 + \tan \theta_1 \tan \theta_2} = \frac{-\frac{1}{2} - \frac{1}{3}}{1 - \frac{1}{2} \cdot \frac{1}{3}} = -\frac{\frac{5}{6}}{\frac{5}{6}} = -1,$$

$$\therefore \phi = 135^\circ.$$

2. Find the equation to a straight line which bisects the angle between the lines

$$5y - 2x = 0 \dots\dots (1), \text{ and } 3y + 4x = 12 \dots\dots (2).$$

Let (x, y) be any point in the required line; then, since this line bisects the angle between (1) and (2), the perpendiculars from (x, y) on these must be equal.

Substituting in the formula $P = \frac{Ax + By + C}{\sqrt{A^2 + B^2}}$, we get the perpendicular on (1)

$$= \frac{-2x + 5y}{\sqrt{4 + 25}},$$

$$\text{and that on (2)} = \frac{4x + 3y - 12}{\sqrt{16 + 9}},$$

$$\therefore \frac{5y - 2x}{\sqrt{29}} = \frac{4x + 3y - 12}{5} \text{ is equation required,}$$

$$i. e. (25 - 3\sqrt{29})y - (10 + 4\sqrt{29})x + 12\sqrt{29} = 0.$$

But $\sqrt{29} = 5.38$, \therefore equation becomes

$$8.86y - 31.52x + 64.56 = 0.$$

3. Find the angle between the lines whose equations are

$$\left. \begin{array}{l} 1 + 3x + 2y = 0 \\ \text{and } 3 + 2x - 3y = 0 \end{array} \right\}.$$

These equations become

$$y = -\frac{3}{2}x - \frac{1}{2} \dots\dots\dots (1),$$

$$\text{and } y = \frac{2}{3}x + 1 \dots\dots\dots (2),$$

therefore, if θ_1 be the angle which (1) makes with the axis of x , and θ_2 the angle which (2) makes with the same,

$$\tan \theta_1 = -\frac{3}{2}, \text{ and } \tan \theta_2 = \frac{2}{3},$$

$$\therefore \tan \theta_1 \cdot \tan \theta_2 = -1;$$

hence the lines are at right angles to each other.

4. Find the length of the perpendicular from the point $x = 3, y = 5$, upon the line $7x - 3y = 9$.

The length of a perpendicular from a point x, y , upon the line $Ax + By + C = 0$, is

$$\frac{Ax + By + C}{\sqrt{A^2 + B^2}}.$$

In the present example $x = 3, y = 5, A = 7, B = -3$, and $C = -9$,

$$\therefore p = \frac{21 - 15 - 9}{\sqrt{49 + 9}} = -\frac{3}{\sqrt{58}},$$

or, independently of the sign, the length of the perpendicular equals $\frac{3}{\sqrt{58}}$.

5. The following equation represents two straight lines at right angles to each other.

$$2y^2 - 3xy - 2x^2 - y + 2x = 0 \dots\dots\dots (1),$$

$$\therefore 2y(y - 2x) + x(y - 2x) - (y - 2x) = 0,$$

$$\therefore (2y + x - 1)(y - 2x) = 0.$$

Hence, equation (1) is resolved into the two

$$\begin{aligned} & 2y + x - 1 = 0 \} \\ \text{and } & y - 2x = 0 \} \end{aligned}$$

which evidently represent two lines at right angles to each other, for they are each equations of the first degree, and therefore represent straight lines; and since the coefficients of x divided by those of y in the two equations when multiplied together $= -1$, the lines are at right angles to each other.


6. In the figure to Euclid, book i. prop. 47, prove that the lines FC, KB, and AL meet in one point.

Take A for the origin, AG and AH for co-ordinate axes.

Now, equation

$$\text{to FC is } \frac{x}{AD} + \frac{y}{AC} = -1,$$

$$\text{and to KB is } \frac{x}{AB} + \frac{y}{AE} = -1,$$



$$\text{and } AD : AC :: FG : GC :: AB : AB + AC,$$

$$\therefore AD = \frac{AB \cdot AC}{AB + AC}, \text{ and } \frac{1}{AD} = \frac{1}{AC} + \frac{1}{AB};$$

$$\text{similarly, } AE : AB :: HK : BH :: AC : AB + AC,$$

$$\therefore AE = \frac{AB \cdot AC}{AB + AC}, \text{ and } \frac{1}{AE} = \frac{1}{AC} + \frac{1}{AB},$$

$$\therefore \text{equation to FC becomes } x \left(\frac{1}{AC} + \frac{1}{AB} \right) + \frac{y}{AC} = -1,$$

$$\text{and to KB } \dots \frac{x}{AB} + y \left(\frac{1}{AC} + \frac{1}{AB} \right) = -1,$$

$$\therefore \text{for P their intersection } \frac{x}{AC} - \frac{y}{AB} = 0,$$

$$\therefore \frac{AN}{NP} = \frac{AC}{AB}, \therefore \text{ANP is similar to ABC,}$$

and \therefore P is in AL, that is, the perpendicular from A on BC.—

7. In the equation $ay^2 + bxy + dx + ey = 0$, find the relations among the coefficients, that it may represent two straight lines.

The condition is, that the equation may be resolvable into two factors of the first degree, and since we observe that if $x = 0$, $y = 0$, and that the coefficient of $x^2 = 0$, we assume the equation to become

$$(Ax + By)(Cy + 1) = 0,$$


$$\therefore ACxy + BCy^2 + Ax + By = 0;$$

By comparing this with the original equation we get the relation amongst the coefficients—viz.,

$$\left. \begin{aligned} a &= BC \\ b &= AC \\ d &= B \\ e &= A \end{aligned} \right\},$$

$\therefore ae = A \cdot B \cdot C = bd$ is the relation required.

8. ABCD is a parallelogram, AE and ED parallelograms about its diagonal CB. Show that the diagonals of these, viz., HF and GL, meet CB in the same point.



Take E for the origin, EF and EL for co-ordinate axes.

let HE = a , EF = b , EL = a' , and EG = b' .

Then equation to HF is $-\frac{x}{a} + \frac{y}{b} = 1$,

$$\dots \quad \text{GL} \quad \dots \quad \frac{x}{a'} - \frac{y}{b'} = 1,$$

$$\text{and} \quad \dots \quad \text{CB} \quad \dots \quad x = \frac{a'}{b} y = \frac{a}{b'} y;$$

therefore, for intersection of CB and HF,

$$x \left(\frac{1}{a'} - \frac{1}{a} \right) = 1, \therefore x = \frac{aa'}{a - a'},$$

$$\text{and } y = \frac{bx}{a'} = \frac{ab}{a - a'};$$

and for intersection of CB and GL,


$$x \left(\frac{1}{a'} - \frac{1}{a} \right) = 1, \therefore x = \frac{aa'}{a-a'},$$

$$\text{and } y = \frac{ab}{a-a'};$$

hence, HF and GL intersect CB in the same point.

9. Prove that right lines drawn from the three angles of a triangle to the middle of the opposite sides intersect in the same point.

Let the axis of x coincide with the base of the triangle, and the axis of y be AY, the axes being rectangular. We



must now determine the equations of the lines AF, BE, CD, but before we do this we must find the co-ordinates of the points A, B, C, D, E, F.

For the co-ordinates of the point A ... ($y = 0, x = 0$).

If $AB = c$, then the co-ordinates of } ($y = 0, x = c$).
the point B }

For the point C ($y = y', x = x'$).

Now, as D, E, F, are in the middle of AB, AC, BC, we have

$$AD = \frac{AB}{2} = \frac{c}{2}, \quad EI = \frac{CH}{2} = \frac{y'}{2}, \quad AI = \frac{x'}{2},$$

$$FG = \frac{CH}{2} = \frac{y'}{2}, \quad GH = \frac{BH}{2} = \frac{c - x'}{2},$$

$$\therefore AG = x' + \frac{c - x'}{2} = \frac{c + x'}{2};$$

this gives the co-ordinates of the points.

$$D \dots\dots\dots \left(y = 0, \quad x = \frac{c}{2} \right),$$

$$E \dots\dots\dots \left(y = \frac{y'}{2}, \quad x = \frac{x'}{2} \right),$$

$$F \dots\dots\dots \left(y = \frac{y'}{2}, \quad x = \frac{c + x'}{2} \right).$$

Since each of the lines AF, BE, CD, passes through two points, and the general equation to a right line through two points is

$$y - y' = \frac{y' - y''}{x' - x''} (x - x'),$$

substituting for x' , y' , y'' , x'' , the corresponding values in the figure under consideration, we shall obtain y and x for each particular line.

Since AF passes through the two points

$$(0, 0) \quad \frac{y'}{2}, \quad \left(\frac{c + x'}{2} \right),$$

$$\text{its equation is } y - 0 = \frac{0 - \frac{y'}{2}}{0 - \frac{c + x'}{2}} (x - 0),$$

$$\therefore y = \frac{y'}{c + x'} x \text{ is the equation to AF} \dots\dots (1);$$

$$\text{and, since BE passes through } (0, c) \left(\frac{y'}{2}, \quad \frac{x'}{2} \right),$$

$$\text{its equation is } y - 0 = \frac{0 - \frac{y'}{2}}{c - \frac{x'}{2}} (x - c),$$

$$\text{or } y = \frac{y'}{-2c + x'} (x - c) \dots\dots\dots (2).$$

$$\text{And since CD passes through } (y', x') \left(0, \quad \frac{c}{2} \right),$$

$$\text{its equation is } (y - y') = \frac{y'}{x' - \frac{c}{2}} (x - x'),$$

$$\text{or } y = y' + \frac{y'}{\left(x' - \frac{c}{2}\right)} (x - x'),$$

$$y = \frac{y'x' - \frac{y'c}{2} + y'x - y'x'}{x' - \frac{c}{2}} \\ = \frac{y'x - \frac{y'c}{2}}{\left(x' - \frac{c}{2}\right)} = \frac{y'(2x - c)}{2x' - c} \dots\dots\dots (3);$$

$$\text{and from these } x = \frac{c + x'}{3}, y = \frac{y'}{3},$$

Or having the co-ordinates of AF, BE, CD, we may proceed as follows:—

Since AF passes through the origin, its equation is of the form $y = ax$, and since it passes through the point $\left(\frac{y'}{2}, \frac{c + x'}{2}\right)$, we have the relation

$$\frac{y'}{2} = a \left(\frac{c + x'}{2}\right), \therefore a = \frac{y'}{c + x'},$$

$$\therefore \text{the equation of AF is } y = \frac{y'}{c + x'} x \dots\dots (1).$$

The right line BE passing through the point B, or $(c, 0)$ its equation is of the form $y = a'(x - c)$. In the equation $y - y' = a'(x - x')$, we must put 0 for y' , and c for x' , and a for a' , but as this line passes through the point E, or

$$\left(\frac{y'}{2}, \frac{x'}{2}\right), \text{ we have } \frac{y'}{2} = a' \left(\frac{x'}{2} - c\right), \therefore a' = \frac{y'}{x' - 2c};$$

hence the equation to the line BE is

$$y = \frac{y'}{x' - 2c} (x - c) \dots\dots\dots (2.)$$

In the same manner for CD,

$$y = \frac{y'}{2x' - c} (2x - c) \dots\dots\dots (3.)$$

From equations (1) and (2),

$$\frac{y'}{c + x'} \cdot x = \frac{y'}{x' - 2c} (x - c),$$

$$x'y' \cdot x - 2cy' \cdot x = cy' \cdot x - c^2y' + x'y' \cdot x - cx'y',$$

or by reduction,

$$3cy' \cdot x = cy' (c + x'), \quad \therefore x = \frac{c + x'}{3},$$

substitute this value in (1), and we have $y = \frac{y'}{3}$.

From equations (1) and (3) we have,

$$\frac{y'}{c + x'} \cdot x = \frac{y'}{2x' - c} (2x - c)$$

$$2x'y' \cdot x - cy' \cdot x = 2cy' \cdot x - c^2y' + 2x'y' \cdot x - cx'y',$$

$$\text{or } 3cy' \cdot x = cy' (c + x'), \quad \therefore x = \frac{c + x'}{3};$$

$$\text{and, consequently, } y = \frac{y'}{3}.$$

Hence we see that the co-ordinates of the point of intersection of the two right lines AF and BE are identical with those of the point of intersection of AF and CD; thus the three lines intersect in the same point.

In reflecting upon the preceding analysis, it is easy to see that the procedure is the same whatever be the inclination of the axes.

However, they become more simple when we take AB for the axis of x , and take for the axis of ordinates a straight line AY parallel to CD, which we can do when the line CD is given in position.

In that case, it is plain that the abscissa x' of the point C = AD, or $\frac{c}{2}$, where $c = 2x'$, and the equations (1), (2), (3), will be as follows:—

$$\text{the equation to the straight line AP, } y = \frac{y'}{3x'}x, *$$

$$\text{the equation of the straight line BE, } y = -\frac{y'}{3x'}(x - 2x'),$$

$$\text{the equation of the straight line CD, } x = x',$$

(since the last line is parallel AY.)

These things being premised, combining the last equation with the first, there results $y = \frac{y'}{3}$; and combining it with the second, there results again $y = \frac{y'}{3}$.

Thus the co-ordinates of the point of intersection of CD, AE, and of CD, BE, are

$$x = x', \quad y = \frac{y'}{3} = \frac{CD}{3}.$$


This proves that the choice of axes influences the simplicity of calculation in resolving questions by Analytical Geometry.

10. To prove that the perpendiculars drawn from the angular points of a triangle cut one another in the same point.

* Since the line AF passes through the origin and makes an angle with axis of abscissæ $= \tan^{-1} \frac{y'}{3x'}$, and since the line BE makes an angle with the same axis $= \tan^{-1} - \frac{y'}{3x'}$, and cuts the axis at a distance $2x'$ from the origin, its equation is $y = -\frac{y'}{3x'}(x - 2x')$.

Referring to rectangular axes, since we have to consider the relation of perpendiculars from two sides, take for the axis of abscissæ the line AB, and for the axis of ordinates a line perpendicular to AX, at A.

Let $c = AB$, x', y' the co-ordinates of the point C, we shall have for the equation of CD, $x = x' (1)$, since it is parallel to the axis of y .



Having found the equations of AP and BE, we must commence to find that of the lines CB, AC, to which they are perpendicular.

Since the line CB passes through two points (y', x') , and $(0, c)$ its equation is $y - y' = a(x - x')$, a having for its value $\frac{y'}{x' - c}$ when we put $y' = 0$, $x' = c$.

That of the line AC passing through the origin is $y = mx$, m being $= \frac{y'}{x'}$, since the point C is on that line.

These things being done, as AF passes through the origin, it has an equation $y = a'x$; and, as it is perpendicular to CB, we deduce the equation $aa' + 1 = 0$, where $a' = -\frac{1}{a} = \frac{c - x'}{y'}$, since the equation of AF is $y = \frac{c - x'}{y'} x$. (2).

The line BE passing through the point B, where the co-ordinates are $0, c$, has for its equation $y = m'(x - c)$.

But since it is perpendicular to AC, there results the relation $mm' + 1 = 0$, where $m' = -\frac{1}{m} = -\frac{x'}{y'}$.

Lastly, therefore, the equation BE is

$$y = \frac{-x'}{y'}(x - c) \dots\dots\dots (3).$$

Now, if we combine the equations (1) and (2),

$$\text{we find for } x = x', \quad y = \frac{(c - x)x'}{y'}.$$

The general equation admits of modification according as we take the origin.

If A, a point in the circumference, be taken as the origin, we have $\alpha^2 + \beta^2 = r^2$.

From the form (1)

$$x^2 - 2\alpha x + \alpha^2 + y^2 - 2\beta y + \beta^2 = r^2,$$

$$\therefore \text{since } (\alpha^2 + \beta^2) = r^2,$$

$$\text{we have } x^2 - 2\alpha x + y^2 - 2\beta y = 0 \dots (3).$$

If O, the centre, be the origin, the equation is (since α and $\beta = 0$)

$$x^2 + y^2 = r^2 \dots (4).$$

If the origin be at A'', the extremity of the diameter which coincides with the axis of x , we have

$$\bullet \quad x^2 + y^2 = 2rx \dots (5).$$

If in (3) we make $y = 0$, there results $x^2 - 2\alpha x = 0$, $\therefore x(x - 2\alpha) = 0$, $\therefore y = 0$ is satisfied by either $x = 0$ or $x = 2\alpha$, which shows that $x = 0$; $y = 0$ is placed on the circumference, as the value 0 of y corresponds to $x = 2\alpha$ as well as to $x = 0$; it follows that the circumference cuts the axis of x at a second point, C, where A'C is double of A'D, therefore the chord A'C is cut into two equal parts by the perpendicular from the centre of the circle, a well-known geometrical property of the circle.

Taking (5), $y^2 = 2rx - x^2 = x(2r - x)$.

$$\text{Now, } y = MR, \quad x = A''R, \quad 2r - x = GR,$$

$$\therefore MR^2 = A''R \times GR,$$

$$\therefore A''R : MR :: MR : GR;$$

that is, the perpendicular let fall from a point in the circumference on a diameter, is a mean proportional between the two segments. The equation may be put $y^2 + x^2 = 2rx$, but if we draw the chord A''M,

$$MR^2 + A''R^2 = A''M^2, \quad 2r = A''G, \quad x = A''R,$$

These things being premised, to express where the two circumferences cut, and to obtain the points of intersection, we must eliminate x and y by combining (1) and (2).

This gives

$$x = \frac{r^2 - r'^2 + d^2}{2d} \dots\dots\dots (3),$$

$$y = \pm \frac{1}{2d} \sqrt{4d^2r^2 - (r^2 - r'^2 + d^2)^2} \dots\dots\dots (4).$$

We now can proceed to the discussion of the point proposed for consideration.

If the quantity (4) under the radical sign is positive, in that case the values of x and y are real, and the circumferences have two common points.

Proving that the two points of intersection are on the same abscissa, OP, having the two ordinates equal, but of opposite signs.

Therefore, at every *cutting* of the two circles the line joining their centres is perpendicular to the common chord, and bisects it.

However, since y may be real or imaginary (x being always real), we will transform the above expression.

The quantity (4) under the radical is evidently

$$= (2dr + r^2 + d^2 - r'^2)(2dr - r^2 - d^2 + r'^2)$$

$$= \{(r + d)^2 - r'^2\} \{r'^2 - (r - d)^2\} =$$

$$(r + r' + d)(r - r' + d)(r + r' - d)(r' - r + d),$$

$$\therefore y = \pm \frac{1}{2d} \sqrt{(r + r' - d)(r - r' + d)(r + r' + d)(r' - r + d)};$$

under that form, as the third factor $r + r' + d$, under the radical sign, is necessarily positive, y will be real, according as the three other factors are positive, or one of the two positive, and the other two negative.

But this last circumstance cannot exist, since, if one of the three factors is negative, the other two are necessarily positive. For example, suppose $r + d - r'$ is < 0 ,

$$\begin{aligned} \therefore r + d < r', \quad \therefore r < r' \text{ and } d < r', \\ \therefore r' - r + d \text{ and } r' - d + r \text{ are positive,} \end{aligned}$$

and, *à fortiori*, the three factors cannot all be negative at once.

Therefore, the only case is where the three factors are positive at once, and in that case y is real; that is to say, the two circles cut one another when the sum of each two of the quantities r, r', d is $>$ the third.

Now, if one of the three factors is $(-)$, and the other two $(+)$, in that case y is imaginary, and the point is not a point of intersection. Therefore, the two circumferences have not a common point.

But suppose that any one of the factors is $= 0$, it gives, therefore, $y = \pm 0$, and the two circumferences have only one common point, which is necessarily on the line joining the centres, where the ordinate is zero; therefore, two circles touch when the distance of the centres is equal to the sum or difference of the radii.

Perhaps no example than this more strikingly shews the advantages resulting from the application of Algebra to Geometry.

(23.) To find the equation to the tangent to a circle.

Let us at first consider a line to pass through two points whose co-ordinates are (x', y') , (x'', y'') , the equation to this line is, by art. 13,

$$y - y' = \frac{y' - y''}{x' - x''} (x - x') \dots\dots\dots (1),$$

and if we combine this with the two equations

$$x'^2 + y'^2 = r^2 \dots\dots\dots (2),$$

$$x''^2 + y''^2 = r^2 \dots\dots\dots (3),$$

these two latter equations hold because each point (x', y') , (x'', y'') , are in the circumference of the circle.

Subtract (3) from (2), and we have

$$y'^2 - y''^2 + x'^2 - x''^2 = 0 \dots\dots\dots (4);$$

but since the difference of the squares of any two quantities

is equal to the product of the sum and difference of the same quantities, we have

$$(y' + y'')(y' - y'') + (x' + x'')(x' - x'') = 0;$$

$$\therefore \frac{y' - y''}{x' - x''} = -\frac{x' + x''}{y' + y''}.$$

If this value be substituted in (1) it becomes

$$y - y' = -\frac{x' + x''}{y' + y''}(x - x') \dots \dots \dots (5).$$


When the two points merge into one, the secant becomes a tangent, and we have $x' = x''$, and $y' = y''$, which gives

$$y - y' = -\frac{x''}{y''}(x - x''), \text{ and } x''^2 + y''^2 = r^2.$$

By means of these two equations we obtain the equation to the tangent in a more simple form. Clearing the former of fractions, we get

$$y y'' + x x'' = x''^2 + y''^2, \text{ or } y y'' + x x'' = r^2, \\ \text{since by the latter, } x''^2 + y''^2 = r^2.$$

This latter form is easily remembered, from its analogy to the equation of the circle. If in this latter form of the tangent we make $y = 0$, then $x x'' = r^2$, or $x = \frac{r^2}{x''}$, which is the abscissa OR of the point where the tangent meets the axis of x' ; also we have $PR = OR - OP$; or,




$$PR = \frac{r^2}{x''} - x'' = \frac{r^2 - x''^2}{x''}.$$

This line PR is called the subtangent,

Transformation of Co-ordinates.

(24.) The most general problem of transformation is as follows.—an equation being given between x and y , referred to any whatsoever axes, to find the equation when we pass to two new axes

Let AY, AX , be two axes to which the curve $MM'M''$ is referred by means of some equation $F(x, y) = 0$; $A'X', A'Y'$, two new axes.



Let $AP, PM = x$, and y , respectively, be the old co-ordinates, and x', y' , or $A'P', P'M'$, the new ones.

We seek now to express x', y' , in terms of x, y , or *vice versa*.

Let $A'X''$ and $P'H$ be drawn parallel to $AX, A'Y''$, and $P'K$ parallel to AY produce $A'Y''$ to B .

Let $AB = a, A'B = b, X'A'X'' = \alpha, Y'A'X'' = \alpha',$ and $Y''A'X'' = \epsilon$.

a and b are known quantities, the co ordinates of the new origin

As to the angle ϵ , it is always given (*a priori*), since it is the angle which the old axes make with each other.

We shall now proceed to give the principal cases

COR. I The most simple of all cases is that in which the two new axes are parallel to the old ones; that is to say, the axes continue in the same direction, the origin alone being different. Then,

$$\begin{cases} x = x' + a \\ y = y' + b \end{cases} \dots\dots (1),$$

$$AP \text{ or } x = AB + BP = a + A'K + P'H,$$

$$PM \text{ or } y = A'B + ML = b + P'K + MH;$$

thus, these being reduced, we can determine $A'K, P'K, P'H$, and MH .

But, in the triangles $\Lambda'P'K$, $MP'H$, we have, by Hann Trigonometry, page 56,

$$(1.) \Lambda'K : \Lambda'P' :: \sin \Lambda'P'K : \sin \Lambda'KP';$$

or because $\Lambda'P'K = P'A'Y'' = \zeta - \alpha$, and

$$\Lambda'KP' = \Lambda'LM = \pi - MLX'' = \pi - \zeta,$$

$$\Lambda'K : x' :: \sin (\zeta - \alpha) : \sin \zeta, \quad \therefore \Lambda'K = \frac{x' \sin (\zeta - \alpha)}{\sin \zeta}.$$

$$(2.) P'K : \Lambda'P' :: P'A'K : \sin \Lambda'KP', \quad \therefore P'K = \frac{x' \sin \alpha}{\sin \zeta}$$

$$(3.) P'H : MP' :: \sin P'MH : \sin P'HM,$$

or because $P'MH = Y'A'Y'' = (\zeta - \alpha')$, and

$$P'HM = \Lambda'LM = \pi - \zeta,$$

$$P'H : y' :: \sin (\zeta - \alpha') : \sin \zeta, \quad \therefore P'H = \frac{y' \sin (\zeta - \alpha')}{\sin \zeta}.$$

$$(4.) MH : MP' :: \sin MP'H : \sin P'HM, \quad \therefore MH = \frac{y' \sin \alpha'}{\sin \zeta},$$

therefore, putting these values in the expressions for x , y we obtain

$$\left. \begin{aligned} x &= \frac{x' \sin (\zeta - \alpha) + y' \sin (\zeta - \alpha')}{\sin \zeta} - a \\ y &= \frac{x' \sin \alpha + y' \sin \alpha'}{\sin \zeta} + b \end{aligned} \right\} (2).$$

These are the formulas most general in the transformation of co-ordinates of which it is easiest to deduce particular corresponding formulas to all the positions of the new origin and to the different directions of the new axes with respect to the old ones, in giving to a , b , these suitable values, positive or negative, and to the angles α and α' all values from 0 up to π .

2. To pass from a system of rectangular to a system of oblique axes.

In this case, let $\mathfrak{C} = \frac{\pi}{2}$, $\therefore \sin \mathfrak{C} = 1$, $\sin (\mathfrak{C} - \alpha) = \cos \alpha$,
and $\sin (\mathfrak{C} - \alpha') = \cos \alpha'$; consequently, we have

$$\left. \begin{aligned} x &= x' \cos \alpha + y' \cos \alpha' + a \\ y &= x' \sin \alpha + y' \sin \alpha' + b \end{aligned} \right\} \dots\dots\dots (3).$$

3. To pass from one rectangular system to another.

In that case, $\mathfrak{C} = \frac{\pi}{2}$, $\alpha' = \frac{\pi}{2} + \alpha$,

$$\sin \mathfrak{C} = 1, \quad \sin (\mathfrak{C} - \alpha) = \cos \alpha,$$

$$\sin (\mathfrak{C} - \alpha') = \sin \left(\frac{\pi}{2} - \frac{\pi}{2} - \alpha \right) = -\sin \alpha,$$

$$\sin \alpha' = \sin \left(\frac{\pi}{2} + \alpha \right) = \cos \alpha,$$

$$\therefore \left. \begin{aligned} x &= x' \cos \alpha - y' \sin \alpha + a \\ y &= y' \cos \alpha + x' \sin \alpha + b \end{aligned} \right\} \dots\dots\dots (4).$$

This is a very useful transformation.

4. To change from an oblique to a rectangular system.

In the general formula $\alpha' = \frac{\pi}{2} + \alpha$, $\sin \alpha' = \cos \alpha$,

$$\sin (\mathfrak{C} - \alpha') = \sin \left(\mathfrak{C} - \frac{\pi}{2} - \alpha \right) = -\cos (\mathfrak{C} - \alpha),$$

$$\therefore \left. \begin{aligned} x &= \frac{x' \sin (\mathfrak{C} - \alpha) - y' \cos (\mathfrak{C} - \alpha)}{\sin \mathfrak{C}} + a \\ y &= \frac{x' \sin \alpha + y' \cos \alpha}{\sin \mathfrak{C}} + b \end{aligned} \right\} \dots\dots\dots (5).$$


If the angle α be negative, then its sine will be negative, but the cosine will remain positive; we have the following modification, which will be found useful;

$$\left. \begin{aligned} x &= \frac{x' \sin \alpha' - y' \cos \alpha'}{\sin (\alpha' - \alpha)} \\ y &= \frac{y' \cos \alpha - x' \sin \alpha}{\sin (\alpha' - \alpha)} \end{aligned} \right\} \dots\dots\dots (6).$$

SECTION IV.—PARABOLA.

(25.) If there be taken the fixed line VF , and a fixed point S , then if $SP = VP$, the perpendicular distance of P from the line VF , the locus of P , is the *parabola*. Through S draw the perpendicular LS , and bisect it in A , then it is evident that A is a point in the curve.

To find the equation to the parabola. Referring to rectangular co-ordinates, let A be the origin, $AN = x$, $NP = y$, $LA = AS = a$;



then $VP = LN = SP$,

$$\therefore LN^2 = (x + a)^2 = S'P^2 = y^2 + (x - a)^2, \therefore y^2 = 4ax.$$

If S be the origin, then it is evident that $y^2 = 4a(x + a)$, for in that case $SN = x$.

If L be the origin, then $y^2 = 4a(x - a)$, since LN is then $= x$.

Polar Equations.

If S be the pole, $ASP = \theta$,

$$\text{then } r = SP = LN = 2a - 2r \cos \theta, \therefore r = \frac{2a}{1 + \cos \theta},$$

$$\text{or } = \frac{a}{\sin^2 \frac{\theta}{2}}. \quad (\text{Hann's Trigonometry, page 27.})$$

It is evident that since $4a$ may be made to equal any constant quantity, the equation to the parabola may be written $y^2 = px$. If $x = a$, then the corresponding value of y is $2a$, therefore the double ordinate through the point is $= 4a$. This double ordinate is generally termed the *parameter*, or *latus rectum*.

PROPERTIES OF THE PARABOLA.

Properties deduced from the Tangent.

(26.) The equation to a straight line passing through (y', x') , (y'', x'') , is

$$y - y' = \frac{y' - y''}{x' - x''} (x - x') \dots\dots\dots (1),$$

and by the property of the parabola

$$y'^2 = 4ax' \dots\dots\dots (2),$$

$$\text{and } y''^2 = 4ax'' \dots\dots\dots (3);$$

therefore, subtracting (3) from (2),

$$(y' + y'')(y' - y'') = 4a(x' - x''),$$

$$\therefore \frac{y' - y''}{x' - x''} = \frac{4a}{y' + y''},$$

hence, by substituting (1), becomes

$$y - y' = \frac{4a}{y' + y''}(x - x').$$


Now, let the points (x', y') , (x'', y'') , approach indefinitely near to one another, then the secant joining these two points becomes a tangent, and its equation is

$$y - y' = \frac{2a}{y'}(x - x') \dots\dots\dots (4).$$

This equation may be put under a different form for multiplying by y' and transposing,

$$\begin{aligned} yy' &= 2ax - 2ax' + y'^2 \\ &= 2ax - 2ax' + 4ax' \\ &= 2a(x + x') \dots\dots\dots (5), \end{aligned}$$

an equation closely analogous to the equation of the curve, and only differing from it in this, that y' and $2x$ are replaced by yy' and $(x + x')$.



In the annexed figure, suppose P to be a point in the curve $AN = x$, $NP = y$, PT a tangent to the curve at P,

cutting the axis of x at T . Then NT is called the *subtangent*, and if PK , perpendicular to PT , be drawn cutting the axis of x at K , PK is termed a *normal*, NK the *subnormal*.

Now, from (5), when $y = 0$, $x = -x'$, therefore, $NT = 2x'$, therefore, the subtangent is equal to twice the abscissa, a very useful and important property.

Since the normal is at right angles to the tangent from (4),

$$(6), \quad y - y' = \frac{-y'}{2a} (x - x') \text{ is the equation to the normal.}$$

To find an expression for the length of the subnormal NK , put $y = 0$ in (6), $\therefore x - x' = 2a = NK$, a *constant*

$$\text{length of tangent is } \sqrt{NP^2 + NT^2} = 2 \sqrt{ax' + x'^2}, \quad (7),$$

$$\text{length of normal is } \sqrt{NP^2 + NK^2} = 2 \sqrt{ax' + a^2}, \quad (8).$$

We now proceed to the investigation of some properties of the parabola.

When $x' = a$, that is, when the tangent passes through the extremity of the parameter, the tangent and normal are equal, and also the subtangent and subnormal.

$$\text{equation of tangent is then } y - 2a = (x - a),$$

$$\therefore y = (x + a).$$

Therefore, any ordinate to the focal tangent = rad. vector of the point where the ordinate cuts the curve, so that the focal tangent cuts off from the tangent through the vertex a part equal to the distance of the vertex from the focus.

Draw the line HP perpendicular upon the directrix.

Then $SP = HP = DN = TS$, and $HTSP$ is a parallelogram,

$$\begin{aligned} \cos \angle HPT &= \frac{HP^2 + PT^2 - ST^2}{2 HP \cdot PT} = \frac{PT^2}{2 \cdot HP \cdot PT} = \frac{PT^2}{2 SP \cdot PT}, \\ &= \frac{PT}{2 SP}. \end{aligned}$$

The triangle HPS being isosceles, PT bisects the angle HPS , and it also bisects HS at right angles in Q ; the point Q is always on the axis of ordinates;

$$\therefore \angle HPT = \angle SPT,$$

therefore, the tangent at any point bisects the angle made by two straight lines, one drawn to the focus, and the other perpendicular to the directrix.

$$ST = AS + AT = a + x,$$

$$SK = SN + NK = x - a + 2a = x + a,$$

$$\therefore ST = SK, \text{ but } ST = SP,$$

$$\therefore SP = ST = SK,$$

From P draw PX' parallel to the axis


$$T'PX' = PTS = SPT,$$

$$KPx' = SPK.$$

The radius vector and the diameter at the point of contact are equally inclined to the tangent.

These are some of the principal parabolic properties, others will be referred to afterwards.

(27.) The focus S and any point P being joined, and a circle described on SP as diameter, this circle shall touch the axis AY.



Through P draw a tangent cutting axis of (y) in Y; through Y draw YC perpendicular to AY, then from equation to tangent

$$yy' = 2a(x + x'),$$

and equation to AY, $x = 0$,

$$\text{we find } AY = y = \frac{2ax'}{y'} = \frac{y'}{2}$$

(since $x'y'$ is a point in parabola),

$$\therefore BC = \frac{PN}{2}, \quad \therefore SC = CP,$$

or C is the centre of circle, and $CY = CP$, for the angle

$\text{CPY} = \text{the angle CYP}$; but CY is perpendicular to AY , therefore the circle touches AY .

(28.) Any chord Pp is drawn through S , the focus to a parabola; shew that

$$4\text{SP} \cdot \text{Sp} = \text{L}(\text{SP} + \text{Sp}),$$

L being the (latus rectum),

$$\text{SP} = \text{AN} + \text{AS},$$

$$\text{Sp} = \text{An} + \text{AS};$$

$$\text{and } \text{SP} : \text{Sp} :: \text{SN} : n\text{S}$$

$$:: (\text{AN} - \text{AS}) : (\text{AS} - \text{An})$$


$$:: (\text{SP} - 2\text{AS}) : (2\text{AS} - \text{Sp}),$$

$$\therefore \text{SP}(2\text{AS} - \text{Sp}) = \text{Sp}(\text{SP} - 2\text{AS}),$$

$$\therefore 2\text{SP} \cdot \text{Sp} = 2\text{AS}(\text{SP} + \text{Sp}).$$

$$\text{But } 4\text{AS} = \text{BB}' = \text{L},$$


$$\therefore 4\text{SP} \cdot \text{Sp} = \text{L}(\text{SP} + \text{Sp}).$$



(29.) In a plane triangle ABC , if $\tan A \tan \frac{B}{2} = 2$, and

AB be fixed, the locus of C is a parabola vertex A and focus B .

Take A for the pole, AB the initial line, $\text{AC} = r$. Find O , the centre of inscribed circle, and join OA , OB , OC , and draw perpendiculars OP , OQ , to the sides of triangle.



$$\text{Now, } r = \text{AP} + \text{PC} = \text{OP} \left\{ \cot \frac{A}{2} + \cot \frac{C}{2} \right\}$$

$$= \text{OP} \left\{ \cot \frac{A}{2} + \tan \frac{A+B}{2} \right\},$$

$$\text{and } OP = OQ = QB \tan \frac{B}{2} = (AB - AQ) \tan \frac{B}{2}$$

$$= \left(AB - OQ \cot \frac{A}{2} \right) \tan \frac{B}{2},$$

$$\therefore OQ \left(1 + \cot \frac{A}{2} \tan \frac{B}{2} \right) = AB \tan \frac{B}{2},$$

$$\therefore OP = OQ = \frac{AB \tan \frac{B}{2}}{1 + \cot \frac{A}{2} \tan \frac{B}{2}},$$

$$\therefore r = \frac{AB \tan \frac{B}{2}}{1 + \cot \frac{A}{2} \tan \frac{B}{2}} \left\{ \cot \frac{A}{2} + \frac{\tan \frac{A}{2} + \tan \frac{B}{2}}{1 - \tan \frac{A}{2} \tan \frac{B}{2}} \right\}$$

$$= \frac{AB \tan \frac{B}{2}}{1 + \cot \frac{A}{2} \tan \frac{B}{2}} \left\{ \frac{\cot \frac{A}{2} + \tan \frac{A}{2}}{1 - \tan \frac{A}{2} \tan \frac{B}{2}} \right\}.$$

$$\text{But } \tan A \cdot \tan \frac{B}{2} = 2,$$

$$\therefore \tan \frac{B}{2} = \frac{1 - \tan^2 \frac{A}{2}}{\tan \frac{A}{2}},$$

$$\therefore \tan \frac{B}{2} \cot \frac{A}{2} = \frac{1 - \tan^2 \frac{A}{2}}{\tan^2 \frac{A}{2}},$$

$$\therefore 1 + \cot \frac{A}{2} \tan \frac{B}{2} = 1 + \frac{1 - \tan^2 \frac{A}{2}}{\tan^2 \frac{A}{2}} = \cot^2 \frac{A}{2},$$

$$\text{and } 1 - \tan \frac{A}{2} \tan \frac{B}{2} = 1 - \left(1 - \tan^2 \frac{A}{2}\right) = \tan^2 \frac{A}{2},$$

$$\therefore r = AB \frac{1 - \tan^2 \frac{A}{2}}{\tan \frac{A}{2}} \left(\cot \frac{A}{2} + \tan \frac{A}{2} \right),$$

$$= AB \left(1 - \tan^2 \frac{A}{2}\right) \left(1 + \cot^2 \frac{A}{2}\right)$$

$$= AB \cdot \frac{2 \cos A}{1 + \cos A} \cdot \frac{2}{1 - \cos A}.$$

Putting for AB (c), and for A (θ), we get

$$r = \frac{4c \cdot \cos \theta}{1 - \cos^2 \theta} = \frac{4c \cdot \cos \theta}{\sin^2 \theta},$$

which is the equation to a parabola vertex A and focus B .


SECTION V.—ELLIPSE.

(30) An ellipse is a curve in which, if P be any point, $SP + PH$ is a constant quantity, S and H being two fixed points.

Bisect S, H in D , draw a perpendicular through D , and make $DB = DA = a$. With centre S , or H , and radius $= a$, cut the perpendicular in the points C and E . It is evident that A, B, C, E , are points in the curve, and that the curve is divided into four equal and similar parts by the lines CDE, ADB .

Let $DS : DA :: e : 1$, therefore, $DS = ae$. If $e = 0$, $DS = 0 = DH$, and therefore the points S and H coincide

with D and the ellipse becomes a circle. If $e = 1$, then $DS = DH = DA = DB$, and it is evident that the points of the line AB are the only ones which satisfy the required condition, since $PA + PB > AB$, whenever the point P is above the line AB. It is evident that $e > 1$ is inconsistent with the conditions of construction.



This quantity e is generally termed the eccentricity.

To find the equation of the curve, let D be the origin,

$$DN = x, NP = y, AD = a, CD = b,$$

$$SP^2 = y^2 + SN^2 = y^2 + (x - ae)^2 = y^2 + x^2 + a^2e^2 - 2aex,$$

$$HP^2 = y^2 + HN^2 = y^2 + (x + ae)^2 = y^2 + x^2 + a^2e^2 + 2aex,$$

$$\therefore HP^2 - SP^2 = (HP + SP) \cdot (HP - SP) = 4aex,$$

$$\therefore HP - SP = 2ex \dots\dots\dots (1),$$

$$HP^2 + SP^2 - 2HP \cdot PS = 4e^2x^2,$$

$$\text{and } HP^2 + SP^2 = 2(y^2 + x^2 + a^2e^2),$$

$$\therefore 2HP \cdot PS = 2y^2 + (2 - 4e^2)x^2 + 2a^2e^2,$$

$$HP^2 + SP^2 + 2HP \cdot PS = 4\{y^2 + (1 - e^2)x^2 + a^2e^2\}$$

$$\text{but } (HP + SP)^2 = 4a^2,$$

$$\therefore (1.) y^2 = (1 - e^2)(a^2 - x^2)$$

is the equation to the ellipse, the origin being at the centre ;

$$(1 - e^2) = \frac{DA^2 - DS^2}{DA^2} = \frac{DC^2}{DA^2} = \frac{b^2}{a^2}, \text{ since}$$

$$e = \frac{DS}{DA}, \text{ and } SC = DA ;$$

$$\therefore y^2 = \frac{b^2}{a^2}(a^2 - x^2), \text{ or } \frac{y^2}{b^2} + \frac{x^2}{a^2} = 1,$$

is the equation, a and b being termed respectively the semi-axis major and the semi-axis minor.

If we had taken CD as the axis of x , the equation would have been

$$\frac{x^2}{b^2} + \frac{y^2}{a^2} = 1.$$

If the origin be taken at S, $SN = x$, or $NP = y$, and for x in equation (1) substitute $(x + ae)$,

$$\begin{aligned} \therefore y^2 &= (1 - e^2)(a^2 - a^2e^2 - x^2 - 2aex) \\ &= (1 - e^2)(b^2 - x^2 - 2aex) \dots\dots\dots (2). \end{aligned}$$

If H be the origin,

$$y^2 = (1 - e^2)(b^2 + x^2 + 2aex) \dots\dots\dots (3).$$

If A be the origin,

$$y^2 = (1 - e^2)(2ax - x^2) \dots\dots\dots (4).$$

Some properties of the ellipse are directly deducible from these equations, from (1),

$$\begin{aligned} NP^2 &= \frac{CD^2}{BD^2} \cdot (BD^2 - DN^2) \\ &= \frac{CD^2}{BD^2} \cdot (BD + DN) \cdot (BD - DN) = \frac{CD^2}{BD^2} \cdot AN \cdot NB, \\ \therefore NP^2 \cdot BD^2 &= CD^2 \cdot AN \cdot NB \dots\dots\dots (A). \end{aligned}$$

$$\text{from (4), } y^2 = \frac{b^2}{a^2} \cdot (2a - x) \cdot x,$$

$$\therefore NP^2 = \frac{CD^2}{DB^2} \cdot BN \cdot AN,$$

$$\therefore NP^2 \cdot DB^2 = CD^2 \cdot AN \cdot BN \dots\dots\dots (B).$$

We may also deduce the polar equations to the ellipse.

Let D be the pole $DN = x$, $NP = y$, $DP = r$, $PDN = \theta$,

$$\therefore x = r \cos \theta, \quad y = r \sin \theta,$$

$$\therefore \text{from equation (1)} \quad r^2 \left(\frac{\sin^2 \theta}{b^2} + \frac{\cos^2 \theta}{a^2} \right) = 1,$$

$$\therefore r^2 = \frac{b^2 a^2}{a^2 \sin^2 \theta + b^2 \cos^2 \theta} = \frac{b^2 a^2}{a^2 - (a^2 - b^2) \cos^2 \theta},$$

$$\therefore r = \frac{b}{\sqrt{1 - e^2 \cos^2 \theta}}.$$

If S be the pole, $SP = r$, $HP = 2a - r$, $PSH = \theta$, then

$$(2a - r)^2 = r^2 + 4a^2 e^2 - 4aer \cos \theta,$$

$$\therefore 4a^2 + r^2 - 4ar = 4a^2 e^2 + r^2 - 4aer \cos \theta,$$

$$\therefore a^2 - ar = a^2 e^2 - aer \cos \theta,$$

$$\therefore a - r = ae^2 - er \cos \theta,$$

$$r = \frac{a(1 - e^2)}{1 - e \cos \theta}.$$

(31). We shall next proceed to the investigation of the equation to a tangent at any point P in the ellipse, since the principal properties depend upon the tangent.

Let us take a point whose co ordinates are x' , y'

Take a contiguous point whose co-ordinates are x'' , y'' .


We have, therefore, from (1),

$$y'^2 = (1 - e^2)(a^2 - x'^2)$$

$$y''^2 = (1 - e^2)(a^2 - x''^2),$$

$$\therefore y'^2 - y''^2 = -(1 - e^2)(x'^2 - x''^2),$$

$$\therefore (y' + y'')(y' - y'') = -(1 - e^2)(x' + x'')(x' - x'').$$



The equation to a line through the two points is (X and Y being the co-ordinates of any point in the line)

$$Y - y' = \frac{y' - y''}{x' - x''}(X - x') = -(1 - e^2) \frac{x' + x''}{y' + y''}(X - x').$$

Now, this line is a secant, but if we suppose the points to be taken indefinitely near one another, ultimately this line becomes a tangent.

Hence, making $x' = x''$, $y' = y''$, we obtain

$$Y - y' = - (1 - e^2) \frac{x'}{y'} (X - x')$$

as the equation to the tangent.

We must now find the values of the subtangent, normal, subnormal, &c.

$$Y - y' = - (1 - e^2) \frac{x'}{y'} (X - x')$$

the equation to the tangent.

Let $\mathbf{y} = 0$, $\therefore -y' + -(1 - e^2) \frac{x'}{y'} (\mathbf{x} - x')$,

$$\text{and } X - x' = \frac{1}{1 - e^2} \cdot \frac{y'^2}{x'}$$

$$X = \frac{y'^2 + (1 - e^2) x'^2}{(1 - e^2) x'} = \frac{(1 - e^2) a^2}{(1 - e^2) x'} = \frac{a^2}{x'} = DT,$$

$$\therefore DT = \frac{BD^2}{DN}.$$

(The student must observe, that in deducing the equation to the curve, we called DN and DP, x and y , respectively, since we were not investigating a relation between any two *particular* co-ordinates, but a relation between each pair. Here, however, P being supposed to be any *particular* point, it is most proper to denote its co-ordinates by x' and y' .)

$$\therefore DT \cdot DN = BD^2, \text{ or } DN : BD :: BD : DT.$$

$$\text{The subtangent } NT = DT - x' = \frac{a^2 - x'^2}{x'},$$

$$\therefore NT = \frac{(a + x')(a - x')}{ND} = \frac{AN \cdot NB}{ND},$$

$$\therefore NT \cdot ND = AN \cdot NB,$$

$$\text{also, } BT = \frac{a^2 - ax'}{x'} = \frac{NB \cdot DB}{ND},$$

$$\therefore BT \cdot ND = NB \cdot DB.$$

In the equation to the tangent, make $X = 0$;

$$\therefore Y - y' = - (1 - e^2) \frac{x'}{y'} (-x'), \therefore Y = \frac{y'^2 + (1 - e^2) x'^2}{y'}$$

$$= \frac{(1 - e^2) \cdot a^2}{y'} = \frac{b^2}{y'} = DG,$$

$$\therefore DG \cdot NP = DC^2;$$

$$\text{also, } CG = \frac{b^2}{y'} - b = \frac{b \cdot (b - y')}{y'} =$$

(if PV be drawn parallel to AB)

$$\frac{CV \cdot CD}{VD}, \therefore VD \cdot CG = CV \cdot CD$$

A similar relation to $BT \cdot ND = NB \cdot DB$; in fact, the two relations together may be thus expressed:—

If a tangent be drawn at any point, if, also, perpendiculars from that point be let fall upon the principal diameters, the distances between the extreme points of the diameters, and the points where the tangent meets the diameters, are to the distances between the extreme points and the feet of the perpendiculars as the diameters are to the distance between the feet of the perpendiculars and the centre.

(32) Since the normal is perpendicular to the tangent, and passes through x', y' ,

$$Y - y' = \frac{y'(X - x')}{x'(1 - e^2)} \text{ is its equation.}$$

$$\text{Making } Y = 0, \quad -y' = \frac{y'}{x'} \cdot \frac{1}{(1 - e^2)} (X - x'),$$

$$\therefore -1 = \frac{X - x'}{(1 - e^2)x'}, \text{ and } X = e^2 x' = DK,$$

$$\therefore \frac{DK}{DN} = \frac{DS^2}{DB^2}, \quad DK \cdot DB^2 = DN \cdot DS^2.$$

Again, the subnormal $NK = (1 - e^2)x'$

$$= \frac{b^2}{a^2} x' = \frac{DC^2}{DB^2} \cdot DN,$$

$$\therefore NK : DN :: DC^2 : DB^2.$$

$$\text{Making } X = 0, \quad Y - y' = \frac{y'}{x'} \cdot \frac{1}{1 - e^2} \cdot -x',$$

$$\therefore Y = \left(1 - \frac{1}{1 - e^2}\right) y' = \frac{-e^2 y'}{1 - e^2} = \frac{-DS^2 y'}{b^2},$$

$$\therefore DR = \frac{DS^2 \cdot PN}{DC^2}, \quad \therefore \frac{DR}{PN} = \frac{DS^2}{DC^2},$$

$$\begin{aligned}
\text{RE} &= b - \frac{e^2 y'}{(1 - e^2)} = b - \frac{\text{DS}^2 y'}{b^2}, \\
\therefore \text{RE} &= b - \frac{(a^2 - b^2) y'}{b^2} \\
&= b + y' - \frac{a^2 y'}{b^2} = \text{VE} - \frac{\text{BD}^2 \cdot \text{NP}}{\text{DC}^2}, \\
\therefore \text{VR} &= \frac{\text{BD}^2 \cdot \text{NP}}{\text{DC}^2}, \\
\therefore \text{VR} : \text{VD} &:: \text{BD}^2 : \text{DC}^2.
\end{aligned}$$

The equations to the tangent and normal may be put under the following form by substituting for $1 - e^2$ its value, viz., $\frac{b^2}{a^2}$,

$$\begin{aligned}
\text{Y} - y' &= -\frac{b^2}{a^2} \cdot \frac{x'}{y'} (\text{X} - x'), \text{ equation to the tangent,} \\
\therefore a^2 \text{Y} y' - a^2 y'^2 &= -b^2 \text{X} x' + b^2 x'^2, \\
\text{or } a^2 \text{Y} y' + b^2 \text{X} x' &= a^2 y'^2 + b^2 x'^2, \text{ but } a^2 y'^2 + b^2 x'^2 = a^2 b^2, \\
\therefore a^2 \text{Y} y' + b^2 \text{X} x' &= a^2 b^2;
\end{aligned}$$

$$\text{Y} - y' = \frac{a^2}{b^2} \cdot \frac{y'}{x'} (\text{X} - x'), \text{ equation to the normal.}$$

(33.) To prove that the normal bisects the angle contained by the focal distances.

Since $\text{DS} = \text{DH} = ae$, and $\text{DK} = e^2 x'$,

$$\begin{aligned}
\text{HK} &= \text{HD} + \text{DK} = ae + e^2 x' = e(a + ex'), \\
\text{SK} &= \text{DS} - \text{DK} = ae - e^2 x' = e(a - ex');
\end{aligned}$$

but, page 44, $\text{HP} + \text{PS} = 2a$, and $\text{HP} - \text{PS} = 2ex'$,

$$\therefore \text{HP} = a + ex', \text{ and } \text{PS} = a - ex';$$

hence, by substituting these values in the above, we have

$$\text{HK} = e \cdot \text{HP}, \text{ and } \text{SK} = e \cdot \text{PS};$$

$$\text{by division, } \frac{HK}{SK} = \frac{e \cdot HP}{e \cdot PS} = \frac{HP}{PS},$$

$$HK : SK :: HP : PS.$$

hence the angle HPS is bisected by the normal.

Cor. It follows from this that the angle HPG is equal to the angle SPT, that is, the tangent makes equal angles with the focal distances.

Ellipse referred to any System of Conjugates.

(31.) The characteristic property of every system of conjugate diameters is, that each bisects every chord drawn parallel to the other; it follows, that if we refer a curve to any similar system, the new equation ought only to contain the square of the co-ordinates, and a constant quantity.

The equation to the ellipse referred to its centre and principal diameter is

$$a^2 y^2 + b^2 x^2 = a^2 b^2.$$

Substituting, in this equation, for x and y , $x \cos \alpha + y \sin \alpha'$, $x \sin \alpha + y \sin \alpha'$ respectively, there results,

$$(a^2 \sin^2 \alpha' + b^2 \cos^2 \alpha') y^2 + 2 \{a^2 \sin \alpha \sin \alpha' + b^2 \cos \alpha \cos \alpha'\} xy + (a^2 \sin^2 \alpha + b^2 \cos^2 \alpha') x^2 = a^2 b^2;$$

this equation belonging to an oblique system having the same origin.

But by hypothesis the equation ought to contain only the squares of the variables and known quantities, we must therefore put the coefficient of $xy = 0$,

$$\therefore a^2 \sin \alpha \sin \alpha' + b^2 \cos \alpha \cos \alpha' = 0 \dots \dots \dots (1).$$

Hence the above equation is reduced to

$$(a^2 \sin^2 \alpha' + b^2 \cos^2 \alpha') y^2 + (a^2 \sin^2 \alpha + b^2 \cos^2 \alpha) x^2 = a^2 b^2 \dots (2)$$

Dividing equation (1) by $\cos \alpha \cos \alpha'$, we have

$$a^2 \tan \alpha \tan \alpha' + b^2 = 0, \quad \therefore \tan \alpha \tan \alpha' = -\frac{b^2}{a^2}.$$

Therefore, as we have only one equation to determine the two angles α, α' , it follows that the number of systems of conjugate diameters is infinite.

Making successively $y = 0, x = 0$, in equation (2), and there results,

$$\text{for } y = 0 \dots\dots\dots x^2 = \frac{a^2 b^2}{a^2 \sin^2 \alpha + b^2 \cos^2 \alpha};$$

$$\text{and for } x = 0 \dots\dots\dots y^2 = \frac{a^2 b^2}{a'^2 \sin^2 \alpha' + b'^2 \cos^2 \alpha'}$$


Which expressions are the squares of the semi-conjugate diameters. Calling these conjugates $2a', 2b'$, respectively, we have these relations;—

$$a'^2 = \frac{a^2 b^2}{a^2 \sin^2 \alpha + b^2 \cos^2 \alpha},$$

$$\therefore a^2 \sin^2 \alpha + b^2 \cos^2 \alpha = \frac{a^2 b^2}{a'^2},$$

$$\text{and } b'^2 = \frac{a^2 b^2}{a'^2 \sin^2 \alpha' + b'^2 \cos^2 \alpha'},$$

$$\therefore a^2 \sin^2 \alpha' + b^2 \cos^2 \alpha' = \frac{a^2 b^2}{b'^2}.$$



Substituting these values in equation (2), there results

$$a'^2 y^2 + b'^2 x^2 = a'^2 b'^2$$

for the equation to the ellipse, referred to any system of conjugate diameters.

Consequently, the centre being the origin, the equation to the ellipse preserves always the same form for any system of conjugates.

Since for $x = \pm a', y^2 = 0$, and for $y = \pm b', x^2 = 0$, it follows that the two straight lines drawn from the points $MM' mm'$, parallel to the two diameters, are tangents to the curve.

As this equation is of the same form as that between rectangular co-ordinates, it is clear that all properties which are independent of the inclination of the co-ordinates will be equally true for the conjugate diameters as for the principal diameters or axes.

The equation to the tangent is

$$a'^2 Y y'' + b'^2 X x'' = a'^2 b'^2,$$

where (x'', y'') are the co-ordinates of the point of contact.

If in this equation we make $Y = 0$; $x = \frac{a'^2}{x''}$, which is the value of OR, the point where the tangent meets the axis x ; and if from this distance we subtract x'' , or OP, we have the subtangent PR,

$$PR = \frac{a'^2}{x''} - x'' = \frac{a'^2 - x''^2}{x''};$$

this is the same result as obtained at art. 31.

Also as at page 48. The rectangle of the ordinate of the ellipse at the point of contact and the ordinate of the tangent at the centre, is equal to the square of that semi-diameter which is taken for the axis of ordinates.

Also, that the rectangle of the abscissa of the point of contact, and of the point where the tangent meets the axis of abscissæ, is equal to the square of that semi-diameter which is taken for the axis of abscissæ.

On Supplementary Chords.


(35.) If from the extremities A and B of the major axis two right lines be drawn to any point of the curve, these lines are called supplementary chords.

The equation of the right line BP, passing through the point B, whose co-ordinates are $y = 0$, and $x = a$, is

$$y = m(x - a).$$

The equation of the right line AP, passing through the point $(y = 0, x = -a)$, is

$$y = m'(x + a).$$



Now, let x', y' , be the co-ordinates of the point P; and, as this point is in both lines and also in the curve, we have the following relations:—

$$y' = m(x' - a) \dots\dots\dots (1),$$

$$y' = m'(x' + a) \dots\dots\dots (2),$$

$$a^2 y'^2 + b^2 x'^2 = a^2 b^2 \dots\dots\dots (3).$$

From (1) $m = \frac{y'}{x' - a}$, and from (2) $m' = \frac{y'}{x' + a}$,

by multiplication $mm' = \frac{y'^2}{x'^2 - a^2} \dots\dots\dots (4).$

From (3) we have $-b^2 x'^2 + a^2 b^2 = a^2 y'^2$

$$-b^2 (x'^2 - a^2) = a^2 y'^2,$$

$$\therefore \frac{y'^2}{x'^2 - a^2} = -\frac{b^2}{a^2},$$

but from (4) $mm' = \frac{y'^2}{x'^2 - a^2}$,

$$\therefore mm' = -\frac{b^2}{a^2} \dots\dots\dots (5).$$

The same relation obtains if we draw the supplementary chords from the extremities of any diameter whatever, E, E'; for let x'', y'' , be the co-ordinates of the point E, those of the point E' will be x'', y'' ; and the equation of the lines drawn from the points E, E', to any point P' whatever in the curve will be

$$y - y'' = m(x - x''),$$

$$y + y'' = m'(x + x''),$$

and, as the point P', or (x', y') , is in both these lines, we have

$$y' - y'' = m(x' - x''),$$

$$y' + y'' = m'(x' + x''),$$

$$\therefore mm' = \frac{y'^2 - y''^2}{x'^2 - x''^2};$$

but the points P' , E , E' , are also in the curve, hence

$$a^2 y'^2 + b^2 x'^2 = a^2 b^2,$$

$$a^2 y''^2 + b^2 x''^2 = a^2 b^2,$$

$$\therefore \frac{y'^2 - y''^2}{x'^2 - y''^2} = -\frac{b^2}{a^2};$$

but from above $\frac{y'^2 - y''^2}{x'^2 - y''^2} = m m',$

$$\therefore m m' = -\frac{b^2}{a^2},$$

which is the same result as before.

(36.) To find the angle contained by the supplementary chords, we have

$$\tan P = \frac{m - m'}{1 + m m'},$$

where m is the tangent of the angle PBX , and m' the tangent of the angle PAX ; we have already shown that

$$m = \frac{y'}{x' - a}, \quad m' = \frac{y'}{x' + a},$$

$$\therefore \frac{m - m'}{1 + m m'} = \frac{\frac{y'}{x' - a} - \frac{y'}{x' + a}}{1 + \frac{y'^2}{x'^2 - a^2}} = \frac{2 a y'}{x'^2 - a^2 + y'^2} \dots (6);$$

and, since the lines meet in the curve, the equation of the curve must be satisfied

$$a^2 y'^2 + b^2 x'^2 = a^2 b^2, \text{ or}$$

$$x'^2 - a^2 = -\frac{a^2 y'^2}{b^2};$$

hence, by substitution, we obtain

$$\frac{m - m'}{1 + mm'} = \frac{2ay'}{-\frac{a^2y'^2}{b^2} + y'^2} = \frac{-2ab^3}{(a^2 - b^2)y'} \dots\dots\dots (7).$$

It is clear from this result, that if we consider any point P of the curve above AB, in which case y' is positive, the tangent of the angle APB is negative; then this angle is necessarily obtuse, as it ought to be, since all the points of the ellipse are within the circumference of a circle described on AB, as a diameter.

As the obtuse angle increases its tangent decreases; hence, when the tangent is least the angle is greatest, and equation (7) will be least when y' is greatest; or when $y' = b$, this value substituted in (7) gives

$$\tan P = \frac{-2ab^2}{b(a^2 - b^2)} = \frac{-2ab}{a^2 - b^2}.$$

On this supposition, the values of m and m' become

$$m = \frac{-b}{a}, \quad m' = \frac{b}{a}, \quad \text{since } y' = b, \text{ gives } x = 0.$$

From the above, it appears that supplementary chords drawn from the extremities of the major axis make the greatest angle when they meet in the point C, the extremity of the minor axis.

(37.) We can deduce some other useful properties by transforming the oblique axes to rectangular axes,

$$x = \frac{x \sin \alpha' - y \cos \alpha'}{\sin (\alpha' - \alpha)},$$

$$y = \frac{y \cos \alpha - x \sin \alpha}{\sin (\alpha' - \alpha)}.$$

Substituting the values in equation (52) page 74, we have

$$(a'^2 \cos^2 \alpha + b'^2 \cos^2 \alpha) y^2 + (-2a' \sin \alpha \cos \alpha - 2b' \sin \alpha' \cos \alpha') xy$$

$$+ (a'^2 \sin^2 \alpha + b'^2 \sin^2 \alpha) x^2 = a'^2 b'^2 \sin^2 (\alpha' - \alpha),$$

$$\text{which must be the same as } a^2 y^2 + b^2 x^2 = a^2 b^2,$$

therefore, the following conditions must have place,

$$a'^2 \cos^2 \alpha + b'^2 \cos^2 \alpha' = a^2 \dots\dots\dots (a),$$

$$a'^2 \sin^2 \alpha + b'^2 \sin^2 \alpha' = b^2 \dots\dots\dots (b),$$

$$a'^2 \sin \alpha \cos \alpha + b'^2 \sin \alpha' \cos \alpha' = 0 \dots\dots\dots (c),$$

$$a'^2 b'^2 \sin^2 (\alpha' - \alpha) = a^2 b^2 \dots\dots (d);$$

add (a) and (b), and there results

$$a'^2 + b'^2 = a^2 + b^2 \dots\dots\dots (e);$$

or the sum of the squares of any system of conjugate diameters is equal to the sum of the squares of the principal diameters.

Equation (d), multiplied by 4, gives

$$4 a' b' \sin (\alpha' - \alpha) = 4 a b,$$

and since $\alpha' - \alpha$ is the inclination of the conjugate axes, the left-hand side of the equation expresses the area of a parallelogram whose sides are those axes, and included angle $\alpha' - \alpha$ (see Hann's "Trigonometry," page 60); the right-hand side gives the rectangle of principal axes; hence, every parallelogram circumscribing an ellipse having the sides parallel to a system of conjugate axes, is equal to the rectangle of the principal axes.

Now, since the area of the parallelogram is

$$4 \text{ MD} \cdot m i,$$

$$\text{we have } 4 \text{ MD} \cdot m i = 4 a b,$$

$$m i = \frac{a b}{\text{MD}},$$

but $m i$ equals a perpendicular on the tangent Tt from D , which call p , then

$$p = \frac{a b}{\text{MD}}, \quad \text{or } p^2 = \frac{a^2 b^2}{\text{MD}^2} = \frac{a^2 b^2}{a^2 + b^2 - a'^2}$$

$$(\text{since } \text{MD}^2 + m D^2 = a^2 + b^2).$$

$$\begin{aligned}
\text{Now } mD^2 &= a^2 + b^2 - DM^2 \\
&= a^2 + b^2 - (x^2 + y^2) \\
&= a^2 + b^2 - \left\{ x^2 + \frac{b^2}{a^2} (a^2 - x^2) \right\} \\
&= a^2 + b^2 - x^2 - b^2 + \frac{b^2}{a^2} x^2 \\
&= a^2 - \left(1 - \frac{b^2}{a^2} \right) x^2 \\
&= a^2 - \left(\frac{a^2 - b^2}{a^2} \right) x^2 \\
&= a^2 - e^2 x^2 \\
&= (a + ex)(a - ex);
\end{aligned}$$

but by art. 33, $a + ex = HP$, and $a - ex = SP$, fig. p. 47.

$$\therefore mD^2 = SP \cdot HP,$$

or the rectangle contained by the focal distances of any point of an ellipse, is equal to the square of the corresponding semi-conjugate diameter.

(38.) In the ellipse

$$p^2 = b^2 \cdot \frac{r}{2a - r},$$

where p is the perpendicular from one of the foci on the tangent: the equation to the tangent is

$$y = \frac{-b^2 x}{a^2 y'} x + \frac{b^2}{y'},$$

perpendicular from the point $(-ae, 0)$ on this

$$\begin{aligned}
&= \frac{\frac{-b^2 x'}{a^2 y'} - ae - \frac{b^2}{y'}}{\sqrt{1 + \frac{b^4 x'^2}{a^4 y'^2}}}
\end{aligned}$$

$$\begin{aligned}
&= \frac{-b^2 \{ae x' + a^2\}}{\sqrt{a^4 y'^2 + b^4 x'^2}} \\
&= \frac{-b^2 a (a + e x')}{\sqrt{a^2 b^2 \left\{ a^2 - x^2 + \frac{b^2}{a^2} x'^2 \right\}}} \\
&= \frac{-b (a + e x')}{\sqrt{a + a x'} \sqrt{a - e x'}} \\
&= \frac{-b \sqrt{a + e x'}}{\sqrt{a - e x'}}, \\
\therefore p^2 &= \frac{b^2 (a + e x')}{a - e x'} \\
&= b^2 \cdot \frac{\text{HP}}{\text{SP}}, \\
\therefore p^2 &= b^2 \cdot \frac{r}{2a - r}.
\end{aligned}$$

In the same manner, if we put p' for the perpendicular from the other focus on the tangent, we have

$$\begin{aligned}
p'^2 &= b^2 \frac{\text{SP}}{\text{HP}}, \\
\therefore p^2 \cdot p'^2 &= b^2 \cdot \frac{\text{SP}}{\text{HP}} \cdot b^2 \frac{\text{HP}}{\text{SP}} = b^4, \\
p \cdot p' &= b^2.
\end{aligned}$$

That is, the rectangle of the perpendiculars from the foci on the tangent is equal to the square of the semi-axis minor.

(39) If a circle be described on the major axis of an ellipse, and if Y be the ordinate to the circle, and y the ordinate to the ellipse corresponding to the same abscissa, then

$$\frac{y}{Y} = \frac{b}{a}.$$

By the equation to the circle

$$Y^2 = a^2 - x^2,$$

$$\text{ellipse } y^2 = \frac{b^2}{a^2} (a^2 - x^2).$$

Divide the second by the first, and we have

$$\frac{y^2}{Y^2} = \frac{b^2}{a^2}, \quad \text{or } \frac{y}{Y} = \frac{b}{a};$$

that is, the ordinate of an ellipse is to the ordinate of a circle described upon the major axis, as the minor axis is to the major axis.

It is evident that this gives an easy method of describing an ellipse by points.

(40.) If xy , and $x'y'$ be any two points on the conjugate diameters, then

$$\frac{xx'}{a^2} + \frac{yy'}{b^2} = 0,$$

$$\text{for } \frac{y}{x} = \tan \alpha, \quad \text{and } \frac{y'}{x'} = \tan \alpha',$$

$$\text{but } \tan \alpha \tan \alpha' = -\frac{b^2}{a^2},$$

$$\therefore \frac{y}{x} \cdot \frac{y'}{x'} = -\frac{b^2}{a^2}$$

$$\frac{yy'}{b^2} = -\frac{xx'}{a^2},$$

$$\text{or } \frac{xx'}{a^2} + \frac{yy'}{b^2} = 0.$$

On the Eccentric Angle in the Ellipse.*


(41.) The co-ordinates of any point xy of an ellipse, may take the form $x = a \cos \phi$, $y = b \sin \phi$.

* O'Brien's "Co-ordinate Geometry," page 111.

If we assume $\frac{x}{a} = \cos \phi$, or, in other words, $\phi = \cos^{-1} \frac{x}{a}$, then, substituting in the equation,

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1,$$

$$\text{we have } \frac{y^2}{b^2} = 1 - \cos^2 \phi = \sin^2 \phi,$$



$$\therefore y^2 = b^2 \sin^2 \phi, \text{ or } y = b \sin \phi.$$

Hence, we may assume $x = a \cos \phi$, and $y = b \sin \phi$.

If we describe a circle, $AP'B$ on AB , as a diameter, produce the ordinate MP to meet it in P' , and join $P'D$; $P'DM$ is the angle whose cosine is $\frac{x}{a}$, for

$$\cos P'DM = \frac{DM}{DP'} = \frac{DM}{DB} = \frac{x}{a};$$

therefore, if we produce the ordinate to meet a circle described on the major axis as a diameter, and draw a line from the point where the ordinate meets the circle to the centre, then ϕ is the angle which that line makes with the major axis.

If PQR be drawn parallel to $P'D$, meeting CD' in the point R , and DB in Q , then $PR = P'D = a$, and $PQB = P'DB = \phi$; since $y = b \sin \phi$, we have

$$b \sin \phi = PM = PQ \sin PQM = PQ \sin \phi,$$

$$\therefore PQ = b.$$

If ϕ and ϕ' be the eccentric angles of the extremities of two conjugate diameters, then $\phi' = \phi + 90^\circ$.

Let (x, y) , and (x', y') , be the co-ordinates of the extremities, P and C' of the two conjugate diameters DP and DC', ϕ and ϕ' the corresponding eccentric angles,

$$\text{then } x = a \cos \phi, \quad y = b \sin \phi, \quad x' = a \cos \phi', \quad y' = b \sin \phi',$$

$$\text{then, by art. 40, } \cos \phi \cos \phi' + \sin \phi \sin \phi = 0,$$

$$\text{or } \cos (\phi' - \phi) = 0,$$

$$\text{which shews that } \phi' - \phi = 90^\circ, \text{ or } \phi' = \phi + 90^\circ.$$

This property of conjugate diameters, with reference to the eccentric angle, is of great use in problems relating to conjugate diameters.

COR. Hence

$$x' = a \cos (\phi + 90^\circ) = -a \sin \phi = -\frac{a}{b} y,$$

$$\text{and } y' = b \sin (\phi + 90^\circ) = b \cos \phi = \frac{b}{a} x,$$

which determine the co-ordinates of P and C.

(42.) The properties deduced in art. 37 can be found in a very simple manner, by using the eccentric angle.

$$\text{Let } DP = r, \quad C'D = r', \quad PDB = \theta, \quad C'DB = \theta',$$

$$r^2 = x^2 + y^2 = a^2 \cos^2 \phi + b^2 \sin^2 \phi \quad \dots\dots\dots (1);$$

and putting $\phi + 90$ for ϕ , we have

$$r'^2 = a^2 \sin^2 \phi + b^2 \cos^2 \phi \quad \dots\dots\dots (2);$$

add (1) and (2),

$$\begin{aligned} r^2 + r'^2 &= a^2 (\sin^2 \phi + \cos^2 \phi) + b^2 (\sin^2 \phi + \cos^2 \phi) \\ &= a^2 + b^2 \quad \dots\dots\dots (4). \end{aligned}$$

$$\begin{aligned} \text{The area of a parallelogram, whose sides are } C'D \text{ and } PD, \\ = rr' \sin (\theta' - \theta) = rr' (\sin \theta' \cos \theta - \cos \theta' \sin \theta) \end{aligned}$$

$$\begin{aligned}
&= r \cos \theta . r' \sin \theta' - r \sin \theta . r' \cos \theta \\
&= xy' - x'y \\
&= ab \sin \phi' \cos \phi - ab \cos \phi' \sin \phi \\
&= ab (\sin \phi' \cos \phi - \cos \phi' \sin \phi) \\
&= ab \sin(\phi' - \phi) = ab \{\text{since } \sin(\phi' - \phi) = \sin 90\} = 1.
\end{aligned}$$

(43.) Given the axes of an ellipse and the inclination of any two conjugate diameters, to determine their length and direction.

In the first place, we shall find a' and b' . By page 57,

$$a'^2 + b'^2 = a^2 + b^2; \quad 2a'b' = \frac{2ab}{\sin \phi};$$

by adding and subtracting these two equations, we have

$$(a' + b')^2 = a^2 + b^2 + \frac{2ab}{\sin \phi},$$

$$(a' - b')^2 = a^2 + b^2 - \frac{2ab}{\sin \phi};$$

$$\therefore a' = \frac{1}{2} \sqrt{a^2 + b^2 + \frac{2ab}{\sin \phi}} + \frac{1}{2} \sqrt{a^2 + b^2 - \frac{2ab}{\sin \phi}},$$

$$b' = \frac{1}{2} \sqrt{a^2 + b^2 + \frac{2ab}{\sin \phi}} - \frac{1}{2} \sqrt{a^2 + b^2 - \frac{2ab}{\sin \phi}}.$$

In the equation, page 51, $a^2 \tan \alpha \tan \alpha' + b^2 = 0$,

put $\phi + \alpha$ for α' , and we have

$$a^2 \tan \alpha \tan (\phi + \alpha) + b^2 = 0,$$

$$\text{but } \tan (\phi + \alpha) = \frac{\tan \phi + \tan \alpha}{1 - \tan \phi \tan \alpha};$$

therefore, by substitution and reduction we have the quadratic

$$a^2 \tan^2 \alpha + (a^2 - b^2) \tan \phi \cdot \tan \alpha + b^2 = 0,$$

$$\therefore \tan \alpha = -\frac{(a^2 - b^2) \tan \epsilon}{2a^2} \pm \frac{1}{2a^2} \sqrt{(a^2 - b^2)^2 \tan^2 \epsilon - 4a^2 b^2},$$

this value of $\tan \alpha$ will be real if $(a^2 - b^2)^2 \tan^2 \epsilon$ be greater or at least equal to $4a^2 b^2$.

$$\text{If } \tan^2 \epsilon = \frac{4a^2 b^2}{(a^2 - b^2)^2}, \text{ or } \tan \epsilon = \pm \frac{2ab}{a^2 - b^2};$$

substituting this value in the $\tan \alpha$, we have

$$\tan \alpha = -\frac{a^2 - b^2}{2a^2} \times \pm \frac{2ab}{a^2 - b^2} = \pm \frac{b}{a},$$

which, since $\tan \alpha \tan \alpha' = -\frac{b^2}{a^2}$, gives

$$\tan \alpha = -\frac{b}{a}, \quad \tan \alpha' = \frac{b}{a}.$$

(44.) Given two conjugate diameters in an ellipse and their inclination to determine the axes.

Since $a^2 + b^2 = a'^2 + b'^2$, and $2ab = 2a'b' \sin \epsilon$;

by adding and subtracting these equations, we have

$$(a + b)^2 = a'^2 + b'^2 + 2a'b' \sin \epsilon,$$

$$(a - b)^2 = a'^2 + b'^2 - 2a'b' \sin \epsilon,$$

$$a = \frac{1}{2} \sqrt{a'^2 + b'^2 + 2a'b' \sin \epsilon} + \frac{1}{2} \sqrt{a'^2 + b'^2 - 2a'b' \sin \epsilon},$$

$$b = \frac{1}{2} \sqrt{a'^2 + b'^2 + 2a'b' \sin \epsilon} - \frac{1}{2} \sqrt{a'^2 + b'^2 - 2a'b' \sin \epsilon}.$$

These values are always real, for $(a' - b')^2 > 0$, then $a'^2 + b'^2 > 2a'b'$, and, *a fortiori*, $a'^2 + b'^2 > 2a'b' \sin \epsilon$. We can determine the angle α from the equation

$$\tan \alpha \cdot \tan (\epsilon + \alpha) = -\frac{b^2}{a^2},$$

as before, in the last article.

(45.) *The Normal in Terms of its Inclination to the Major Axis.*

The equation to the normal is

$$\begin{aligned} y - y' &= \frac{a^2 y'}{b^2 x'} (x - x'), \\ \text{or } y &= \frac{a^2 y'}{b^2 x'} x + y' - \frac{a^2 y'}{b^2} \\ &= \frac{a^2 y'}{b^2 x'} x + \frac{b^2 - a^2}{b^2} \cdot y' \dots\dots (1). \end{aligned}$$

Let the equation to the normal be

$$y = mx + h \dots\dots\dots (2),$$

then, comparing (1) and (2),

$$m = \frac{a^2 y'}{b^2 x'}, \text{ and } h = \frac{b^2 - a^2}{b^2} \cdot y' \dots\dots (3),$$

$$m = \frac{a^2}{b^2} \cdot \frac{b}{a} \cdot \frac{\sqrt{a^2 - x'^2}}{x'}$$

$$\frac{b^2 m^2}{a^2} = \frac{a^2 - x'^2}{x'^2},$$

$$\therefore x'^2 = \frac{a^4}{a^2 + m^2 b^2},$$

$$\begin{aligned} \therefore a^2 - x'^2 &= a^2 - \frac{a^4}{a^2 + m^2 b^2} \\ &= \frac{a^4 + a^2 m^2 b^2 - a^4}{a^2 + m^2 b^2}, \end{aligned}$$

$$\therefore \sqrt{a^2 - x'^2} = \frac{a m b}{\sqrt{a^2 + m^2 b^2}},$$

$$\begin{aligned}
 \therefore h &= \frac{b^2 - a^2}{b^2} \cdot \frac{b}{a} \sqrt{a^2 - x'^2} \\
 &= \frac{b^2 - a^2}{ab} \cdot \frac{am b}{\sqrt{a^2 + m^2 b^2}} \\
 &= - (a^2 - b^2) \cdot \frac{m}{\sqrt{a^2 + m^2 b^2}};
 \end{aligned}$$

$$\therefore (2) \text{ becomes } y = mx - (a^2 - b^2) \cdot \frac{m}{\sqrt{a^2 + m^2 b^2}},$$

$$\text{or } (y - mx) \sqrt{a^2 + m^2 b^2} + m \cdot (a^2 - b^2) = 0.$$

The equation to the tangent in terms of the angle that it makes with the major axis, is

$$y = mx \pm \sqrt{m^2 a^2 + b^2}.$$

(46.) In the ellipse, to find the locus of the middle points of a series of parallel chords.

The equation to the ellipse is

$$A^2 y^2 + B^2 x^2 = A^2 B^2 \dots\dots\dots (1).$$

Let MM' be any right line whose equation is

$$y = mx + h \dots\dots\dots (2).$$

Substitute in (1) for y its value from (2),

$$(A^2 m^2 + B^2) x^2 - 2 A^2 m h \cdot x + A^2 h^2 - A^2 B^2 = 0,$$

the roots of this equation will give the abscissæ of the two points MM',

$$x^2 - \frac{2 A^2 m h \cdot x}{A^2 m^2 + B^2} + \frac{A^2 h^2 - A^2 B^2}{A^2 m^2 + B^2} = 0;$$

let $x'y'$, and $x''y''$, be the co-ordinates of the two points, and α, \mathfrak{C} , those of the middle point N,

$$\text{then } \alpha = \frac{x' + x''}{2}, \quad \mathfrak{C} = \frac{y' + y''}{2},$$

since in a quadratic equation the coefficient of the second term taken with a contrary sign is equal to the sum of the roots, we have

$$x' + x'' = -\frac{2A^2mh}{A^2m^2 + B^2},$$

$$\therefore \alpha = -\frac{A^2mh}{A^2m^2 + B^2}.$$

To find the corresponding value of \mathfrak{C} , we must remark that α and \mathfrak{C} is a point in the line whose equation is

$$y = mx + h;$$

hence, substituting \mathfrak{C} for y , and α for x , we obtain

$$\mathfrak{C} = m\alpha + h;$$

substitute the value of α in this equation

$$\mathfrak{C} = -\frac{A^2m^2h}{A^2m^2 + B^2} + h = \frac{+B^2h}{A^2m^2 + B^2}.$$

Dividing \mathfrak{C} by α , we get

$$\frac{\mathfrak{C}}{\alpha} = \frac{-B^2}{A^2m}.$$

Now, as this result is independent of h , which fixes the position of the chord MM' , it follows that if we put α', \mathfrak{C}' , α'', \mathfrak{C}'' , the co-ordinates of the middle points of the other chords parallel to the first, we shall in like manner find

$$\frac{\mathfrak{C}'}{\alpha'} = \frac{-B^2}{A^2m^2}, \quad \frac{\mathfrak{C}''}{\alpha''} = \frac{-B^2}{A^2m^2},$$

therefore, in general, representing the co-ordinates of the middle points by x and y , we have the relation

$$\frac{y}{x} = \frac{-B^2}{A^2 m'}, \text{ or } y = -\frac{B^2}{A^2 m'} \cdot x,$$

which is evidently a straight line passing through the origin.


Hence, all the diameters of an ellipse pass through the centre; and, conversely, every line which passes through the centre is a diameter, if we put m' for the tangent of the angle which the line forms with the axis of x ,

$$m' = \frac{-B^2}{A^2 m}, \quad \therefore m = \frac{-B^2}{A^2 m'}.$$

The same process applies to the hyperbola, by changing $-B^2$ into $+B^2$.

SECTION VI.—HYPERBOLA.

(47.) If S and H be two fixed points, and if the difference of HP and SP be a constant quantity $= 2a$ suppose, the locus of P is the hyperbola.



Bisect SH in D , make

$$DB = DA = a;$$

then, A and B are manifestly points in the curve.

Let $\frac{DH}{DB} = e$, a quantity greater than unity; then $DH = ae$.

To find the equation to the curve.

Let D be the origin $DN = x$, $NP = y$;

$$\text{then } HN = ae + x, \quad SN = ae - x,$$

$$HP^2 = y^2 + (ae + x)^2, \quad SP^2 = y^2 + (ae - x)^2,$$

$$\therefore HP^2 + SP^2 = 2y^2 + 2a^2e^2 + 2x^2,$$

$$HP^2 - SP^2 = (HP - SP)(HP + SP) = 4aex,$$

$$\therefore \text{HP} + \text{SP} = 2cx, \text{ and } \text{HP}^2 + \text{SP}^2 + 2\text{HP} \cdot \text{PS} = 4e^2x^2,$$

$$\therefore 2\text{HP} \cdot \text{PS} = (4e^2 - 2)x^2 - 2y^2 - 2a^2e^2,$$

$$\text{and } \text{HP}^2 + \text{SP}^2 - 2\text{HP} \cdot \text{PS} = 4a^2 = 4(y^2 + a^2e^2) + 4(1 - e^2)x^2;$$

herefore the equation to the hyperbola, the point D being the origin, is

$$y^2 = (1 - e^2)(a^2 - x^2).$$

This expression, $1 - e^2$, is generally put $= -\frac{b^2}{a^2}$, and then

$$\frac{y^2}{b^2} - \frac{x^2}{a^2} = -1, \text{ or } a^2y^2 - b^2x^2 = -a^2b^2 \dots (1),$$

is the hyperbolic equation.

If we had taken the point A for the origin, y has the same value as in (1), but AN is $= \text{DN} - \text{DA} = \text{DN} - a$;

$$\therefore \frac{y^2}{b^2} - \frac{(x + a)^2}{a^2} = -1,$$

$$\text{or } y^2 = (1 - e^2)(-2ax - x^2) = (e^2 - 1)(2ax + x^2) \dots (2),$$

and if we take S the origin, $\text{SN} = \text{DN} - ae$,

$$\therefore y^2 = (e^2 - 1)(2aex + x^2) \dots \dots \dots (3).$$

(48.) We now proceed to the deduction of the polar equations to the hyperbola. If D be the pole, $\text{DP} = r$, $\text{HDP} = \theta$, then, in equation (1), $x = -r \cos \theta$, $y = r \sin \theta$; and we have

$$r^2 \sin^2 \theta = (1 - e^2)(a^2 - r^2 \cos^2 \theta),$$

$$\therefore r^2 \{\sin^2 \theta + (1 - e^2) \cos^2 \theta\} = a^2 \cdot (1 - e^2),$$

$$\text{and } r^2 = \frac{a^2(1 - e^2)}{1 - e^2 \cos^2 \theta},$$

$$\therefore r = \frac{a \sqrt{e^2 - 1}}{\sqrt{e^2 \cos^2 \theta - 1}} = \frac{b}{\sqrt{e^2 \cos^2 \theta - 1}};$$

and if H be the pole, $\angle \text{PHS} = \theta$, $\text{HP} = r$,

$$\text{SP}^2 = (2a - r)^2 = r^2 + 4a^2e^2 - 4aer \cos \theta,$$

$$\therefore 4a^2 + r^2 - 4ar = r^2 + 4a^2e^2 - 4aer \cos \theta,$$

$$\therefore a^2 - ar = a^2e^2 - aer \cos \theta, \therefore r = \frac{a(1 - e^2)}{1 - e \cos \theta}.$$

Now, it is evident that there exists a curve similarly situated with regard to H, as AP with regard to S; these two curves are called opposite hyperbolæ, though, properly speaking, the *hyperbola* is the system compounded of the two.

(49.) We now proceed to the discussion of a few hyperbolic properties; but it will be convenient to first investigate the equation to the tangent, since the most important properties depend upon it.

Let (y', x') , (y'', x'') , be two points in the curve,

$$\text{then, } y'^2 = (1 - e^2)(a^2 - x'^2), \quad y''^2 = (1 - e^2)(a^2 - x''^2),$$

$$\therefore y'^2 - y''^2 = (y' + y'')(y' - y'') = (1 - e^2)(x''^2 - x'^2) \\ = (e^2 - 1)(x' + x'')(x' - x''),$$


and the equation to a straight line passing through (y', x') , (y'', x'') , is, if Y and X be the co-ordinates of the line,

$$Y - y' = \frac{y' - y''}{x' - x''}(X - x'),$$

$$\text{or } Y - y' = \frac{x' + x''}{y' + y''} \cdot (e^2 - 1)(X - x').$$

Now, supposing the points (y', x') , (y'', x'') , to approach continually, the line ultimately becomes a tangent at the point (y', x') , and its equation is then

$$Y - y' = \frac{x'}{y'} \cdot (e^2 - 1) \cdot (X - x').$$



We have now to find the subtangent, subnormal, &c.

$$Y - y' = \frac{(e^2 - 1)x'}{y'} \cdot (X - x'),$$

the equation to the tangent at the point (y', x') .

$$\text{If } X = 0, \quad Y - y' = \frac{(1 - e^2)x'^2}{y'},$$

$$\therefore Y = \frac{y'^2 + (1 - e^2)x'^2}{y'} = DT' = \frac{(1 - e^2)a^2}{y'} = -\frac{b^2}{y'},$$

this is the point where the tangent meets the conjugate axes below D,

$$\therefore DT' : DC :: DC : PN.$$

Also, if $Y = 0$,

$$-y' = \frac{(e^2 - 1)x'}{y'} (X - x'), \quad \therefore X - x' = \frac{y'^2}{(1 - e^2)x'},$$

$$X = \frac{(1 - e^2)x'^2 + y'^2}{(1 - e^2)x'} = \frac{(1 - e^2)a^2}{(1 - e^2)x'} = \frac{a^2}{x'} = TD,$$

$$\therefore TD : DA :: DA : DN,$$

$$DN - DT = TN = \frac{x'^2 - a^2}{x'} = \text{subtangent},$$

another property of the hyperbola.

(50.) The equation to the normal PK now remains to be found.

$Y - y' = \frac{(e^2 - 1)x'}{y'} (X - x')$, is the equation to the tangent; since the normal is perpendicular to the tangent and passes also through y', x' , its equation consequently is

$$Y - y' = \frac{y'}{(1 - e^2)x'} (X - x').$$

If in this equation we make $X = 0$,

$$Y - y' = \frac{y'}{(1 - e^2)x'} (-x'),$$

$$TN = \frac{DN^2 - AD^2}{DN} \dots\dots\dots (3),$$

$$\frac{DG}{NP} = \frac{DS^2}{BS \cdot AS}, \therefore DG : (NP \cdot DS) :: DS : (BS \cdot AS) \dots (4),$$

$$NK = \frac{CD^2}{AD^2} \cdot DN \dots\dots\dots (5);$$

or, which is the same,

$$DG = \frac{AD^2 + CD^2}{CD^2} \cdot PN.$$

On Directrices and Parameters.

The hyperbola and ellipse have each two directrices, but the parabola has only one.

In the parabola the distance of the directrix from the vertex is $= a$.

In the ellipse and hyperbola the distance of the directrix from the centre $= \frac{a}{e}$.

The parameter, or latus rectum, is the double ordinate drawn through the focus in each of the three curves. In the parabola it is $= 4a$, a being the distance between the vertex and focus. In the ellipse and hyperbola it is $= \frac{2b^2}{a}$, when a and b are the major and minor axes.

(51.) *On the Asymptote to the Hyperbola.*

$$y^2 = (1 - e^2)(a^2 - x^2),$$

$$\therefore y = \sqrt{e^2 - 1} \cdot \left\{ x - \frac{a^2}{2 \cdot x} \right\} \&c. \dots\dots\dots (1),$$

is the equation to the curve.

From the form of this equation, it seems that there are relations between the curve and the line

$$y = (\sqrt{e^2 - 1})x, \text{ which are worth attending to.}$$

It is evident that this line passes through the origin, making an angle $= \tan^{-1} \sqrt{e^2 - 1}$ with the principal diameter;

$$\begin{aligned} \text{and since } y^2 &= (e^2 - 1)x^2 \text{ in the line,} \\ \text{and } y^2 &= (e^2 - 1)x^2 - (e^2 - 1)\frac{a^2}{x^2} \text{ in the curve,} \end{aligned}$$

it is evident that the lineal ordinate is greater than the hyperbolic corresponding to the same abscissa.

But on considering the equation to the curve when expanded into the series (1) above, it is evident that as x increases the hyperbolic and lineal ordinate approximate nearer and nearer.

So that we may say, that when $x = a$ the line touches the hyperbola, and is therefore an asymptote.

Since in the equation $y = \sqrt{e^2 - 1} \cdot x$, the radical may be taken with either sign, and x may be either positive or negative; there are evidently four asymptotes, all cutting at D, a pair belonging to each hyperbolic branch.

It will be an instructive exercise for the student to endeavour to deduce the asymptotes from the tangential equation

$$Y - y' = \frac{(e^2 - 1) \cdot x'}{y'} (X - x'),$$

considering the asymptote to be a tangent at an infinite distance.

(52) To prove that the normal bisects the exterior angle considering the focal distances at any point in the hyperbola.

$$\begin{aligned} \text{By art. 47,} \quad \text{HP} + \text{SP} &= 2ex, \\ \text{and } \text{HP} - \text{SP} &= 2a; \end{aligned}$$

by adding and subtracting we have

$$\text{HP} = ex + a, \text{ and } \text{SP} = ex - a,$$

$$\text{SK} = \text{DK} - \text{DS} = e^2 x' - ae = e(ex' - a) = e \cdot \text{SP},$$

$$\text{HK} = \text{DK} + \text{DH} = e^2 x + ae = e(ex' + a) = e \cdot \text{HP},$$

$$\therefore \frac{\text{SK}}{\text{HK}} = \frac{\text{SP}}{\text{HP}},$$

$$\therefore \text{PK bisects the angle } \text{SPP}'.$$

From this it is clear that the focal distances make equal angles with the tangent, that is, the angle SPT is equal to the angle HPT ; for $\text{KPT} = \text{KPT}'$, being right angles, and $\text{SPK} = \text{KPP}'$, from above; the latter, being taken from the former, leaves the remaining angles SPT and $\text{P'PT}''$ equal, but $\text{P'PT}''$ is equal to its vertical opposite angle HPT ,

$$\therefore \text{HPT} = \text{SPT}.$$

(53.) By proceeding in the same manner as for the ellipse, page 58, we have $p^2 = b^2 \frac{r}{2a + r}$, and therefore we have in the same manner

$$pp' = b^2,$$

or the rectangle of the perpendiculars from the foci on the tangent is equal to the square of the semi-axis minor in the hyperbola as well as in the ellipse.

(54.) When the hyperbola is related to its conjugate diameters, the process is exactly the same as for the ellipse.

The equation referred to its axis is $a^2 y^2 - b^2 x^2 = -a^2 b^2$, and substituting for x and y , as in the ellipse, we get

$$a^2 \sin a \sin a' - b^2 \cos a \cos a' = 0 \dots \dots \dots (1),$$

$$\text{and } (a^2 \sin^2 a' - b^2 \cos^2 a') y^2$$

$$+ (a^2 \sin^2 a - b^2 \cos^2 a) x^2 = -a^2 b^2 \dots \dots \dots (2),$$

from (1) we have, dividing by $\cos a \cos a'$,

$$a^2 \tan a \tan a' - b^2 = 0, \text{ or } \tan a \tan a' = \frac{b^2}{a^2} \dots \dots \dots (3),$$

$$\tan a' = \frac{b^2}{a^2 \tan a} \dots \dots \dots (4).$$

In equation (2) make $y = 0$ and $x = 0$ successively, and we have

$$x^2 = \frac{-a^2 b^2}{a^2 \sin^2 a - b^2 \cos^2 a}$$

$$y^2 = \frac{-a^2 b^2}{a^2 \sin^2 a' - b^2 \cos^2 a'}.$$

We must observe that, of these two squares, one is positive, and the other is negative, for any value of x corresponding to $y = 0$ is real; but that of y corresponding to $x = 0$ is imaginary; but that they have different signs may be shewn as follows:—multiplying these two equations together, we have

$$x^2 y^2 = \frac{a^4 b^4}{(a^2 \sin^2 a - b^2 \cos^2 a)(a^2 \sin^2 a' - b^2 \cos^2 a')},$$

E 2

multiplying out the denominator

$$a^4 \sin^2 \alpha \sin^2 \alpha' + b^4 \cos^2 \alpha \cos^2 \alpha' \\ - a^2 b^2 (\sin^2 \alpha' \cos^2 \alpha + \sin^2 \alpha \cos^2 \alpha');$$

but by equation (1), $a^2 \sin \alpha \sin \alpha' - b^2 \cos \alpha \cos \alpha' = 0$,

$$\therefore a^4 \sin^2 \alpha \sin^2 \alpha' + b^4 \cos^2 \alpha \cos^2 \alpha' = 2 a^2 b^2 \sin \alpha \sin \alpha' \cos \alpha \cos \alpha';$$

therefore, the denominator becomes

$$a^2 b^2 (2 \sin \alpha \sin \alpha' \cos \alpha \cos \alpha' - \sin^2 \alpha' \cos^2 \alpha - \sin^2 \alpha \cos^2 \alpha') \\ = -a^2 b^2 (\sin \alpha' \cos \alpha - \sin \alpha \cos \alpha')^2 = -a^2 b^2 \sin^2 (\alpha' - \alpha);$$

$$\therefore x^2 y^2 = \frac{a^4 b^4}{-a^2 b^2 \sin^2 (\alpha' - \alpha)} = -\frac{a^2 b^2}{\sin^2 (\alpha' - \alpha)} \dots\dots (5).$$

This result, being essentially negative, proves that x^2 and y^2 have different signs.

Taking the axis of x for the transverse diameter in this case, the expression for x^2 is positive, but that of y^2 is negative; putting a'^2 for the first, and $-b'^2$ for the second, we have

$$a'^2 = \frac{-a^2 b^2}{a^2 \sin^2 \alpha - b^2 \cos^2 \alpha}$$

$$-b'^2 = \frac{-a^2 b^2}{a^2 \sin^2 \alpha' - b^2 \cos^2 \alpha'},$$

$$\therefore a^2 \sin^2 \alpha - b^2 \cos^2 \alpha = -\frac{a^2 b^2}{a'^2}$$

$$a^2 \sin^2 \alpha' - b^2 \cos^2 \alpha' = \frac{a^2 b^2}{b'^2};$$

from these values of a'^2 and $-b'^2$, equation (2) becomes

$$a^2 y^2 - b'^2 x^2 = -a'^2 b'^2,$$

the equation to the hyperbola related to oblique axes originating at the centre, and making angles α and α' with the primitive axis of x .

If we put in this equation

$$x = 0, \quad y = \sqrt{-b^2} = \pm b \sqrt{-1}, \quad \text{and} \quad y = 0, \quad x = \pm a',$$

since the value of y , when $x = 0$, is impossible, the curve does not meet the axis of y , but as $\pm a'$ is the value of x when $y = 0$, the curve meets the axis of x at the extremities of the diameter $2a'$.

Now, if in equation (5) we put a'^2 for x^2 and $-b^2$ for y^2 , we obtain the relation

$$-a'^2 b'^2 = \frac{-a^2 b^2}{\sin^2(a' - a)},$$

$\therefore ab \sin(a' - a) = ab$, which is the same as in the ellipse.

(55.) In the general properties of the ellipse, if we put $-b^2$ for b^2 , we have the corresponding general properties of the hyperbola.

(56.) In the values of a'^2 , b'^2 , obtained in art. 54, replacing the expressions for $\sin^2 a$, $\cos^2 a$, $\sin^2 a'$, $\cos^2 a'$, by their values in terms of $\tan a$ and $\tan a'$, viz. —

$$\cos^2 a = \frac{1}{1 + \tan^2 a}, \quad \sin^2 a = \frac{\tan^2 a}{1 + \tan^2 a},$$

$$\cos^2 a' = \frac{1}{1 + \tan^2 a'}, \quad \sin^2 a' = \frac{\tan^2 a'}{1 + \tan^2 a'},$$

$$\text{we obtain } a'^2 = \frac{-a^2 b^2 (1 + \tan^2 a)}{a^2 \tan^2 a - b^2} \dots\dots\dots (1),$$

$$-b'^2 = \frac{-a^2 b^2 (1 + \tan^2 a')}{a^2 \tan^2 a' - b^2},$$

$$\text{or } b'^2 = \frac{a^2 b^2 (1 + \tan^2 a')}{a^2 \tan^2 a' - b^2} \dots\dots\dots (2).$$

$$\text{From (1),} \quad \tan^2 a = \frac{b^2}{a^2} \left(\frac{a'^2 - a^2}{a'^2 + b^2} \right).$$

From (2), $\tan^2 \alpha' = \frac{b^2}{a^2} \left(\frac{a^2 + b'^2}{b'^2 - b^2} \right);$

multiplying these together, we have

$$\tan^2 \alpha \cdot \tan^2 \alpha' = \frac{b^4}{a^4} \left\{ \frac{(a'^2 - a^2)(a^2 + b'^2)}{(a'^2 + b^2)(b'^2 - b^2)} \right\},$$

$$\text{but } \tan \alpha \tan \alpha' = \frac{b^2}{a^2} \quad (\text{page } 75),$$

$$\text{or } \tan^2 \alpha \tan^2 \alpha' = \frac{b^4}{a^4},$$

$$\therefore \frac{b^4}{a^4} \left\{ \frac{(a'^2 - a^2)(a^2 + b'^2)}{(a'^2 + b^2)(b'^2 - b^2)} \right\} = \frac{b^4}{a^4},$$

$$\text{or } \frac{(a'^2 - a^2)(a^2 + b'^2)}{(a'^2 + b^2)(b'^2 - b^2)} = 1,$$

$$\therefore (a'^2 - a^2)(a^2 + b'^2) = (a'^2 + b^2)(b'^2 - b^2),$$

$$a'^2 a^2 - a^4 + a'^2 b'^2 - a^2 b'^2 = a'^2 b'^2 + b^2 b'^2 - a'^2 b^2 - b^4;$$

transposing and reducing,

$$a'^2 (a^2 + b^2) - b'^2 (a^2 + b^2) = a^4 - b^4,$$

$$\text{or } (a'^2 - b'^2)(a^2 + b^2) = a^4 - b^4 = (a^2 + b^2)(a^2 - b^2),$$

$$\therefore a'^2 - b'^2 = a^2 - b^2.$$

This might have been deduced, as well as the latter part of art. 54, more elegantly, by transforming the oblique ordinates to rectangular, as we have already done in the ellipse, substituting for x and y the values

$$\frac{x \sin' - y \cos \alpha}{\sin(\alpha' - \alpha)}, \quad \frac{y \cos \alpha - x \sin \alpha}{\sin(\alpha' - \alpha)};$$

in the equation $a'^2 y - b'^2 x = -a'^2 b'^2$, we obtain, as in the ellipse,

$$a'^2 \cos^2 \alpha - b'^2 \cos^2 \alpha' = a^2 \dots\dots\dots (1),$$

$$a'^2 \sin^2 \alpha - b'^2 \sin^2 \alpha' = -b^2 \dots\dots\dots (2),$$

$$-a'^2 b'^2 \sin^2 (\alpha' - \alpha) = -a^2 b^2 \dots\dots\dots (3),$$

by adding (1) and (2), we obtain

$$a'^2 - b'^2 = a^2 - b^2 \dots\dots\dots (4).$$

$$\text{From equation (3), } 4 a' b' \sin (\alpha' - \alpha) = 4 a b \dots\dots (5).$$

Equation (4) shews that the difference of the squares of any two conjugate diameters is equal to the difference of the square of the principal axes.

Equation (5) shews that the rectangle described on any system of conjugate diameters, is equal to the rectangle on the axes.

(57.) We have already shewn, that all we have hitherto done for the ellipse will, with slight modifications, apply to the hyperbola; but the converse of this is not true, for there are many properties of the hyperbola which cannot belong to the ellipse; these are the properties relative to the asymptotes. To complete the whole, we shall proceed to find the equation to the hyperbola referred to its asymptotes.

The equation of the hyperbola referred to its axes is

$$a^2 y^2 - b^2 x^2 = -a^2 b^2 \dots\dots\dots (1).$$

We must now pass from a system of rectangular to a system of oblique axes; to do this, we must substitute in the above equation, for x and y , the following values (see art. 24, eq. 3.):—


$$x = x \cos \alpha + y \cos \alpha',$$

$$y = x \sin \alpha + y \sin \alpha',$$

and we have

$$\left. \begin{aligned} & (a^2 \sin^2 \alpha - b^2 \cos^2 \alpha) y^2 \\ & + (2 a^2 \sin \alpha \sin \alpha' - 2 b^2 \cos \alpha \cos \alpha') x y \\ & + (a^2 \sin^2 \alpha - b^2 \cos^2 \alpha) x^2 \end{aligned} \right\} = -a^2 b^2 \dots (2).$$

Here we take for the new axis of x , the asymptote DK ;



and for the axis of y , the asymptote DL ; the angles α and α' are determined by the equations

$$\tan \alpha = -\frac{b}{a}, \quad \tan \alpha' = \frac{b}{a},$$

$$\therefore \cos \alpha = \frac{a}{\sqrt{a^2 + b^2}}, \quad \sin \alpha' = -\frac{b}{\sqrt{a^2 + b^2}},$$

$$\cos \alpha' = \frac{a}{\sqrt{a^2 + b^2}}, \quad \sin \alpha = \frac{b}{\sqrt{a^2 + b^2}},$$

$$\cos^2 \alpha' = \frac{a^2}{a^2 + b^2}, \quad \text{and} \quad \sin^2 \alpha' = \frac{b^2}{a^2 + b^2};$$

multiply the latter by a^2 , and the former by b^2 , then, by subtraction, we have

$$a^2 \sin^2 \alpha' - b^2 \cos^2 \alpha' = \frac{a^2 b^2 - a^2 b^2}{a^2 + b^2} = 0.$$

In the same way,

$$a^2 \sin^2 \alpha - b^2 \cos^2 \alpha = \frac{a^2 b^2 - a^2 b^2}{a^2 + b^2} = 0.$$

Also,

$$2a^2 \sin \alpha \sin \alpha' - 2b^2 \cos \alpha \cos \alpha' = -\frac{4a^2 b^2}{a^2 + b^2}.$$

Hence we see the terms involving x^2 and y^2 vanish, and if we substitute in equation (2) the value of the coefficient of xy , just found, we have

$$xy = \frac{a^2 + b^2}{4} \dots\dots\dots (3).$$

The equation which contains the rectangle of the variables, and the known quantity $\frac{a^2 + b^2}{4}$, is called the equation to the hyperbola, referred to its asymptotes as co-ordinate axes.

(58.) If the hyperbola be equilateral, then

$$a = b, \quad \text{and} \quad xy = \frac{a^2}{2}.$$

(59.) If we wished conversely to determine from equation (2) the angles α and α' , so that the terms involving x^2 and y^2 should disappear, we have then the following conditions:—

$$a^2 \sin^2 \alpha' - b^2 \cos^2 \alpha' = 0, \quad a^2 \sin^2 \alpha - b^2 \cos^2 \alpha = 0.$$

From the first of these we have $\tan \alpha' = \pm \frac{b}{a}$,

from the second $\tan \alpha = \pm \frac{b}{a}$.

This shews, that if we take α' for the angle whose tangent is $+\frac{b}{a}$, we must take α for the angle whose tangent is $-\frac{b}{a}$. Therefore, the new axes, with respect to which the equation is brought to the form $xy = k^2$, are the asymptotes of the curve, where k^2 is put for $\frac{a^2 + b^2}{4}$.

It is clear from this equation, or $y = \frac{k^2}{x}$, or $x = \frac{k^2}{y}$, that as we increase x the more is y diminished; and if we suppose x to be greater than any assignable quantity, y will become less than any assignable quantity, and conversely; so that the new axes have the characteristic property of the asymptotes.

Let ϵ be the angle LDK , which the asymptotes make with each other; then, multiplying both sides of the equation $xy = k^2$ by $\sin \epsilon$, we obtain

$$xy \sin \epsilon = k^2 \sin \epsilon.$$

Now, if P be any point in the curve, MP and MD the co-ordinates of this point, parallel to the asymptotes, then the parallelogram $DMPQ$ has for its area

$$DM \cdot PH = xy \sin \epsilon;$$

this is equal to $k^2 \sin \epsilon$, which is a constant quantity, and is independent of the position of the point P .

Therefore, all parallelograms described upon the co-ordinates parallel to the asymptotes are equal. It can be easily shown that this constant area is equal to half the rectangle $DBEC$ contained by the semi-axes, for $\epsilon = 2\alpha'$,

$$\therefore \sin \epsilon = \sin 2\alpha' = 2 \sin \alpha' \cos \alpha';$$

$$\text{by art. 57, } \cos \alpha' = \frac{a}{\sqrt{a^2 + b^2}}, \quad \sin \alpha' = \frac{b}{\sqrt{a^2 + b^2}},$$

$$\therefore \sin \epsilon = 2 \sin \alpha' \cos \alpha' = \frac{2ab}{\sqrt{a^2 + b^2}},$$

$$\therefore xy \sin \epsilon = \frac{a^2 + b^2}{4} \cdot \sin \epsilon \text{ becomes}$$

$$xy \sin \epsilon = \frac{a^2 + b^2}{4} \cdot \frac{2ab}{\sqrt{a^2 + b^2}} = \frac{ab}{2}.$$

To the extremities of the minor axis draw the lines BC, BD, AC, AD; the figure thus formed, having its sides parallel to the asymptotes, is four times the area of the figure DIBI', constructed on the co-ordinates of the point A.

Therefore, if from any number of points in the hyperbola, lines be drawn parallel to the asymptotes, and terminating therein, the parallelograms so formed will all be equal to each other, and to half the rectangle of the principal semi-axes.

(60.) To find the equation of the tangent to any point of the hyperbola referred to its asymptotes.

Let x'', y'' , be the co-ordinates of the point P; x', y' , the co-ordinates of another point in the curve; the equation to the hyperbola being

$$xy = k^2 \dots\dots\dots (1).$$

The secant will be represented by

$$y - y'' = \frac{y' - y''}{x' - x''}(x - x'') \dots\dots\dots (2),$$

and $x' y'$, and $x'' y''$, being both points in the curve,

$$x' y' = k^2 \dots\dots\dots (3),$$

$$x'' y'' = k^2 \dots\dots\dots (4).$$

To find the coefficient $\frac{y' - y''}{x' - x''}$, subtract (4) from (3), and we have $x'y' - x''y'' = 0$; and adding $y''x'$ and subtracting $x'y''$, which are identical, we obtain

$$x'(y' - y'') + y''(x' - x'') = 0,$$

$$\therefore \frac{y' - y''}{x' - x''} = -\frac{y''}{x'};$$

the secant therefore becomes

$$y - y'' = -\frac{y''}{x'}(x - x').$$


Now, in order that this secant may become a tangent, we must suppose $x' = x''$, and $y' = y''$,

$$\therefore y - y'' = -\frac{y''}{x'}(x - x')$$

is the equation to the tangent,

$$\text{or } y''x + x'y = 2x'y' = \frac{1}{2}(a^2 + b^2).$$

(61.) To find the area of a parabola.



$$\left. \begin{aligned} pn^2 &= 4a'mp \dots\dots\dots (1) \\ PN^2 &= 4a' \cdot MP \dots\dots\dots (2) \end{aligned} \right\} \text{by the equation to the parabola,}$$

$$\therefore pn^2 - PN^2 = 4a'(mp - MP),$$

$$\text{or } (pn + PN) \cdot (pn - PN) = 4a' \cdot Nn,$$

$$\overline{pn} + \text{PN} \cdot \text{Mm} = \frac{\text{PN}^2}{\text{MP}} \cdot \text{Nn}, \text{ from (2),}$$

$$\therefore \text{MP} \cdot \text{Mm} = \frac{\text{PN}^2}{pn + \text{PN}} \cdot \text{Nn}, \text{ (ultimately,)}$$

$$= \frac{1}{2} \text{PN} \cdot \text{Nn},$$

$$\text{or the rectangle Pm} = \frac{1}{2} \text{Pn};$$

and so on for any other rectangle; therefore the space Anp is double the space Apm ,

$$\text{or } \text{Anpm} = 3 \text{Apm}; \quad \therefore \text{Apm} = \frac{1}{3} \text{Anpm},$$

$$\text{or } \text{Anp} = \frac{2}{3} \text{Anpm}.$$

Hence, the parabola is $\frac{2}{3}$ of its circumscribing rectangle.

To find the Area of an Ellipse.

(62.) The area of an ellipse may be found from the constant ratio $\frac{b}{a}$, of the ordinate of an ellipse to that of the circle described upon the major axis.

Conceive any polygon whatever to be inscribed in a circle, of which one of the sides is MM' ; from M and M' let fall the perpendiculars MP and $\text{M}'\text{P}'$, these perpendiculars cut the ellipse in N and N' ; join NN' , and we thus obtain a polygon inscribed in the ellipse, of which NN' is one of its sides.

Let Y and Y' be the co-ordinates of the points M and M' , and y and y' the co-ordinates of the points N and N' , corresponding to the same abscissæ x , x' ; the trapeziums $\text{MM}'\text{PP}'$, $\text{NN}'\text{PP}'$, give

$$\text{MM}'\text{PP}' = \frac{\text{Y} + \text{Y}'}{2} (x - x'), \quad \text{NN}'\text{PP}' = \frac{y + y'}{2} (x - x'),$$


$$\therefore \frac{NN'PP'}{MM'PP'} = \frac{y + y'}{Y + Y'}.$$

Now, since, by art. 39,

$$y = \frac{b}{a} \cdot Y, \quad y' = \frac{b}{a} \cdot Y',$$

$$\therefore \frac{y + y'}{Y + Y'} = \frac{b}{a},$$

$$\therefore \frac{NN'PP'}{MM'PP'} = \frac{b}{a}.$$



In the same manner, each of the trapeziums which compose the polygon inscribed in the ellipse, is to the corresponding trapezium of the polygon inscribed in the circle as b is to a ; hence, we conclude that the sum of all the first trapeziums, or the polygon inscribed in the ellipse, is to the second polygon in the same ratio, or

$$\frac{P}{P'} = \frac{b}{a};$$

p and P being the polygons respectively. As this relation is true, whatever be the number of the sides of the two polygons, it is true in their limits, which are the areas of the ellipse and circle; hence, if s and S be the respective areas, we have

$$\frac{s}{S} = \frac{b}{a}, \quad \text{or } s = \frac{b}{a} \cdot S;$$

but π represents the ratio of the circumference of a circle to its diameter, or the area of a circle whose radius is 1, πa^2 is the area of a circle whose radius is a , and, consequently,

$$s = \frac{b}{a} \cdot S = \frac{b}{a} \pi a^2 = \pi a b,$$

which equals the area of the ellipse.

For the hyperbola the same relation $s = \frac{b}{a} \cdot S$, obtains, s being the area comprised between the curve and any chord

parallel to the second axis, & the corresponding area of an equilateral hyperbola, described upon the first axis. But this does not enable us to find the area of the hyperbola, since we must at first know that of an equilateral hyperbola.

On the Right Line, Circle, and various properties of the Conic Sections.

(63.) In the note, page 16, it is shewn that the length of the perpendicular from a point $x'y'$ on $Ax + By + C = 0$, is

$$\frac{Ax' + By' + C}{\sqrt{A^2 + B^2}};$$

by this we can shew that the length of the perpendicular from $x'y'$ on the line $x \cos \alpha + y \sin \alpha - p = 0$, is

$$x' \cos \alpha + y' \sin \alpha - p;$$

$$\text{for } \frac{A}{\sqrt{A^2 + B^2}} x' + \frac{B}{\sqrt{A^2 + B^2}} y' + \frac{C}{\sqrt{A^2 + B^2}} =$$

is of the same form, since we may take

$$\frac{A}{\sqrt{A^2 + B^2}} = \cos \alpha, \text{ and } \frac{B}{\sqrt{A^2 + B^2}} = \sin \alpha,$$

as they satisfy the condition $\cos^2 \alpha + \sin^2 \alpha = 1$.

This is useful in solving a large class of problems.

Given the base and area of a triangle, to find the locus of the vertex *.

Let the equation of the given base be

$$x \cos \alpha + y \sin \alpha - p = 0,$$

and $x \cos \alpha + y \sin \alpha - p$ is the length of the perpendicular from any point x, y , on the base; if the given base = a , and given area = m^2 , the equation of the locus is

$$x \cos \alpha + y \sin \alpha - p = \frac{2m^2}{a},$$

since the perpendicular of any triangle equals twice the area divided by the base.

* See Salmon's "Conic Sections," page 41.

(64.) The equation of a line bisecting the angle contained by the lines

$$x \cos \alpha + y \sin \alpha - p = 0, \text{ and } x \cos \beta + y \sin \beta - p' = 0,$$

$$\text{is } x \cos \alpha + y \sin \alpha - p = x \cos \beta + y \sin \beta - p',$$

which represents a right line passing through the intersection of these two lines, for it is evidently the equation to a right line; and since the co-ordinates of the points of intersection must satisfy the two former equations, they must also satisfy the latter equation.

(65) The general equation to the circle referred to rectangular axes being

$$(x - \alpha)^2 + (y - \beta)^2 = r^2,$$

$$\text{or } x^2 + y^2 - 2\alpha x - 2\beta y + \alpha^2 + \beta^2 - r^2 = 0 \dots\dots (1);$$

every equation of the second degree of the form

$$x^2 + y^2 + Ax + By + C = 0 \dots\dots\dots (2),$$

having the coefficients of x and y either unity or equal, and not containing the rectangle xy of the variables, is the equation to a circle.

Comparing (1) and (2),

$$-2\alpha = A, \quad -2\beta = B, \quad \text{and } \alpha^2 + \beta^2 - r^2 = C,$$

$$\alpha = -\frac{A}{2}, \quad \beta = -\frac{B}{2},$$

$$r = \sqrt{\alpha^2 + \beta^2 - C} = \sqrt{\frac{A^2 + B^2}{4} - C},$$

replacing α, β, r , by the above values,

$$\left(x + \frac{A}{2}\right)^2 + \left(y + \frac{B}{2}\right)^2 = \frac{A^2 + B^2}{4} - C.$$

Or by completing the squares in equation (2) we obtain the same result, thus

$$x^2 + Ax + y^2 + By = -C;$$

completing the square in each,

$$x^2 + Ax + \left(\frac{A}{2}\right)^2 + y^2 + By + \left(\frac{B}{2}\right)^2 = \frac{A^2}{4} + \frac{B^2}{4} - C,$$

$$\text{or } \left(x + \frac{A}{2}\right)^2 + \left(y + \frac{B}{2}\right)^2 = \frac{A^2 + B^2}{4} - C \dots\dots (3).$$

Examples.

$$\dots\dots\dots 2x^2 + 2y^2 - 3x + 4y - 1 = 0,$$

$$x^2 + y^2 - \frac{3x}{2} + 2y = \frac{1}{2};$$

putting the equation in the following form,

$$x^2 - \frac{3x}{2} + y^2 + 2y = \frac{1}{2},$$

complete the squares,

$$\begin{aligned} x^2 - \frac{3x}{2} + \left(\frac{3}{4}\right)^2 + y^2 + 2y + 1 &= \frac{9}{16} + 1 + \frac{1}{2} \\ &= \frac{9}{16} + \frac{16}{16} + \frac{8}{16} = \frac{33}{16}, \quad \left(x - \frac{3}{4}\right)^2 + (y + 1)^2 = \frac{33}{16}, \end{aligned}$$

which is the equation to a circle whose radius is $\frac{\sqrt{33}}{4}$,

and the co-ordinates of the centre $\frac{3}{4}$ and -1 .

Since A, B, C, may be any quantities whatever, we may sometimes have $\frac{A^2 + B^2}{4} - C = 0$, or indeed negative.

If it be equal 0 then the circle is reduced to a point, and if it be negative the radius is imaginary; thus, the equation

$$4x^2 + 4y^2 - 12x - 8y + 13 = 0$$

represents a point whose co-ordinates are $\frac{3}{2}$ and 1, for it can be transformed into

$$\left(x - \frac{3}{2}\right)^2 + (y - 1)^2 = 0.$$

The equation $x^2 + y^2 + 4x - 2y + 7 = 0$,

$$x^2 + 4x + y^2 - 2y = -7,$$

completing the square;

$$x^2 + 4x + 4 + y^2 - 2y + 1 = 4 + 1 - 7 = -2$$

$$(x + 2)^2 + (y - 1)^2 = -2,$$

an equation of which the first member, being the sum of two positive squares, cannot be equal to a negative quantity.

(66.) In the parabola, if PSP' be a chord through the focus, and PSN = θ , then, page 37,

$$SP = \frac{l}{1 - \cos \theta}, \quad \text{and} \quad SP' = \frac{l}{1 + \cos \theta},$$

$$\frac{1}{SP} + \frac{1}{SP'} = \frac{1 - \cos \theta}{l} + \frac{1 + \cos \theta}{l} = \frac{2}{l},$$

where $l = 2a =$ half the latus rectum.

(67.) In the parabola, the parameter of any diameter = 4 times (distance of the vertex of the diameter from the focus).

The parameter is the double ordinate passing through the focus.

$$\begin{aligned} SQ &= \frac{2a}{1 + \cos ASQ}, & sq &= \frac{2a}{1 + \cos(\pi + qsx)} \\ & & &= \frac{2a}{1 + \cos(\pi + ASQ)} \\ & & &= \frac{2a}{1 - \cos AS} \end{aligned}$$

$$\pi = \text{ASP} + 2(\pi - \text{ASQ}), \quad \therefore \text{ASQ} = \frac{\pi}{2} + \frac{\text{ASP}}{2},$$

$$\therefore \text{ASQ} = \frac{\pi}{2} + \frac{\text{ASP}}{2},$$

$$\text{Qq} = \frac{4a}{\cos^2 \frac{\text{ASP}}{2}} = 4 \text{SP}.$$

(68.) Find the area included between the parabola $y^2 = 4ax$ and the straight line $x = y + a$.


Since $x = a$ gives $y = 0$, we see that the given line passes through the focus of the parabola; and because the coefficient of x is 1, the line makes an angle of 45° with the axis of x .

Hence, in the figure,

APP' is the area required,

and $\angle \text{PSN} = 45^\circ$,

$\therefore \text{SN} = \text{PN}$, and $\text{SN}' = \text{P}'\text{N}'$.



Now, $\text{APP}' = \text{ASP} + \text{ASP}'$

$$= \text{APN} - \text{SPN} + \text{AN}'\text{P}' + \text{SP}'\text{N}'$$

$$= \frac{2}{3} (\text{AN} \cdot \text{PN} + \text{AN}' \cdot \text{P}'\text{N}') - \frac{1}{2} (\text{PN}^2 - \text{P}'\text{N}'^2).$$

Find x from equations $y^2 = 4ax$

$$\text{and } x = y + a,$$

we get $y^2 = (x - a)^2 = 4ax$, $\therefore x^2 - 6ax + a^2 = 0$

$$\therefore x = 3a \pm 2\sqrt{2} \cdot a = (3 \pm 2.8284)a,$$

$$\therefore \text{AN} = (5.8284), \text{AN}' = (0.1716)a,$$

and $y = x - a$, $\therefore \text{PN} = 4.8284$, $\text{P}'\text{N}' = (0.8284)a$,


$$\therefore AP = \frac{2}{3} \cdot a^2 \{ (5.8284)(4.8284) + (0.1716)(0.8284) \}$$

$$= \frac{1}{2} a^2 \{ (4.8284)^2 - (0.8284)^2 \}$$

$$= a^2 (18.80124 - 11.31360) = 7.48764 \times a^2. \quad \text{Answer.}$$

(69.) Draw a normal at the extremity of the latus rectum of a parabola whose equation is $y^2 = 4a(x - a)$, and find its distance from the origin of co-ordinates.

If $x = a$, we find from the given equation that $y = 0$; therefore, the curve cuts the axis of x at a distance a from the origin. If $x < a$, then y is impossible; therefore, no portion of the curve lies to the left of $(a, 0)$, that is, A. Hence, A is the vertex of the parabola. Transfer the origin to this point, and we find $y'^2 = 4ax'$,



since $y' = y$ and $x' = x - a$,

therefore X, that is, the old origin, is the foot of the directrix. To draw a normal, then, at the extremity of the latus rectum, take $SG = SP = 2a$, and join PG ; this is the normal required,

for $\angle SPG = \angle SGP = \angle GPx$.

Join XP , then this is the distance of the normal from the origin, and

$$XP = \sqrt{SP^2 + XS^2} = \sqrt{8AS^2} = 2\sqrt{2} \cdot a.$$

(70.) Find the locus of the intersection of two tangents to a hyperbola which meet one another at right angles.

The equation to the tangent to a hyperbola, in terms of the angle which it makes with the axis of x , is

$$y = mx \mp \sqrt{m^2 a^2 - b^2};$$

consequently, for another which intersects this at right angles, we get

$$y = -\frac{1}{m} x \mp \sqrt{\frac{a^2}{m^2} - b^2}$$

Now, this equation becomes

$$my + x = \mp \sqrt{a^2 - m^2 b^2},$$

$$\therefore m^2 y^2 + 2mxy + x^2 = a^2 - m^2 b^2 \dots\dots (1);$$

and from the first equation

$$y - mx = \mp \sqrt{m^2 a^2 - b^2},$$

$$\therefore y^2 - 2mxy + m^2 x^2 = m^2 a^2 - b^2 \dots\dots\dots (2);$$

adding (1) and (2), we get

$$(y^2 + x^2)(m^2 + 1) = (a^2 - b^2)(m^2 + 1),$$

$$\therefore x^2 + y^2 = a^2 - b^2.$$

Hence, the locus is a circle whose centre is the centre of the hyperbola, and radius $= \sqrt{a^2 - b^2}$.

(71.) Shew that the parameter belonging to any diameter of a parabola varies inversely as the square of the sine of the angle at which the corresponding ordinates are inclined to it.

Let Px' be any diameter, SP its parameter, PT a tangent at its extremity, α the angle at which the ordinates to Px' are inclined, and, consequently, $PTS = \alpha$.

Then we know that it is a property of the parabola


that $TPS = PTS = \alpha$,

$$\therefore PSN = 2\alpha.$$

$$\text{Now, } \frac{SN}{SP} = \cos 2\alpha = 1 - 2\sin^2 \alpha,$$

$$\text{and } SN = AN - AS = AN + AS - 2AS = SP - 2AS,$$

$$\therefore \frac{SP - 2AS}{SP} = 1 - 2\sin^2 \alpha, \quad \therefore AS = SP \sin^2 \alpha,$$



or $SP = \frac{AS}{\sin^2 \alpha}$, $\therefore SP \propto \frac{1}{\sin^2 \alpha}$, since AS is constant.

(72.) If CP' , CQ' , be diameters of an ellipse, making angles α, β , respectively with the axis major, then

$$\frac{CP'^2}{CQ'^2} = \frac{a'^2}{a''^2} = \frac{a^2 \sin^2 \beta + b^2 \cos^2 \beta}{a^2 \sin^2 \alpha + b^2 \cos^2 \alpha}.$$

Let $xy, x'y'$, be the co-ordinates of P' and Q' respectively,

then $x = a' \cos \alpha$, and $x' = a'' \cos \beta$,

$y = a' \sin \alpha$, $y' = a'' \sin \beta$;

therefore, substituting in the equation to the ellipse at the points $xy, x'y'$,

$$a^2 b^2 = a''^2 \{a^2 \sin^2 \beta + b^2 \cos^2 \beta\},$$

$$a^2 b^2 = a'^2 \{a^2 \sin^2 \alpha + b^2 \cos^2 \alpha\};$$

or dividing the former by the latter,

$$1 = \frac{a''^2}{a'^2} \cdot \frac{a^2 \sin^2 \beta + b^2 \cos^2 \beta}{a^2 \sin^2 \alpha + b^2 \cos^2 \alpha},$$

$$\therefore \frac{a'^2}{a''^2} = \frac{a^2 \sin^2 \beta + b^2 \cos^2 \beta}{a^2 \sin^2 \alpha + b^2 \cos^2 \alpha}.$$

(73.) From any point P in an ellipse draw the ordinate PN; join P and the centre C of the ellipse, and also N and the extremity B of the conjugate axis. The locus of their intersection O is required.

Let x', y' , be the co-ordinates of P, then


$$y = \frac{y'}{x'} x$$

is equation to PC (1),

$$\text{and } y = -\frac{b}{x'} x + b$$

is equation to BN (2),

$$x'y = -bx + bx',$$



$$x'(y - b) = -bx,$$

$$x' = -\frac{bx}{y-b}, \quad \therefore x'^2 = \frac{b^2 x^2}{(y-b)^2};$$

$$\text{and from (1), } x'y = xy'$$

$$y' = \frac{x'y}{x},$$

$$\begin{aligned} \therefore y'^2 &= \frac{x'^2 y^2}{x^2} = \frac{b^2 x^2 y^2}{x^2 (y-b)^2} \\ &= \frac{b^2 y^2}{(y-b)^2}. \end{aligned}$$

Also, since $x'y'$ is a point in the ellipse, we have

$$a^2 y'^2 + b^2 x'^2 = a^2 b^2;$$

$$\text{substituting } a^2 y'^2 + b^2 x'^2 = a^2 (y-b)^2$$

$$= a^2 y^2 - 2a^2 by + a^2 b^2$$


$$= b^2 x^2 + 2a^2 y = a^2 b^2, \quad \therefore \text{locus required is a parabola.}$$

(74.) At what point in an ellipse is the angle formed by the two focal distances greatest?

Let P be the required point.

$$\angle SPH = \theta,$$

$$CN = x \quad NP = y.$$



$$\text{Now } \tan \theta = -\tan (PHS + PSH)$$

$$= \frac{\tan PHS + \tan PSH}{\tan PHS \cdot \tan PSH - 1}$$

$$= \frac{\frac{y}{ae-x} + \frac{y}{ae+x}}{\frac{y^2}{a^2 e^2 - x^2} - 1} = \frac{2aey}{y^2 - a^2 e^2 + x^2},$$

which must be a maximum when

$$y^2 - a^2 e^2 + x^2 = 0;$$

$$i.e., \text{ when } x^2 + y^2 = a^2 e^2 = CH^2.$$


$$\text{But } x^2 + y^2 = CP^2 = CH^2, \quad \therefore CP = CH.$$

Hence, with centre C and radius CH, describe a circle cutting the ellipse in P, this will then be the required point.

(75.) The area between two normals to a parabola at the extremities of a focal chord and the curve $= \frac{20 a^2}{3 \sin^4 2\theta}$, θ being the angle which one of the normals makes with the axis.

Let PP' be the chord,

PQ and P'Q the normals, which, since PP' passes through the focus, are at right angles to one another.



Take x, y , the co-ordinates of P and x', y' , those of P'; also the angle $PSG = \theta$, $\therefore \angle SPG = \theta$.

Now,

$$\begin{aligned} \text{area APP'} &= \frac{2}{3} \{xy + x'y'\} + \frac{1}{2} \{(a - x')y' - (x - a)y\} \\ &= \frac{1}{6} \{xy + x'y'\} + \frac{a}{2} \{y + y'\} \\ &= \frac{1}{24a} \{y^3 + y'^3\} + \frac{a}{2} \{y + y'\}, \end{aligned}$$

since xy and $x'y'$ are points in the parabola.

$$\begin{aligned} \text{But } y &= PS \sin 2\theta, \text{ and } PS = a + r = 2a + PS \cos 2\theta \\ &= 2a - PS \cos 2\theta, \end{aligned}$$

$$\therefore PS = \frac{2a}{1 + \cos 2\theta} = \frac{a}{\cos^2 \theta},$$

$$\text{and } y' = P'S \cdot \sin 2\theta,$$

Likewise, $P'S = 2a + P'S \cos 2\theta$,

$$\therefore P'S = \frac{2a}{1 - \cos 2\theta} = \frac{a}{\sin^2 \theta},$$

$$\therefore y = 2a \tan \theta, \text{ and } y' = 2a \cot \theta,$$

$$\begin{aligned} \therefore APP' &= \frac{8a^3}{24a} \{\tan^3 \theta + \cot^3 \theta\} + a^2 \{\tan \theta + \cot \theta\} \\ &= \frac{a^2}{3} \left\{ \frac{\sin^3 \theta + \cos^3 \theta + 3 \sin^2 \theta \cos \theta}{\sin^3 \theta \cos^3 \theta} \right\} \\ &= \frac{a^2}{3} \left\{ \frac{\sin^4 \theta + 2 \sin^2 \theta \cos^2 \theta + \cos^4 \theta}{\sin^3 \theta \cos^3 \theta} \right\} = \frac{a^2}{3 \sin^3 \theta \cdot \cos^3 \theta}. \end{aligned}$$

Again,

$$\begin{aligned} \Delta PQP' &= \frac{PQ \cdot P'Q}{2} = \frac{PP' \cos \theta}{2} \frac{PP' \sin \theta}{2} = \frac{PP'^2}{2} \sin \theta \cos \theta \\ &= \frac{1}{2} \{PS^2 + 2PS \cdot P'S + P'S^2\} \sin \theta \cdot \cos \theta \\ &= \frac{a^2}{2} \left\{ \frac{\sin \theta}{\cos^3 \theta} + \frac{2}{\sin \theta \cdot \cos \theta} + \frac{\cos \theta}{\sin^3 \theta} \right\} \\ &= \frac{a^2}{2} \left\{ \frac{\sin^4 \theta + 2 \sin^2 \theta \cos^2 \theta + \cos^4 \theta}{\sin^3 \theta \cdot \cos^3 \theta} \right\} + \frac{a^2}{2 \sin^3 \theta \cdot \sin^3 \theta}, \end{aligned}$$

therefore, the area

$$\begin{aligned} \Delta PQP' &= APP' + PQP' = \left(\frac{a^2}{3} + \frac{a^2}{2} \right) \cdot \frac{1}{\sin^3 \theta \cdot \cos^3 \theta} \\ &= \frac{5a^2}{6 \sin^3 \theta \cdot \cos^3 \theta} = \frac{20a^2}{3 \cdot (8 \sin^3 \theta \cdot \cos^3 \theta)} \\ &= \frac{20a^2}{3 \sin^3 2\theta}. \end{aligned}$$

EQUATIONS OF THE SECOND DEGREE*.

(76.) The general form for equations of the second degree, being those in which the ordinates xy are involved to the second power, is

$$Ax^2 + By^2 + Cxy + ax + by + c = 0,$$

wherein each of the constants A, B, C, a, b, c , may be either *positive* or *negative*.

Let us, in the first place, transfer the equation to two other rectangular axes parallel to the original ones, and having their origin at a point whose ordinates are $x'y'$; and (24) by substituting $x + x'$ and $y + y'$ for x and y , we shall find the corresponding equation to be

$$\begin{aligned} &A(x^2 + 2x'x + x'^2) + B(y^2 + 2y'y + y'^2) \\ &+ C(xy + y'x + x'y' + x'y') \\ &+ a(x + x') + b(y + y') + c = 0; \end{aligned}$$

which, arranged for x and y , becomes

$$\begin{aligned} &Ax^2 + By^2 + Cxy \\ &+ (2Ax' + Cy' + a)x + (2By' + Cx' + b)y \\ &+ (Ax'^2 + By'^2 + Cx'y' + ax' + by' + c) = 0. \end{aligned}$$

(77.) The first three coefficients, A, B, C , stand unaffected with the new constants, x', y' , by which we observe that they are independent of the position of the origin; and hence the position of the origin of any equation of the second degree depends entirely on the values of the three last coefficients a, b, c .

(78.) We may now assume the values of the two ordinates x', y' , at pleasure, since the position of the new origin is entirely arbitrary; and consequently, by the principles of algebra, we may fulfil any two possible conditions which involve them;

* See Woolhouse's "Algebraic Geometry."

let us therefore put the coefficients of x and y each equal to nothing, viz. :—

$$2 A x' + C y' + a = 0,$$

$$2 B y' + C x' + b = 0;$$

and thence

$$x' = \frac{C b - 2 B a}{4 A B - C^2}, y' = \frac{C a - 2 A b}{4 A B - C^2};$$

hence also, by substitution, the last term

$$A x'^2 + B y'^2 + C x' y' + a x' + b y' + c = \frac{C a b - A b^2 - B a^2}{4 A B - C^2} + c;$$

or by assuming

$$C a b - A b^2 - B a^2 + c (4 A B - C^2) = G,$$

it becomes

$$= \frac{G}{4 A B - C^2}$$

The equation is thus transformed into

$$A x^2 + B y^2 + C x y + \frac{G}{4 A B - C^2} = 0 \dots\dots\dots (a),$$

in which the fourth and fifth terms are wanting.

(79.) Let us now transfer this equation to two other rectangular axes, inclined at an angle ω with the former, and retaining the same origin; and (24) substituting $x \cos \omega - y \sin \omega$ and $x \sin \omega + y \cos \omega$ for x and y , we get for the corresponding equation

$$\begin{aligned} & A (x' \cos^2 \omega + y'^2 \sin^2 \omega - 2 x' y' \cos \omega \sin \omega) \\ & + B (x'^2 \sin^2 \omega + y'^2 \cos^2 \omega + 2 x' y' \cos \omega \sin \omega) \\ & + C \{x'^2 \cos \omega \sin \omega - y'^2 \cos \omega \sin \omega + x' y' (\cos^2 \omega - \sin^2 \omega)\} \\ & + \frac{G}{4 A B - C^2} = 0, \end{aligned}$$

which, arranged for x and y , observing that

$$\cos^2 \omega - \sin^2 \omega = \cos 2\omega \text{ and } 2 \cos \omega \sin \omega = \sin 2\omega,$$

becomes

$$\begin{aligned} & (A \cos^2 \omega + B \sin^2 \omega + C \cos \omega \sin \omega) x^2 \\ & + (A \sin^2 \omega + B \cos^2 \omega - C \cos \omega \sin \omega) y^2 \\ & + \{C \cos 2\omega - (A - B) \sin 2\omega\} xy \\ & + \frac{G}{4AB - C^2} = 0. \end{aligned}$$

By taking the value of ω so as to exterminate xy ,

$$C \cos 2\omega - (A - B) \sin 2\omega = 0,$$

$$\text{and } \tan 2\omega = \frac{C}{A - B},$$

which reduces the equation to

$$\begin{aligned} & (A \cos^2 \omega + B \sin^2 \omega + C \cos \omega \sin \omega) x^2 \\ & + (A \sin^2 \omega + B \cos^2 \omega - C \cos \omega \sin \omega) y^2 \\ & + \frac{G}{4AB - C^2} = 0; \end{aligned}$$

and hence it appears that every line of the second order may be referred to two determinate rectangular axes so that its equation shall be transformed into the above form. By assuming

$$A \cos^2 \omega + B \sin^2 \omega + C \cos \omega \sin \omega = A'',$$

$$A \sin^2 \omega + B \cos^2 \omega - C \cos \omega \sin \omega = B'',$$

it becomes

$$A'' x^2 + B'' y^2 + \frac{G}{4AB - C^2} = 0 \dots\dots\dots (b).$$

(80.) Now if the principal semi-diameters of an ellipse and hyperbola be denoted by a' , b' , and the former be taken for
F 2

the axis of x and the origin at the centre, their equations will be as follow:

For the ellipse,

$$\frac{x^2}{a'^2} + \frac{y^2}{b'^2} = 1, \text{ or } b'^2 x^2 + a'^2 y^2 - a'^2 b'^2 = 0;$$

and for the hyperbola,

$$\frac{x^2}{a'^2} - \frac{y^2}{b'^2} = \pm 1, \text{ or } b'^2 x^2 - a'^2 y^2 \mp a'^2 b'^2 = 0*,$$

the under sign representing the conjugate hyperbola.

The signs may be all changed if necessary.

By means of these two equations and the foregoing transformed equation, (b), we deduce the following particulars relative to the general equation.

(81.) 1st. When A'', B'' , are both negative, and $G, 4AB - C^2$ have the same sign, the equation determines an *ellipse*; and when A'', B'' , are both of them positive, and G and $4AB - C^2$ have different signs, the locus is also an *ellipse*†.

(82.) 2nd. When A'', B'' , are of different signs, and G not = 0, the locus is an *hyperbola*.

(83.) 3rd. In each of these cases the squares of the principal semi-diameters are equal to

$$\frac{\pm G}{A''(4AB - C^2)}, \quad \frac{\pm G}{B''(4AB - C^2)},$$

the under sign being for the ellipse, and either sign for the hyperbola,

(84.) 4th. The values of G, A'', B'' , are determined from the equations

$$G = Cab - Ab^2 - Ba^2 + c(4AB - C^2) \dots\dots\dots (1),$$

$$\tan 2\omega = \frac{C}{A - B} \dots\dots\dots (2),$$

$$\left. \begin{aligned} A'' &= A \cos^2 \omega + B \sin^2 \omega + C \cos \omega \sin \omega \\ B'' &= A \sin^2 \omega + B \cos^2 \omega - C \cos \omega \sin \omega \end{aligned} \right\} \dots\dots\dots (3),$$

* For the immediate values of A'', B'' , see article 103.

† The conjugate hyperbolas have the conjugate axis of the other hyperbolas for the transverse, and transverse of the others for the conjugate.

wherein ω is the angle included between the original axis of x and the principal diameter of the curve.

(85.) 5th. The position of the centre of the curve is determined by

$$x' = \frac{Cb - 2Ba}{4AB - C^2}, y' = \frac{Ca - 2Ab}{4AB - C^2}.$$

(86.) 6th. When the equation is of the form

$$Ax^2 + By^2 + Cx + c = 0,$$

wherein the fourth and fifth terms of the general equation are wanting, we have $a = 0$, $b = 0$, and thence $x' = 0$, $y' = 0$, which therefore shews the origin to be at the centre of the curve. This agrees with equation (a), article 78, where the origin is transferred to the centre.

(87.) 7th. By adding the equations (3), article 84, we find

$$A'' + B'' = A + B.$$

Hence we see that, whatever be the position of the axes of co-ordinates, the sum of the coefficients of x^2 and y^2 will be the same.

(88.) 8th. When $G = 0$ and also A'' and B'' of different signs, the general equation defines a straight line.

For in this case the transformed equation (b), article 79, becomes

$$A''x^2 + B''y^2 = 0,$$

which gives

$$\frac{y}{x} = \pm \sqrt{-\frac{A''}{B''}};$$

and this value is real when A'' , B'' , have different signs.

(89.) 9th. In the two following cases it will be found that no real values of x and y can possibly fulfil the equation (b); and consequently that the equation can have no locus.

First. When G and $4AB - C^2$ are of the same sign, and A'' , B'' , both of them positive.

Second. When G and $4AB - C^2$ are of different signs, and A'' , B'' , are both negative.

(90.) 10th. When $G = 0$, and A'' , B'' , have the same sign, no real values of x and y can satisfy the equation (b), except the particular case of $x = 0$, $y = 0$. In this case, therefore, the locus is the single point corresponding with the new origin $x'y'$.

(91.) 11th. It appears that by changing the position of the origin to the centre $x'y'$, the equation

$$Ax^2 + By^2 + Cxy + ax + by + c = 0$$

is transformed into the form

$$Ax^2 + By^2 + Cxy + h = 0,$$

$$\text{wherein } h = \frac{G}{4AB - C^2}.$$

Also, that by taking two other axes of co-ordinates making an angle α with these, so that

$$\tan 2\alpha = \frac{C}{A - B},$$

the equation

$$Ax^2 + By^2 + Cxy + h = 0$$

becomes of the form

$$A''x'^2 + B''y'^2 + h = 0,$$

wherein $A'' + B'' = A + B$ and the constant h is unchanged.

(92.) 12th. Let x'' , y'' , be the two semi diameters of the curve

$$Ax^2 + By^2 + Cxy + h = 0,$$

which coincide with the axes of co-ordinates to which it is referred, and they will be determined by taking first $y = 0$ and then $x = 0$ in the equation, the results being

$$x''^2 = -\frac{h}{A}, y''^2 = -\frac{h}{B}.$$

Let also a', b' , be the principal semi-diameters which coincide with the axes to which the equation

$$A''x^2 + B''y^2 + h = 0$$

appertains; and we similarly have

$$a'^2 = -\frac{h}{A''}, \quad b'^2 = -\frac{h}{B''}.$$

Hence, as $A'' + B'' = A + B$, we have

$$\frac{1}{x'^2} + \frac{1}{y'^2} = \frac{1}{a'^2} + \frac{1}{b'^2};$$

That is, the sum of the reciprocals of the squares of any two semi-diameters of a curve of the second order, which are perpendicular to each other, is the same; and, in reference to the general equation, is =

$$-\frac{A+B}{h} = -\frac{A+B}{G}(4AB - C^2).$$

(93.) When $4AB - C^2 = 0$, we have (55) x', y' , both of them *infinite*, which shews the centre of the curve to be infinitely remote from the origin. It becomes hence necessary to consider this case separately.

$$\text{Let } Ax^2 + By^2 + Cxy + ax + by + c = 0$$

be the general equation, in which $4AB - C^2 = 0$.

Then, transferring the origin to a point $x'y'$, the corresponding equation (76) is

$$\begin{aligned} & Ax^2 + By^2 + Cxy \\ & + (2Ax' + Cy' + a)x + (2By' + Cx' + b)y \\ & + (Ax'^2 + By'^2 + Cx'y' + ax' + by' + c) = 0. \end{aligned}$$

Let $x'y'$ determine some point in the curve, so that

$$Ax'^2 + By'^2 + Cx'y' + ax' + by' + c = 0,$$

and the equation becomes

$$Ax^2 + By^2 + Cxy \\ + (2Ax' + Cy' + a)x + (2By' + Cx' + b)y = 0.$$

But, since $4AB - C^2 = 0$, and $\therefore C = 2\sqrt{AB}$, we have

$$Ax^2 + By^2 + Cxy = (x\sqrt{A} + y\sqrt{B})^2.$$

Hence the reduced equation is equivalent to

$$(x\sqrt{A} + y\sqrt{B})^2 \\ + (2Ax' + Cy' + a)x + (2By' + Cx' + b)y = 0.$$

(94.) We shall now, as in article 79, transfer this equation to two other rectangular axes proceeding from the same origin, and making an angle, ω , with the former; and, (eq. 4, p. 36), putting $x\cos\omega - y\sin\omega$ and $x\sin\omega + y\cos\omega$ for x and y , the resulting equation is

$$\{(\cos\omega\sqrt{A} + \sin\omega\sqrt{B})x - (\sin\omega\sqrt{A} - \cos\omega\sqrt{B})y\}^2 \\ + \{(2Ax' + Cy' + a)\cos\omega + (2By' + Cx' + b)\sin\omega\}x \\ - \{(2Ax' + Cy' + a)\sin\omega - (2By' + Cx' + b)\cos\omega\}y = 0.$$

Let ω satisfy the condition

$$\cos\omega\sqrt{A} + \sin\omega\sqrt{B} = 0,$$

which will give

$$\tan\omega = -\sqrt{\frac{A}{B}}, \quad \cos\omega = -\sqrt{\frac{B}{A+B}}, \quad \sin\omega = \sqrt{\frac{A}{A+B}},$$

and thence

$$\sin\omega\sqrt{A} - \cos\omega\sqrt{B} = \sqrt{A+B}; \\ (2Ax' + Cy' + a)\cos\omega + (2By' + Cx' + b)\sin\omega \\ = -\frac{a\sqrt{B} - b\sqrt{A}}{\sqrt{A+B}},$$

$$\text{and } (2Ax' + Cy' + a)\sin\omega - (2By' + Cx' + b)\cos\omega =$$

$$2x' \sqrt{A(A+B)} + 2y' \sqrt{B(A+B)} + \frac{a \sqrt{A} + b \sqrt{B}}{\sqrt{A+B}}$$

$$= 2 \sqrt{A+B} \left\{ x' \sqrt{A} + y' \sqrt{B} + \frac{a \sqrt{A} + b \sqrt{B}}{2(A+B)} \right\}.$$

The equation thus becomes

$$(A+B)y^2 - \frac{a \sqrt{B} - b \sqrt{A}}{\sqrt{A+B}} x$$

$$- 2 \sqrt{A+B} \left\{ x' \sqrt{A} + y' \sqrt{B} + \frac{a \sqrt{A} + b \sqrt{B}}{2(A+B)} \right\} y = 0.$$

(95.) We have (93) assumed $x'y'$ to determine a point in the curve, but not restricted ourselves to any particular point; we may, therefore, take this point where the curve is intersected by a straight line whose equation is

$$x \sqrt{A} + y \sqrt{B} + \frac{a \sqrt{A} + b \sqrt{B}}{2(A+B)} = 0,$$

by means of which we shall have

$$x' \sqrt{A} + y' \sqrt{B} + \frac{a \sqrt{A} + b \sqrt{B}}{2(A+B)} = 0,$$

which reduces the equation to

$$(A+B)y^2 - \frac{a \sqrt{B} - b \sqrt{A}}{\sqrt{A+B}} \cdot x = 0,$$

$$\text{or } y^2 - \frac{a \sqrt{B} - b \sqrt{A}}{(A+B)^{\frac{1}{2}}} \cdot x = 0 \dots\dots (c).$$

But the equation of a parabola, whose parameter is p , taking the origin at the vertex and the principal axis for the axis of x , is

$$y^2 = px, \text{ or } y^2 - px = 0.$$

Hence the following particulars:—

(96.) 1st. When $a\sqrt{B} - b\sqrt{A} \neq 0$, the locus is a *parabola* whose parameter is equal to

$$\frac{a\sqrt{B} - b\sqrt{A}}{(A + B)^{\frac{1}{2}}}.$$

(97.) 2nd. According to article 8, the equation

$$x\sqrt{A} + y\sqrt{B} + \frac{a\sqrt{A} + b\sqrt{B}}{2(A + B)} = 0$$

defines a straight line inclined to the original axis of x at an angle whose tangent $= -\sqrt{\frac{A}{B}}$, and which is therefore equal to ω , the inclination to the axis of the curve with the axis of x ; this line (95) also passing through the vertex $x'y'$, it must coincide with the axis of the curve. Therefore, the above equation properly represents the *principal diameter* of the curve; by uniting it with the original equation, we may hence find the co-ordinates $x'y'$ of its intersection with the curve, or the vertex.

(98.) 3rd. If $a\sqrt{B} - b\sqrt{A} = 0$, or $a\sqrt{B} = b\sqrt{A}$, the equation (c) gives simply

$$y = 0,$$

which shews the locus in this case to be a straight line corresponding with the new axis of x , the equation of which is given (97).

(99.) 4th. The equation $4AB - C^2 = 0$, giving $C^2 = \pm 2\sqrt{AB}$, the values of the constants A, B , must have the same sign to make C real; that is, they must be either both of them positive or both negative; and hence we may consider them both positive, for, when negative, they can be made so by preliminarily changing all the signs of the original equation. If, under this consideration, C be negative we shall have $C = -2\sqrt{AB}$ instead of $+2\sqrt{AB}$; in this case, the foregoing operations hold good by either substituting $-\sqrt{A}$ instead of \sqrt{A} or $-\sqrt{B}$ for \sqrt{B} , or by considering

either \sqrt{A} or \sqrt{B} to have a negative value; and $\tan \omega$ will become hence $= + \sqrt{\frac{A}{B}}$ instead of $- \sqrt{\frac{A}{B}}$.

Thus we see that, when C is *negative*, $\tan \omega$ is *positive*, and $\omega < \frac{\pi}{2}$; and that, when C is *positive*, $\tan \omega$ is *negative* and

$$\therefore \omega < \frac{\pi}{2}.$$

The foregoing investigations lead immediately to the solutions of the three following propositions.

(100.) *To express the equations of the principal diameters of a curve of the second order, which is determined by the general equation.*

The co-ordinates of the centre (85) are

$$x' = \frac{C b - 2 B a}{4 A B - C^2}, y' = \frac{C a - 2 A b}{4 A B - C^2}.$$

Let ω denote the inclination of one of the principal diameters of the curve with the co-ordinate axis of x ; and (84)

$$\tan 2\omega = \frac{C}{A - B},$$

from which

$$\tan \omega = \frac{\sec 2\omega - 1}{\tan 2\omega} = \sqrt{\frac{\{(A - B)^2 + C^2\} - (A - B)}{C}}.$$

Now the diameter being inclined to the co-ordinate axis of x at the angle ω , and also passing through the centre $x' y'$ of the curve, its equation (13) is

$$y - y' = (x - x') \tan \omega.$$

Hence, by substitution, we have

$$y - \frac{C a - 2 A b}{4 A B - C^2} = \left(x - \frac{C b - 2 B a}{4 A B - C^2} \right) \cdot \sqrt{\frac{\{(A - B)^2 + C^2\} - (A - B)}{C}} \dots (x),$$

for the equation of one of the principal diameters.

The other diameter passing through the centre $x'y'$ perpendicular to this, its equation (15) is

$$y - \frac{Ca - 2Ab}{4AB - C^2} = - \left(x - \frac{Cb - 2Ba}{4AB - C^2} \right) \cdot \sqrt{\frac{C}{\{(A-B)^2 + C^2\} - (A-B)}},$$

or, which is the same,

$$y - \frac{Ca - 2Ab}{4AB - C^2} = - \left(x - \frac{Cb - 2Ba}{4AB - C^2} \right) \cdot \sqrt{\frac{\{(A-B)^2 + C^2\} + (A-B)}{C}} \dots (y).^4$$

(101.) *Cor. 1.* If the origin of the ordinates be the centre of the curve, its equation (86) will be of the form

$$Ax^2 + By^2 + Cxy + c = 0;$$

and we shall have $a = 0$, $b = 0$. In this case, therefore, the equations of the principal diameters are

$$y = \sqrt{\frac{\{(A-B)^2 + C^2\} - (A-B)}{C}} \cdot x,$$

$$\text{and } y = - \sqrt{\frac{\{(A-B)^2 + C^2\} + (A-B)}{C}} \cdot x.$$

(102.) *Note.* The equation

$$\tan 2\omega = \frac{C}{A-B}$$

* By uniting these equations of the principal diameters with the given equation of the curve we may thence find the positions of the vertices.

applies equally to both diameters. For, if 2ω fulfil this equation, it will also hold good when $2\omega \pm \pi$ is substituted; and ω denoting the inclination of one of the diameters, $\omega \pm \frac{\pi}{2}$ will evidently be that of the other.

From this equation we derive generally

$$\tan \omega = \frac{\sec 2\omega - 1}{\tan 2\omega} = \pm \sqrt{\frac{\{(A-B)^2 + C^2\} - (A-B)}{C}},$$

the upper sign appertaining to one of the axes, and the under sign to the other.

Thus, by making use of the under sign, the equation (x) will become the same as the equation (y), and *vice versa* —, because

$$-\sqrt{\frac{\{(A-B)^2 + C^2\} - (A-B)}{C}} \cdot \sqrt{\frac{\{(A-B)^2 + C^2\} - (A-B)}{C}} = -1.$$

When $4AB - C^2 = 0$, see article 97.

(103.) *The equation of a curve of the second order being given to find the values of its principal semi-diameters.*

The squares of the semi-diameters are (83) equal to

$$\frac{\pm G}{A''(4AB - C^2)}, \quad \frac{\pm G}{B''(4AB - C^2)},$$

wherein (84)

$$G = Cab - Ab^2 - Ba^2 + c(4AB - C^2);$$

$$A'' = A \cos^2 \omega + B \sin^2 \omega + C \cos \omega \sin \omega,$$

$$B'' = A \sin^2 \omega + B \cos^2 \omega - C \cos \omega \sin \omega,$$

$$\text{and } \tan 2\omega = \frac{C}{A-B}.$$

From the last we deduce

$$\cos^2 \omega = \frac{1}{2} \left(1 + \frac{1}{\sec 2\omega} \right) = \frac{1}{2} \left(1 + \frac{A - B}{\sqrt{\{(A - B)^2 + C^2\}}} \right),$$

$$\sin^2 \omega = \frac{1}{2} \left(1 - \frac{1}{\sec 2\omega} \right) = \frac{1}{2} \left(1 - \frac{A - B}{\sqrt{\{(A - B)^2 + C^2\}}} \right),$$

$$\cos \omega \sin \omega = \frac{C}{2 \sqrt{\{(A - B)^2 + C^2\}}};$$

and hence we get

$$A'' = \frac{A + B + \sqrt{\{(A - B)^2 + C^2\}}}{2},$$

$$B'' = \frac{A + B - \sqrt{\{(A - B)^2 + C^2\}}}{2}.$$

These and the foregoing value of G substituted, the squares of the principal semi-diameters of the curve are found equal to

$$\begin{aligned} & \pm \frac{2 \{C a b - A b^2 - B a^2 + c (4 A B - C^2)\}}{(4 A B - C^2) [A + B + \sqrt{\{(A - B)^2 + C^2\}}]}, \\ & \pm \frac{2 \{C a b^2 - A b^3 - B a^2 + c (4 A B - C^2)\}}{(4 A B - C^2) [A + B - \sqrt{\{(A - B)^2 + C^2\}}]}, \end{aligned}$$

the under sign being for the ellipse and either sign for the hyperbola (83).

(104.) When the origin is at the centre of the curve (91) $a = 0$, $b = 0$; and therefore, in this case, the squares of the principal semi-diameters are equal to

$$\frac{\pm 2 c}{A + B + \sqrt{\{(A - B)^2 + C^2\}}}, \frac{\pm 2 c}{A + B - \sqrt{\{(A - B)^2 + C^2\}}}.$$

(105.) *To determine the particular description of a curve of the second order from the immediate relative values of the constants which belong to its equation.*

In (81), (82), and the subsequent articles, the different cases are severally stated, throughout the various relations of A'' , B'' , G , $\pm AB - C^2$, &c., where A'' , B'' , are (84) expressed in terms of the coefficients A , B , C , by means of the arc ω as a subsidiary. It is hence only necessary to transfer the relations of A'' , B'' , to those of the immediate coefficients A , B , C , which may be easily effected from their values which have already been found (103), viz.:

$$A'' = \frac{A + B + \sqrt{\{(A - B)^2 + C^2\}}}{2},$$

$$B'' = \frac{A + B - \sqrt{\{(A - B)^2 + C^2\}}}{2}.$$

Thus it is evident that, when $(A + B)^2$ is greater than $(A - B)^2 + C^2$, the sign of $A + B$ cannot be affected with either the addition or subtraction of $\sqrt{\{(A - B)^2 + C^2\}}$, and, consequently, that the values of A'' , B'' , will both have the same sign with $A + B$. But, when $(A + B)^2$ is greater than $(A - B)^2 + C^2$, we shall have

$$(A + B)^2 - \{(A - B)^2 + C^2\} = 4AB - C^2 \text{ positive.}$$

Hence, when $4AB - C^2$ is *positive*, A'' and B'' will both of them have the same sign with $A + B$, that is, they will both be positive when $A + B$ is positive, and both negative when $A + B$ is so.

It is also pretty obvious that, when $(A + B)^2$ is less than $(A - B)^2 + C^2$, the values of A'' , B'' , will have different signs, that is, the one will be positive and the other negative.

In this case we shall have

$$(A + B)^2 - \{(A - B)^2 + C^2\} = 4AB - C^2 \text{ negative.}$$

Thus we see, when $4AB - C^2$ is *negative*, that A'' , B'' , are of different signs*.

Again, under the class $4AB - C^2 = 0$, when the value of $a\sqrt{B} - b\sqrt{A} = 0$, we shall have

$$2\sqrt{A}(a\sqrt{B} - b\sqrt{A}) = 2a\sqrt{AB} - 2Ab = 0,$$

$$\text{or } Ca - 2Ab = 0.$$

Hence, also, when $a\sqrt{B} - b\sqrt{A}$ not $= 0$, we shall have

$$Ca - 2Ab \text{ not } = 0.$$

By carefully comparing these relations with the articles (51), (52), (58), (59), (60), (66), and (68), we find the different descriptions of the curve to be as in the following arrangement, wherein

$$G = Cab - Ab^2 - Ba^2 + c(4AB - C^2).$$

* These relations are also pretty evident from the equations

$$A'' + B'' = A + B,$$

$$4A''B'' = 4AB - C^2.$$

When $4AB - C^2$ is positive, and $A + B$ and G are of different signs, the locus is an ELLIPSE.

| | | | | | |
|-----------------|-------------|----------------------------|------------------------|---|------------------|
| " | " | " | are of the same signs, | " | IMPOSSIBLE. |
| " | " | and $A + B$ and $G = 0$, | " | " | a POINT. |
| " | is negative | " and G not $= 0$, | " | " | an HYPERBOLA. |
| " | " | " $G = 0$, | " | " | a STRAIGHT LINE. |
| $4AB - C^2 = 0$ | " | and $Ca - 2Ab$ not $= 0$, | " | " | a PARABOLA. |
| " | " | $Ca - 2Ab = 0$, | " | " | a STRAIGHT LINE. |

Note.—When the origin of the axes is at the centre of the curve, and, consequently, the equation is

$$Ax^2 + By^2 + Cxy + c = 0, \text{ the value of } G \text{ is simply } c(4AB - C^2).$$

(106.) *Parabola related to its Conjugate Diameters.*

Substitute for x and y respectively, the values

$$a + x \cos \alpha + y \cos \alpha', \text{ and } b + x \sin \alpha + y \sin \alpha';$$

in the equation $y^2 = px$, p being the parameter, and we have

$$\begin{aligned} & b^2 + (x \sin \alpha + y \sin \alpha')^2 + 2b(x \sin \alpha + y \sin \alpha') \\ &= b^2 + x^2 \sin^2 \alpha + y^2 \sin^2 \alpha' + 2xy \sin \alpha \sin \alpha' + 2bx \sin \alpha \\ & \quad + 2by \sin \alpha' = p(a + x \cos \alpha + y \cos \alpha'), \text{ or} \end{aligned}$$

$$\left. \begin{aligned} & y^2 \sin^2 \alpha' + x^2 \sin^2 \alpha + 2xy \sin \alpha \sin \alpha' + b^2 - ap \\ & + (2b \sin \alpha - p \cos \alpha)x + (2b \sin \alpha' - p \cos \alpha')y \end{aligned} \right\} = 0 \dots (1).$$


That this equation may be of the form $y^2 = kx$, the following conditions must obtain

$$\sin \alpha \sin \alpha' = 0, \quad \sin^2 \alpha = 0, \quad 2b \sin \alpha' - p \cos \alpha' = 0,$$

$$\text{and } b^2 - ap = 0 \dots \dots \dots (2);$$

when these are fulfilled the equation becomes

$$y^2 = \frac{p}{\sin^2 \alpha'} \cdot x \dots \dots \dots (3).$$



As the second of the above conditions establishes the first, it follows that, to determine α , α' , a , b , we have only three distinct equations. Thus the number of systems of axes with respect to which the equation preserves the above form is infinite. The relation $\sin \alpha = 0$ shows, besides, that the angle contained by the old and new axis of $x = 0$, or the new axis of x , is parallel to the principal axis. Therefore, in the parabola *all the diameters are parallel to the principal axis*.

In the second place $b^2 - ap = 0$, being that which $y^2 - px$, or $y^2 - px = 0$, becomes when we replace x and y , for the co-ordinates, a , b , of the new origin, we conclude that this origin is on the curve. In giving to a any arbitrary value, we find from the equation $b^2 - pa = 0$, the corresponding

value of b , and the point A' determined by these values represents the new origin.

Lastly, the equation $2b \sin \alpha' - p \cos \alpha' = 0$, gives

$$\tan \alpha' = \frac{p}{2b} \dots\dots\dots (4),$$

which is an expression similar to $a = \frac{p}{2y''}$, which has been found for the tangent to a parabola, which shews that *the new axis of y is a tangent to the curve.*

By equation (4),

$$\tan^2 \alpha' = \frac{p^2}{4b^2}, \quad \therefore \cos^2 \alpha' = \frac{4b^2}{4b^2 + p^2},$$

$$\sin^2 \alpha' = \tan^2 \alpha' \cos^2 \alpha' = \frac{p^2}{4b^2} \cdot \frac{4b^2}{4b^2 + p^2}$$

$$= \frac{p^2}{4b^2 + p^2} = \frac{p^2}{4pa + p^2} = \frac{p}{4a + p} \text{ (since } b^2 = pa),$$

$$\therefore \frac{p}{\sin^2 \alpha'} = 4a + p = 4 \left(a + \frac{p}{4} \right).$$

Now, if we suppose AG to be the abscissa of the new origin A' related to the old axes, and we draw the radius vector FA' , we know that this radius vector is expressed by $a + \frac{p}{4}$, there-

fore $\frac{p}{\sin^2 \alpha'} = 4AF$, that is, the parameter of the parabola related to a system of conjugate axes is four times the distance of the focus from the new origin*.

Putting p' for the new parameter, we have the equation $y' = p'x$ for the equation of the parabola related to its diameters.

The equation $y^2 = p'x$, or $\frac{y^2}{x} = p'$, shews that, for any system of conjugates, the squares of the ordinates are proportional to the corresponding abscissæ.

* This has been done in a different way at page 89.

Examples.

(1.) Let a circle be described, radius R , about a triangle formed by three tangents to a parabola, whose latus rectum is a , and let x_1, x_2, x_3 , be the rectangular abscissæ of the three points of contact, measured from the directrix, then

$$aR^2 = x_1 x_2 x_3.$$

(2.) Given the base of a triangle and the sum of the tangents of the angles at the base, to determine the locus of the vertex.

(3.) If an ellipse the greatest possible be inscribed within a given triangle, and within the ellipse the greatest triangle, and again within the triangle the greatest ellipse, &c., *ad infinitum*, find the sum of the area of all the triangles and of all the ellipses.

(4.) Given the base of a triangle and the difference of the angles at the base, to determine the locus of the vertex.

(5.) Find the locus of the intersection of the tangent to a parabola, and the perpendicular upon it from the vertex.

(6.) Shew that the locus of the focus of a parabola which shall touch a given straight line, and have a given vertex, is a parabola.

(7.) If any two conic sections have a common focus, their intersections range upon two straight lines passing through the intersection of the directrices.

(8.) Given the base and sum of the sides of a triangle, to find the locus of the centre of the inscribed circle.

(9.) Given the base of a triangle and the sum of the other two sides, to find the locus of the inscribed circle.

(10.) Tangents to a parabola form a given angle with each other; what is the locus of their point of intersection?

(11.) A series of circles being described touching the double branches of the cissoid; the first also touching the directrix, the second the first, the third the second, and so on, it is required to determine the abscissa of the point of contact of the n th circle.

(12.) Let seven points in a conic section be connected in any order by the successive lines $a_1, a_2, a_3 \dots a_7$, forming *any* closed heptagon; let P_{mn} be the intersection of a_m and a_n : then the three pairs of lines $P_{14} P_{37}$ and $P_{15} P_{46}$, $P_{35} P_{46}$ and $P_{26} P_{37}$, $P_{14} P_{35}$ and $P_{15} P_{26}$, will intersect in the same straight line.

(13.) Let

$$y^2 = ax^2 + 2bx + c \dots\dots\dots (1),$$

$$y^2 = a_1x^2 + 2b_1x + c \dots\dots\dots (2),$$

be the equations to two conic sections; then, if

$$(b - b_1)^2 = b^2 - ac \pm 2(b^2 - ac)^{\frac{1}{2}}(b_1^2 - a_1c)^{\frac{1}{2}},$$

an indefinite number of triangles may be inscribed within (1) whose sides shall be tangents to the curve (2).

(14.) If three points be taken in a parabola, the ordinates of which are mp , np , and $-\frac{mn}{m+n} \cdot p$, the circle which circumscribes the triangle formed by tangents at these points, will pass through the vertex and focus: if the ordinates be mp , np , and $-\frac{mn+1}{m+n} \cdot p$, it will pass through the focus and the intersection of the axis and directrix: if mp , np , and $-\frac{mn-1}{m+n} \cdot p$, it will touch the axis at the focus; p being the semi-parameter, and m and n any numbers.

(15.) If two conic sections, whose axes are parallel, intersect one another, the lines joining the points of intersection are equally inclined to the axis.

(16.) Any *seven* points being given, two others can be assigned, such that the *thirty-six* connecting lines of the system shall be tangents *by six together of twelve* conic sections. In how many ways can these points be found?

(17.) A conic section is cut in four points by a circle, and two lines, each passing through two of the points of intersection, are made the axes of co-ordinates, their point of meeting being the origin; shew that the equation to the conic section is of the form $x^2 + bxy + y^2 + dx + ey + f = 0$.

(18.) Describe a circle touching three semicircles, the distance of its centre from the diameter = 2 (its radius).

(19.) The tangents at the extremity of any focal chord are perpendicular to one another.

(20.) One circle touches another internally, and another circle touches both of them; find the locus of the centre of this latter circle.

(21.) In the parabola, if PT and QT be tangents to it at P and Q , respectively, intersecting in T and PQ , a normal at Q , then will the directrix bisect PT .

(22.) The locus of the feet of the perpendiculars, dropped from the focus upon the tangent to a parabola, is the line that touches the parabola at its vertex.

$$y - y' = \frac{2a}{y'} (x - x'),$$

$$\therefore y = \frac{2a}{y'} x + \frac{2ax'}{y'},$$

$$p = -\sqrt{a^2 + ax'}, \quad \therefore p^2 = a^2 + ax'.$$

Let $x = 0$ in the equation, $yy' = 2a(x + x')$,

$$\text{and } yy' = 2ax',$$

$$y = \frac{2ax'}{y'},$$

$$\therefore p^2 = \frac{4a^2x'^2}{y'^2} + a^2 = ax' + a^2, \text{ as before,}$$

\therefore the locus is as above.

(23.) From the vertex of a parabola a straight line is drawn, inclined at an angle of 45° to the tangent at any point; find the equation to the curve which is the locus of their intersection.

Let x', y' , be the co-ordinates of P, then, since it is in the parabola,

$$y'^2 = 4ax' \dots\dots\dots (1),$$

$$\text{equation to the tangent at P is } yy' = 2a(x + x') \dots (2),$$

$$\angle RAT = 45^\circ - RTA,$$

$$\therefore \tan RAT = -\tan RAX = \frac{1 - \frac{2a}{y'}}{1 + \frac{2a}{y'}} = \frac{y' - 2a}{y' + 2a},$$

$$\text{or } \tan RAX = \frac{2a - y'}{2a + y'},$$

\therefore equation to RA , passing through the origin, is

$$y = \frac{2a - y'}{2a + y'} x \dots\dots\dots (3),$$

$$\text{from (2), } y' = 2a \left(\frac{x + x'}{y} \right) \dots\dots\dots (4),$$

and from (3), $2ay + yy' = 2ax - xy'$,

$$\text{or } (x + y)y' = 2a(x - y),$$

$$\therefore y' = 2a \left(\frac{x - y}{x + y} \right) \dots\dots\dots (5),$$

$$\therefore (4) \text{ and } (5) \quad \frac{x + x'}{y} = \frac{x - y}{x + y}, \quad \therefore x' = - \frac{x^2 + y^2}{x + y}.$$

$$\text{By (4), substituting for } x', \quad y' = 2a \left(\frac{x - y}{x + y} \right).$$

Substituting these values of x', y' , in (1), we obtain

$$a(x - y)^2 + (x + y)(x^2 + y^2) = 0, \text{ the equation required.}$$

(24.) Prove that the length of the longer normal to the ellipse is $\sqrt{a^2 + \frac{c^2}{e^2}}$.

(25.) Find the equation to the normal to a parabola under the form $y + 2am = mx - am'$.

(26.) Determine the locus of a point within a plane triangle, so that the sum of the squares of the straight lines drawn from it to the angular points is constant.

(27.) Given the base of a plane triangle and the difference of the angles at the base; find the curve traced by the vertex.

(28.) Two given unequal circles touch one another externally; shew that the locus of the centre of a circle which

always touches the other two is a hyperbola. Find the axes and e , and shew what the hyperbola becomes when the given circles are equal.

(29.) Shew that, by transforming the axes through 30° , the equation

$$ay = 3 \cdot x^2 + 2\sqrt{3} \cdot xy + y$$

belongs to the common parabola.

(30.) If, with the co-ordinates of any point in the quadrant of an ellipse as semi-axes, another quadrant be described, with the same centre, the chord of the former will always be a tangent to the latter.

Let xy be the co-ordinates of P,

XY those of T.

Then equation to the exterior ellipse is

$$y^2 = \frac{b^2}{a^2} (a^2 - x^2) \dots \dots \dots (1),$$

and $Y^2 = \frac{y^2}{x^2} (x^2 - X^2)$ is equation to the interior ;

substituting from (1) this becomes

$$Y^2 = \frac{b^2}{a^2} \cdot \frac{a^2 - x^2}{x^2} (x^2 - X^2) \dots \dots \dots (2);$$

equation to A B is

$$Y - b = -\frac{b}{a} X,$$

$$\therefore Y = b - \frac{b}{a} X, \quad X = \frac{b}{a} (a - X) \dots \dots \dots (3);$$

combining (3) with (2)

$$(a - X)^2 = \frac{a^2 - x^2}{x^2} (x^2 - X^2)$$

$$x^4 - 2aXx^2 + a^2X^2 = 0,$$

$$(x^2 - aX)^2 = 0,$$

$$x^2 - aX = 0,$$

$$X = \frac{x^2}{a}.$$

Hence the chord meets the quadrant in only one part, and therefore a tangent to it.

(31.) The side of an equilateral hexagon inscribed in an ellipse eccentricity e , with two sides parallel to the axis major, is to the side of one inscribed in a circle on major axis $\therefore 4 - 2e^2 : 4 - e^2$.

Let AP be the side of the hexagon in ellipse, then x and y being the co-ordinates of P, and AC = a .

$$\begin{aligned} AP &= \sqrt{(a-x)^2 + y^2} = PQ, \\ &= 2x, \end{aligned}$$

$$\therefore (a-x)^2 + y^2 = 4x^2,$$

$$y^2 = 4x^2 - (a-x)^2 = (1-e^2)(a^2 - x^2) \text{ by the ellipse,}$$

$$(4-e^2)x^2 + 2ax = (2+e^2)a^2,$$

$$x^2 + \frac{2a}{4-e^2}x = \frac{2-e^2}{4-e^2}a^2,$$

$$x^2 + \frac{2a}{4-e^2}x + \frac{a^2}{(4-e^2)^2} = \left\{ \frac{2-e^2}{4-e^2} + \frac{1}{(4-e^2)^2} \right\} a^2$$

$$= \frac{8-4e^2-2e^2+e^4+1}{(4-e^2)^2} a^2$$

$$= \frac{9-6e^2+e^4}{(4-e^2)^2} a^2,$$

$$\therefore x + \frac{a}{4 - e^2} = \frac{3 - e^2}{4 - e^2} a,$$

$$x = \frac{3 - e^2 - 1}{4 - e^2} a = \left(\frac{2 - e^2}{4 - e^2} \right) a,$$

$$AP = 2x = \text{side of hexagon} = 2a \left(\frac{2 - e^2}{4 - e^2} \right),$$

and side of hexagon in the circle on major axis = a ,

$$\therefore AP : a = 4 - 2e^2 : 4 - e^2.$$

(32.) If ξ, ξ' , radii vectores, be at right angles to each other from the centre of an ellipse, $\frac{1}{\xi^2} + \frac{1}{\xi'^2} = \frac{1}{a^2} + \frac{1}{b^2}$.

$$a^2 y^2 + b^2 x^2 = a^2 b^2,$$

$$\therefore \xi^2 = \frac{a^2 b^2}{a^2 \sin^2 \theta + b^2 \cos^2 \theta},$$

$$\text{and } \xi'^2 = \frac{a^2 b^2}{a^2 \cos^2 \theta + b^2 \sin^2 \theta},$$

$$\frac{1}{\xi^2} = \frac{a^2 \sin^2 \theta + b^2 \cos^2 \theta}{a^2 b^2} = \frac{b^2 - (b^2 - a^2) \sin^2 \theta}{a^2 b^2},$$

$$\text{also } \frac{1}{\xi'^2} = \frac{a^2 + (b^2 - a^2) \sin^2 \theta}{a^2 b^2},$$

$$\frac{1}{\xi^2} + \frac{1}{\xi'^2} = \frac{a^2 + b^2}{a^2 b^2} = \frac{1}{a^2} + \frac{1}{b^2}.$$

(33.) If two parabolas have a common axis, a straight line touching the interior, and bounded by the exterior, will be bisected in the point of contact.

(34.) If C be the centre of an ellipse, and in the normal to any point P , PQ be taken equal to the semi-conjugate at P , Q will trace out a circle round C .

(35.) An ellipse has a square described, touching it at the extremities of the minor axis; an ellipse upon the major axis circumscribes this square. This ellipse is dealt with in the same way as before, and these operations are continued till there are $(n + 1)$ ellipses altogether; prove that if the original

eccentricity equals $\sqrt{\frac{n}{n+1}}$ the last ellipse becomes a circle.

The side of the square must evidently be $2b$, and if $b_1 =$ semi-minor axis of the first circumscribing ellipse,

$$\therefore b^2 = \frac{b_1^2}{a^2} \{a^2 - b^2\},$$

$$a^2 b^2 = b_1^2 \{a^2 - b^2\},$$

$$b_1^2 = \frac{a^2 b^2}{a^2 - b^2};$$

$$\text{similarly, } b_2^2 = \frac{a^2 b^2}{a^2 - 2b^2},$$

$$b_n^2 = \frac{a^2 b^2}{a^2 - nb^2}.$$

But if this last ellipse be a circle, $b_n^2 = a^2$,

$$\therefore a^2 - nb^2 = b^2,$$

$$(1 + n) b^2 = a^2,$$

$$\frac{b^2}{a^2} = \frac{1}{1 + n}.$$

$$\text{But } e^2 = 1 - \frac{b^2}{a^2} = 1 - \frac{1}{1 + n} = \frac{n}{1 + n},$$

$$\therefore e = \sqrt{\frac{n}{n+1}}.$$

(36.) If $y^2 - x^2 = a^2$, and $2x'y' = a^2$, represent the same curve referred to rectangular axes with the same origin, find the angle the axis of one system makes with the axis of the other.

Let θ equal the angle required; then, since

$$x = x \cos \theta - y \sin \theta,$$

$$y = x \sin \theta + y \cos \theta;$$

therefore, substituting in the first

$$(x \sin \theta + y \cos \theta)^2 - (x \cos \theta - y \sin \theta)^2 = a^2,$$

$$\text{or } (x^2 - y^2)(\sin^2 \theta - \cos^2 \theta) + 4xy \sin \theta \cos \theta = a^2,$$

and this is to have the form $a^2 = 2xy$,

$$\therefore \sin^2 \theta - \cos^2 \theta = 0, \quad 2 \sin \theta \cos \theta = 1,$$

$$\text{or } \sin^2 \theta = \cos^2 \theta, \quad \text{and } \sin \theta \cdot \cos \theta = \frac{1}{2},$$

$$\therefore \theta = \frac{\pi}{4}.$$

(37.) Shew that a conic section is the locus of a point whose distance from a given point is a linear function of co-ordinate of the former point.

$$r = A + Bx + Cy,$$

$$r^2 = x^2 + y^2 = (A + Bx + Cy)^2,$$

an equation of the second degree; therefore, that to a conic section.

(38.) Shew that the angle between two conjugate diameters in an ellipse can never be acute, and the angle between any two conjugate diameters in the hyperbola can never be obtuse.

If γ be the angle between the conjugate diameters,

$$y = mx,$$

$$y = m'x,$$

$$\tan \gamma = \tan(-\alpha + \beta) = \frac{m' - m}{1 + mm'};$$

and this in the ellipse

$$= \frac{-\left(1 + \frac{b^2}{m^2 a^2}\right)m}{1 - \frac{b^2}{a^2}},$$

and is negative; therefore γ is obtuse.

In the hyperbola, $mm' = \frac{b^2}{a^2}$, therefore, m and m' have the same sign or the conjugate diameters lie in the same quadrant; therefore, the angle between them is never obtuse.

(39.) In the hyperbola, if the straight line joining the points of contact of two tangents which intersect in a point (hk) (external) always pass through the focus, find the locus of the point.

The equation to the line joining the points of contact is

$$a^2 k y - b^2 h x = -a^2 b^2,$$

$$x = ae, \quad y = 0;$$

therefore, substituting $-b^2 h a e = -a^2 b^2$,

$$\text{or } h a e = a^2, \quad h = \frac{a}{e},$$

therefore the locus is the directrix.

(40.) Find the condition that the curve $ax^2 + bxy + cy^2 = d$ may have an asymptote.

(41.) Find the equation to a circle referred to two tangents as co-ordinate axes, and obtain also the equation to the tangent to the circle thus referred.

Let Ax , Ay , be the co-ordinated axes, P any point of the circle whose co-ordinates are x , y , and c = the radius of the circle whose centre is C . Join CP ; then

$$(x - c)^2 + (y - c)^2 = c^2,$$

$$\text{or, } x^2 + y^2 - 2c(x + y) + c^2 = 0,$$

is the equation to the circle.

Again, from C draw CV parallel to Ax , meeting the tangent PT at the point P , in V and draw PN parallel to Ay , then

$$\angle PTA = \angle CPN, \text{ and } \tan PTA = -\tan PTx,$$

$$\therefore \tan PTx = -\frac{x - c}{y - c},$$

consequently the equation to PT passing through the point $P(x, y)$, is

$$Y - y = -\frac{x - c}{y - c}(X - x),$$

$$\text{or, } (Y - y)(y - c) + (X - x)(x - c) = 0,$$

which is the equation to the tangent.

(42.) In an ellipse, if a be the vectorial angle of the point of contact referred to the focus, the polar equation to the tangent is

$$a(1 - e^2) = r\{e \cos \theta + \cos(\theta - a)\};$$

hence, shew that the straight line joining the focus and the intersection of two tangents bisects the angle between the radii vectores of their points of contact,

$$ASP = a.$$

If x be the abscissa of P , the point of contact from the centre of the ellipse,

$$\left. \begin{array}{l} SP = a - ex \\ \therefore ex = a - SP \end{array} \right\} \therefore ST = \frac{a^2}{x} - ae = \frac{ae \cdot SP}{a - SP}.$$

The equation to a straight line is $\frac{1}{r} = A \cos \theta + B \sin \theta$.

If $\theta = 0$, $A = \frac{1}{ST} = \frac{a - SP}{ae \cdot SP} = (\text{when } \theta = a) \frac{\cos a + e}{a(1 - e^2)}$,

from equation $r = SP = \frac{a(1 - e^2)}{1 + e \cos \theta}$,

$$\text{and } B = \frac{\sin a}{a(1 - e^2)},$$

$$\therefore \frac{1}{r} = \frac{e \cos \theta + \cos(\theta - a)}{a(1 - e^2)}.$$

For the point of intersection of two tangents to the points whose vectorial angles are α, β , we have

$$e \cos \theta + \cos(\theta - a) = e \cos \theta + \cos(\theta - \beta),$$

$$\cos(\theta - a) = \cos(\theta - \beta),$$

$$\theta - a = \beta - \theta.$$

$$\theta = \frac{a + \beta}{2}.$$

(43.) Find the relation among the coefficients in the general equation of the second order, that all diameters may be parallel to one another.

(44.) Let lines be drawn from the vertex of a parabola to the angles of a triangle formed by the intersection of three tangents to the curve, make angles with the axis denoted by $\theta', \theta'', \theta'''$; also let the lines drawn from the vertex to the points of contact make angles with the axis denoted by $\theta_1, \theta_2, \theta_3$. Then

$$\tan \theta' + \tan \theta'' + \tan \theta''' = \tan \theta_1 + \tan \theta_2 + \tan \theta_3.$$

(45.) Construct the curve whose equation is

$$xy - 2y + x - 1 = 0.$$


(46.) Find the position of the curve of which the equation is

$$y^2 - 2xy + 3x^2 + 2y - 4x - 3 = 0.$$

(17.) TP and TP' are two tangents to a parabola, at right angles to each other,

$$TP = b, \text{ and } TP' = c :$$

find the latus rectum



(48.) Given the base of a triangle, and the product of the tangents of the base angles, to find the locus of the vertex.

(49.) Given the base of a triangle, and the difference of the tangents of the angles at the base, to find the locus of the vertex.

(50.) Given the diameter of a parabola, and a tangent through its vertex, to find the locus of the vertex.

(51.) If $(\alpha x + \beta y = \gamma)$ be the equation of any tangent to the ellipse, to determine what relation must hold between α , β , and γ .

(52.) If θ and θ' be the angles which two conjugate semi-diameters r and r' of an ellipse make with the axis major, to shew that

$$\frac{\sin(\theta' + \theta)}{\sin(\theta' - \theta)} = \frac{r^2 - r'^2}{a^2 - b^2}.$$

(53.) The product of the two perpendiculars let fall from any point of the hyperbola upon the asymptotes is invariable.

NEW [AND PERMANENT] LIST OF WORKS

PUBLISHED BY

JOHN WEALE, 59, HIGH HOLBORN, LONDON.

SERIES OF RUDIMENTARY WORKS

FOR THE USE OF BEGINNERS.

1856.

1. CHEMISTRY, by Prof. Fownes, F.R.S., including Agricultural Chemistry, for the use of Farmers. 4th edition 1s.
2. NATURAL PHILOSOPHY, by Charles Tomlinson. 2nd edition . . . 1s.
3. GEOLOGY, by Lieut.-Col. Portlock, F.R.S., &c. 2nd edition . . . 1s. 6d.
- 4, 5. MINERALOGY, by D. Varley, 2 vols. 2nd edition 2s.
6. MECHANICS, by Charles Tomlinson. 2nd edition 1s.
7. ELECTRICITY, by Sir William Snow Harris, F.R.S. 3rd edition . 1s. 6d.
- 8, 9, 10. MAGNETISM, by the same, 3 vols. 3s. 6d.
- 11, 11* ELECTRIC TELEGRAPH, History of the, by E. Highton, C.E., double Part 2s.
12. PNEUMATICS, by Charles Tomlinson. 2nd edition 1s.
- 13, 14, 15. CIVIL ENGINEERING, by Henry Law, C.E., 3 vols.; and 15* Supplement 4s. 6d.
16. ARCHITECTURE (Orders of), by W. H. Leeda. 2nd edition . . . 1s.
17. ARCHITECTURE (Styles of), by T. Bury, Architect. 2nd edition, with additional cuts 1s. 6d.
- 18, 19. ARCHITECTURE (Principles of Design in), by E. L. Garbett, Architect, 2 vols. 2s.
- 20, 21. PERSPECTIVE, by G. Pyne, Artist, 2 vols. 3rd edition . . . 2s.
22. BUILDING, Art of, by E. Dobson, C.E. 2nd edition 1s.
- 23, 24. BRICK-MAKING, TILE-MAKING, &c., Art of, by the same, 2 vols. 2s.
- 25, 26. MASONRY AND STONE-CUTTING, Art of, by the same, with illustrations of the preceding, in 16 4to. atlas plates 2s.
- 27, 28. PAINTING, Art of, or a GRAMMAR OF COLOURING, by George Field, Esq., 2 vols. 2s.
29. DRAINING DISTRICTS AND LANDS, Art of, by G. R. Dempsey, C.E. . 1s.
30. DRAINING AND SEWAGE OF TOWNS AND BUILDINGS, Art of, by G. R. Dempsey, C.E. 1s. 6d.
31. WELL-SINKING AND BORING, Art of, by G. R. Burnell, C.E. 2nd edition 1s.
32. USE OF INSTRUMENTS, Art of the, by J. F. Heather, M.A. 3rd edition 1s.
33. CONSTRUCTING CRANES, Art of, by J. Glynn, F.R.S., C.E. . . . 1s.
34. STEAM ENGINE, Treatise on the, by Dr. Lardner 1s.
35. BLASTING ROCKS AND QUARRYING, AND ON STONE, Art of, by Lieut.-Gen. Sir J. Burgoyne, K.C.B., R.E. 2nd edition . . . 1s.

RUDIMENTARY WORKS.

- 36, 37, 38, 39. **DICTIONARY OF TERMS** used by Architects, Builders, Civil and Mechanical Engineers, Surveyors, Artists, Ship-builders, &c., 4 vols. 4s.
40. **GLASS-STAINING, Art of**, by Dr. M. A. Gessert 1s.
41. **PAINTING ON GLASS, Essay on**, by E. O. Fromberg 1s.
42. **COTTAGE BUILDING, Treatise on**, 2nd edition 1s.
43. **TUBULAR AND GIRDER BRIDGES, and others, Treatise on**, more particularly describing the Britannia and Conway Bridges, with Experiments 1s.
44. **FOUNDATIONS, &c., Treatise on**, by E. Dobson, C.E. 1s.
45. **LIMES, CEMENTS, MORTARS, CONCRETE, MASTICS, &c., Treatise on**, by Geo. R. Burnell, C.E. 1s.
46. **CONSTRUCTING AND REPAIRING COMMON ROADS, Treatise on the Art of**, by H. Law, C.E. 1s.
- 47, 48, 49. **CONSTRUCTION AND ILLUMINATION OF LIGHTHOUSES, Treatise on the**, by Alan Stevenson, C.E., 3 vols. 3s.
50. **LAW OF CONTRACTS FOR WORKS AND SERVICES, Treatise on the**, by David Gibbons, Esq. 1s.
- 51, 52, 53. **NAVAL ARCHITECTURE, Principles of the Science, Treatise on**, by J. Peake, N.A., 3 vols. 3s.
54. **MASTING, MAST-MAKING, AND RIGGING OF SHIPS, Treatise on**, by R. Kipping, N.A. 1s. 6d.
- 55, 56. **NAVIGATION, Treatise on: THE SAILOR'S SEA-BOOK.—How to keep the log and work it off—Latitude and longitude—Great Circle Sailing—Law of Storms and Variable Winds; and an Explanation of Terms used**, with coloured illustrations of Flags, 2 vols. 2s.
- 57, 58. **WARMING AND VENTILATION, Treatise on the Principles of the Art of**, by Chas. Tomlinson, 2 vols. 2s.
59. **STEAM BOILERS, Treatise on**, by R. Armstrong, C.E. 1s.
- 60, 61. **LAND AND ENGINEERING SURVEYING, Treatise on**, by T. Baker, C.E., 2 vols. 2s.
62. **RAILWAY DETAILS, Introductory Sketches of**, by R. M. Stephenson, C.E. 1s.
- 63, 64, 65. **AGRICULTURAL BUILDINGS, Treatise on the Construction of, on Motive Powers, and the Machinery of the Steading; and on Agricultural Field Engines, Machines, and Implements**, by G. H. Andrews, 3 vols. 3s.
66. **CLAY LANDS AND LOAMY SOILS, Treatise on**, by Prof. Donaldson, A.E. 1s.
- 67, 68. **CLOCK AND WATCH-MAKING, AND ON CHURCH CLOCKS, Treatise on**, by E. B. Denison, M.A., 2 vols. 2s.
- 69, 70. **MUSIC, Practical Treatise on**, by C. C. Spencer, 2 vols. 2s.
71. **PIANO-FORTE, Instruction for Playing the**, by the same 1s.
- 72, 73, 74, 75. **RECENT FOSSIL SHELLS, Treatise (A Manual of the Mollusca) on**, by Samuel P. Woodward, and illustrations, 4 vols. 4s.
- 76, 77. **DESCRIPTIVE GEOMETRY, Treatise on**, by J. F. Heather, M.A., 2 vols. 2s.
- 77*. **ECONOMY OF FUEL, Treatise on**, particularly with reference to Reverberatory Furnaces for the Manufacture of Iron and Steam

NEW LIST OF WORKS.

RUDIMENTARY WORKS.

- 78, 79. STEAM AS APPLIED TO GENERAL PURPOSES AND LOCOMOTIVE ENGINES, Treatise on, by J. Sewell, C.E., 2 vols. 2s.
- 79* RUDIMENTARY WORK ON PHOTOGRAPHY, containing full instructions in the Art of producing Photographic Pictures on any material and in any colour; and also Tables of the Composition and Properties of the Chemical Substances used in the several Photographic Processes. By Dr. H. Halleur, of Berlin. Translated from the German, by the advice of Baron A. von Humboldt, by Dr. Strauss 1s.
- 80, 81. MARINE ENGINES, AND ON THE SCREW, &c., Treatise on, by R. Murray, C.E., 2 vols. 2s.
- 80*, 81*. EMBANKING LANDS FROM THE SEA, the Practice of, treated as a Means of Profitable Employment of Capital, by John Wiggins, F.G.S., Land Agent and Surveyor, 2 vols. 2s.
- 82, 82*. POWER OF WATER, AS APPLIED TO DRIVE FLOUR-MILLS, Treatise on the, by Joseph Glynn, F.R.S., C.E. 2s.
83. BOOK-KEEPING, Treatise on, by James Haddon, M.A. 1s.
- 82**, 83*. COAL GAS, Practical Treatise on the Manufacture and Distribution of, by Samuel Hughes, C.E., 3 vols. 3s.
- 83**. CONSTRUCTION OF LOCKS, Treatise on the, with illustrations . . . 1s. 6d.
- 83 bis. PRINCIPLES OF THE FORMS OF SHIPS AND BOATS, by W. Bland, Esq. 1s.
84. ARITHMETIC, Elementary Treatise on, the Theory, and numerous Examples for Practice, and for Self-Examination, by Prof. J. R. Young . . 1s. 6d.
- 84*. KEY to the above, by Prof. J. R. Young 1s. 6d.
85. EQUATIONAL ARITHMETIC: Questions of Interest, Annuities, and General Commerce, by W. Hipsley, Esq. 1s.
- 86, 87. ALGEBRA, Elements of, for the use of Schools and Self-Instruction, by James Haddon, M.A., 2 vols. 2s.
- 88, 89. GEOMETRY, Principles of, by Henry Law, C.E., 2 vols. 2s.
90. GEOMETRY, ANALYTICAL, by James Hann 1s.
- 91, 92. PLANE AND SPHERICAL TRIGONOMETRY, Treatises on, by the same, 2 vols. 2s.
93. MENSURATION, Elements and Practice of, by T. Baker, C.E. 1s.
- 94, 95. LOGARITHMS, Treatise on, and Tables for facilitating Astronomical, Nautical, Trigonometrical, and Logarithmic Calculations, by H. Law, C.E., 2 vols. 2s.
96. POPULAR ASTRONOMY, Elementary Treatise on, by the Rev. Robert Main, M.R.A.S. 1s.
97. STATICS AND DYNAMICS, Principles and Practice of, by T. Baker, C.E. . 1s.
- 98, 98*. MECHANISM, AND PRACTICAL CONSTRUCTION OF MACHINES, Elements of, by the same, 2 vols, 2s.
- 99, 100. NAUTICAL ASTRONOMY AND NAVIGATION, Theory and Practice of, by H. W. Jeans, R.N.C., Portsmouth, 2 vols. 2s.
101. DIFFERENTIAL CALCULUS, by W. S. B. Woolhouse, F.R.A.S. 1s.
102. INTEGRAL CALCULUS, by Homersham Cox, M.A. 1s.
103. INTEGRAL CALCULUS, Collection of Examples of the, by James Hann . . 1s.
104. DIFFERENTIAL CALCULUS, Collection of Examples of the, by J. Haddon, M.A. 1s.
105. ALGEBRA, GEOMETRY, AND TRIGONOMETRY, First Mnemonical Lessons in, by the Rev. Thomas Penington Kirkman, M.A. 1s. 6d.

NEW SERIES OF EDUCATIONAL WORKS;

OR

Volumes intended for Public Instruction and for Reference :

THE public favour with which the Rudimentary Works on scientific subjects have been received induces the Publisher to commence a New Series, somewhat different in character, but which, it is hoped, may be found equally serviceable. The DICTIONARIES of the MODERN LANGUAGES are arranged for facility of reference, so that the English traveller on the Continent and the Foreigner in England may find in them an easy means of communication, although possessing but a slight acquaintance with the respective languages. They will also be found of essential service for the desk in the merchant's office and the counting-house, and more particularly to a numerous class who are anxious to acquire a knowledge of languages so generally used in mercantile and commercial transactions.

The want of small and concise GREEK and LATIN DICTIONARIES has long been felt by the younger students in schools, and by the classical scholar who requires a book that may be carried in the pocket; and it is believed that the present is the first attempt which has been made to offer a complete Lexicon of the Greek Language in so small a compass.

In the volumes on ENGLAND, GREECE and ROME, it is intended to treat of History as a Science, and to present in a connected view an analysis of the large and expensive works of the most highly valued historical writers. The extensive circulation of the preceding Series on the pure and applied Sciences amongst students, practical mechanics, and others, affords conclusive evidence of the desire of our industrious classes to acquire substantial knowledge, when placed within their reach; and this has induced the hope that the volumes on History will be found profitable not only in an intellectual point of view, but, which is of still higher importance, in the social improvement of the people; for without a knowledge of the principles of the English constitution, and of those events which have more especially tended to promote our commercial prosperity and political freedom, it is impossible that a correct judgment can be formed by the mass of the people of the measures best calculated to increase the national welfare, or of the character of men best qualified to represent them in Parliament; and this knowledge becomes indispensable in exact proportion as the elective franchise may be extended and the system of government become more under the influence of public opinion.

The scholastic application of these volumes has not been overlooked, and a comparison of the text with the examinations for degrees given, will show their applicability to the course of historic study pursued in the Universities of Cambridge and London.

- | | | |
|-------|---|---------|
| 1, 2. | CONSTITUTIONAL HISTORY OF ENGLAND, 2 vols., by W. D. Hamilton | 2s. |
| 3, 4. | DOWN TO VICTORIA, by the same | 2s. |
| 5. | OUTLINE OF THE HISTORY OF GREECE, by the same | 1s. |
| 6. | TO ITS BECOMING A ROMAN PROVINCE, by the same | 1s. 6d. |
| 7. | OUTLINE HISTORY OF ROME | 1s. |
| 8. | TO THE DECLINE | 1s. 6d. |

NEW LIST OF WORKS.

RUDIMENTARY WORKS.

- 9, 10. A CHRONOLOGY OF CIVIL AND ECCLESIASTICAL HISTORY, LITERATURE, SCIENCE, AND ART, from the earliest time to a late period, 2 vols., by Edward Law. 2s.
11. GRAMMAR OF THE ENGLISH LANGUAGE, for use in Schools and for Private Instruction, by Hyde Clarke, Esq., D.C.L. 1s.
- 12, 13. DICTIONARY OF THE ENGLISH LANGUAGE. A new and compressed Dictionary of the English Tongue, as Spoken and Written, including above 100,000 words, or 50,000 more than in any existing work, and including 10,000 additional Meanings of Old Words, 2 vols. in 1, by the same 3s. 6d.
14. GRAMMAR OF THE GREEK LANGUAGE, by H. C. Hamilton 1s.
- 15, 16. DICTIONARY OF THE GREEK AND ENGLISH LANGUAGES, by H. R. Hamilton, 2 vols. in 1 2s.
- 17, 18. ——— ENGLISH AND GREEK LANGUAGES, 2 vols. in 1, by the same 2s.
19. GRAMMAR OF THE LATIN LANGUAGE 1s.
- 20, 21. DICTIONARY OF THE LATIN AND ENGLISH LANGUAGES 2s.
- 22, 23. ——— ENGLISH AND LATIN LANGUAGES 1s. 6d.
24. GRAMMAR OF THE FRENCH LANGUAGE, by Dr. Strauss, late Lecturer at Besançon 1s.
25. DICTIONARY OF THE FRENCH AND ENGLISH LANGUAGES, by A. Elwes 1s.
26. ——— ENGLISH AND FRENCH LANGUAGES, by the same 1s. 6d.
27. GRAMMAR OF THE ITALIAN LANGUAGE, by the same 1s.
- 28, 29. DICTIONARY OF THE ITALIAN, ENGLISH, AND FRENCH LANGUAGES, by the same 2s.
- 30, 31. ——— ENGLISH, FRENCH, AND ITALIAN LANGUAGES, by the same 2s.
- 32, 33. ——— FRENCH, ITALIAN, AND ENGLISH LANGUAGES, by the same 2s.
34. GRAMMAR OF THE SPANISH LANGUAGE, by the same 1s.
- 35, 36. DICTIONARY OF THE SPANISH AND ENGLISH LANGUAGES, by the same 2s.
- 37, 38. ——— ENGLISH AND SPANISH LANGUAGES, by the same 2s.
39. GRAMMAR OF THE GERMAN LANGUAGE, by Dr. Strauss 1s.
40. CLASSICAL GERMAN READER, from the best Authors, by the same 1s.
- 41, 42, 43. DICTIONARIES OF THE ENGLISH, GERMAN, AND FRENCH LANGUAGES, by N. E. S. A. HAMILTON, 3 vols. 3s.
- 44, 45. DICTIONARY OF THE HEBREW AND ENGLISH AND HEBREW LANGUAGES, containing all the new Biblical and Rabbinical words, 2 vols. (together with the Grammar, which may be had separately for 1s.), by Dr. Bresslau, Hebrew Professor 7s.

SUPPLEMENTARY TO THE SERIES.

DOMESTIC MEDICINE ; or complete and comprehensive Instructions for Self-aid by simple and efficient Means for the Preservation and Restoration of Health ; originally written by M. Raspail, and now fully translated and adapted to the use of the British public. 1s. 6d.

| | |
|--|---------|
| 7*. GALVANISM, by Sir William Snow Harris | 1s. 6d. |
| 62. RAILWAYS, being vol. ii. | 1s. |
| 75*. Supplement to RECENT and FOSSIL SHELLS | 1s. 6d. |
| 78*. Rudimentary Treatise on LOCOMOTIVE ENGINES | 1s. |
| 79*. Atlas to the preceding; examples of the several Engines in detail, 4to | 3s. 6d. |
| 82***. On WATER WORKS for the Supply of Towns, 3 vols. | 3s. |
| 86, 87*. Key to the ELEMENTS OF ALGEBRA | 1s. 6d. |
| 101*. On WEIGHTS AND MEASURES OF ALL NATIONS | 1s. 6d. |

THE WORK ON

BRIDGES OF STONE, IRON, TIMBER,
AND WIRE.

In 4 Vols., bound in 3, described in the larger Catalogue of Publications; to which the following is the Supplement, now completed, entitled

SUPPLEMENT TO "THE THEORY, PRACTICE, AND
ARCHITECTURE OF BRIDGES OF STONE, IRON,
TIMBER, WIRE, AND SUSPENSION,"

In one large 8vo volume, with explanatory Text and 68 Plates, comprising details and measured dimensions, in Parts as follows:—

| | |
|--------------------|------|
| Part I. | 6s. |
| „ II. | 6s. |
| „ III. | 6s. |
| „ IV. | 10s. |
| „ V. & VI. | 20s. |

Bound in half-morocco, uniform with the larger work, price 2l. 10s., or in a different pattern at the same price.

LIST OF PLATES.

| | |
|--|---|
| Cast-iron girder bridge, Ashford, Rye and Hastings Railway. | Mr. Fairbairn's hollow-girder bridge at Blackburn. |
| Details, ditto. | Waterford and Limerick Railway truss bridge. |
| Elevation and plan of truss of St. Mary's Viaduct, Cheltenham Rail- way. | Hollow-girder bridge over the River Medlock. |
| Iron road bridge over the Railway at Chalk Farm. . | Railway bridge over lagoons of Venice. |

BRIDGES OF STONE, &C.

- Viaduct at Beangency, Orleans and Tours Railway.
- Oblique cast-iron bridge, on the system of M. Polonceau, over the Canal St. Denis.
- Blackwall Extension Railway, Commercial Road bridge.
- Ditto, enlarged elevation of outside girders, with details.
- Ditto, details.
- Ditto, ditto, and sections.
- Ditto, ditto, ditto.
- Richmond and Windsor main line, bridge over the Thames.
- Ditto, details.
- Ditto, ditto, and sections.
- Orleans and Bordeaux Railway bridge.
- Ditto, sections and details.
- Rouen and Havre Railway timber bridge.
- Ditto, details.
- Ditto, ditto, and sections.
- Viaduct of the Valley of Malauncey, near Rouen.
- Hoop-iron suspension bridge over the Seine at Suresne, department de la Seine.
- Hoop-iron suspension foot bridge at Abainville.
- Suspension bridge over the Douro, iron wire suspension cables.
- Ditto, details.
- Glasgow and South-Western Railway bridge over the water of Ayr.
- Ditto, sections and details.
- Plan of the cities of Ofen and Pesth.
- Sections and soundings of the River Danube.
- Longitudinal section of framing.
- No. 1 coffer-dam.
- Transverse framing of coffer-dam.
- Sections of Nos. 2 and 3 of coffer-dam.
- Plan of No. 3 coffer-dam and ice-breakers.
- Plan and elevation of the construction of the scaffolding, and the manner of hoisting the chains.
- Line of soundings,—dam longitudinal sections.
- Dam sections.
- Plan and elevation of the Pesth suspension bridge.
- Elevation of Nos. 2 and 3 coffer-dams.
- End view of ditto.
- Transverse section of No. 2 ditto.
- Transverse section of coffer-dam, plan of the 1st course, and No. 3 pier.
- Vertical section of Nos. 2 and 3 piers, showing vertical bond-stones.
- Vertical cross section of ditto.
- Front elevation of Nos. 2 and 3 piers.
- End elevation of ditto.
- Details of chains.—Ditto.
- Ditto and plan of nut, bolt, and retaining-links.
- Plan and elevation of roller-frames.
- Elevation and section of main blocks for raising the chains.
- Ditto, longitudinal section of fixture pier, showing tunnel for chains.
- Plan and elevation of retaining-plates, showing machine for boring holes for retaining-bars.
- Retaining link and bar.
- Longitudinal plan and elevation of cast-iron beam with truss columns.
- Longitudinal elevation and section of trussing, &c.
- Plan of pier at level of footpath.
- Detail of cantilevers for supporting the balconies round the towers.
- Elevation and section of cantilevers.
- Detail of key-stone & Hungarian arms.
- Front elevation of toll-houses and wing walls.
- Longitudinal elevation of toll-house, fixture pier, wing wall, and pedestal.
- Vertical section of retaining-piers.
- Section at end of fixture pier, showing chain-holes.
- Lamp and pedestal at entrance of bridge.
- Lamp and pedestal at end of wing walls.

Separately sold from the above in a volume, price half-bound in morocco £1. 12s.

An ACCOUNT, with Illustrations, of the SUSPENSION BRIDGE ACROSS the RIVER DANUBE,

BY WILLIAM TIERNEY CLARK, C.E., F.R.S.

With Forty Engravings.

GREAT EXHIBITION BUILDING.

The BUILDING erected in HYDE PARK for the
GREAT EXHIBITION of the WORKS of
INDUSTRY of ALL NATIONS, 1851:

Illustrated by 28 large folding Plates, embracing plans, elevations, sections, and details, laid down to a large scale from the working drawings of the Contractors, Messrs. FOX, HENDERSON, and Co., by CHARLES DOWNES, Architect; with a scientific description by CHARLES COWPER, C.E.

In 4 Parts, royal quarto, now complete, price £1. 10s.,
or in cloth boards, lettered, price £1. 11s. 6d.

. This work has every measured detail so thoroughly made out as to enable the Engineer or Architect to erect a construction of a similar nature, either more or less extensive.

SIR JOHN RENNIE'S WORK

ON THE

**THEORY, FORMATION, AND CONSTRUCTION
OF BRITISH AND FOREIGN HARBOURS.**

Copious explanatory text, illustrated by numerous examples, 2 Vols., very neat
in half-morocco, £18

The history of the most ancient maritime nations affords conclusive evidence of the importance which they attached to the construction of secure and extensive Harbours, as indispensably necessary to the extension of commerce and navigation, and to the successful establishment of colonies in distant parts of the globe.

To this important subject, and more especially with reference to the vast extension of our commerce with foreign nations, the attention of the British Government has of late years been worthily directed; and as this may be reasonably expected to enhance the value of any information which may add to our existing stock of knowledge in a department of Civil Engineering as yet but imperfectly understood, its contribution at the present time may become generally useful to the Engineering Profession.

The Plates are executed by the best mechanical Engravers; the Views finely engraved under the direction of Mr. Pye: all the Engineering Plates have dimensions, with every explanatory detail for professional use.

In octavo, cloth boards, price 9s.

HYDRAULIC FORMULÆ, CO-EFFICIENTS, AND TABLES,

For finding the Discharge of Water from Orifices, Notches, Weirs, Short Tubes, Diaphragms, Mouth-pieces, Pipes, Drains, Streams, and Rivers.

BY JOHN NEVILLE,

ARCHITECT AND C.E., MEMBER ROYAL IRISH ACADEMY, MEMBER INST. C.E. IRELAND, MEMBER GEOLOGICAL SOC. IRELAND, COUNTY SURVEYOR OF LOUTH, AND OF THE COUNTY OF THE TOWN OF DROGHEDA.

This work contains above 150 different hydraulic formulæ (the Continental ones reduced to English measures), and the most extensive and accurate Tables yet published for finding the mean velocity of discharge from triangular, quadrilateral, and circular orifices, pipes, and rivers; with experimental results and co-efficients;—effects of friction; of the velocity of approach; and of curves, bends, contractions, and expansions;—the best form of channel;—THE DRAINAGE EFFECTS OF LONG AND SHORT WEIRS, AND WEIR-BASINS;—extent of back-water from weirs; contracted channels;—catchment basins;—hydrostatic and hydraulic pressure;—water-power, &c.

TREDGOLD ON THE STEAM ENGINE.

Published in 74 Parts, price 2s. 6d. each, in 4to, illustrated by very numerous engravings and wood-cuts, a new and much extended edition, now complete in 3 vols. bound in 4, in elegant half-morocco, price Nine Guineas and a Half.

THE STEAM ENGINE,

IN ITS PROGRESSIVE AND PRESENT STATE OF IMPROVEMENT;

Practically and amply elucidating, in every detail, its modifications and applications, its duties and consumption of fuel, with an investigation of its principles and the proportions of its parts for efficiency and strength; including examples of British and American recently constructed engines, with details, drawn to a large scale.

The well-known and highly appreciated Treatise, Mr. TREDGOLD's national Work on the STEAM ENGINE, founded on scientific principles and compiled from the practice of the best makers—showing also easy rules for construction, and for the calculation of its power in all cases—has commanded a most extensive sale in the several English editions, and in Translations on the Continent. These editions being *now out of print*, the proprietor has been induced to

TREDGOLD ON THE STEAM ENGINE.

enlarge and extend the present edition by practical examples of all kinds, with the most recent improvements in the construction and practical operations of the steam engine both at home and abroad.

The work is divided into the sections named below, either of which may be *purchased separately*: working engineers will be thus enabled to select those portions which more especially apply to the objects upon which they may be respectively employed.

Several scientific men, extensively and practically employed, have contributed original and really practical papers of the utmost utility; by which the value of this extended edition is much increased. A copious INDEX for reference is added.

DIVISION A. Locomotive Engines, 41 plates and 55 wood-cuts, complete, making Vol. I. In half-morocco binding, price £2. 12s. 6d.

DIVISION B. Marine Engines, British and American, numerous plates and wood-cuts, making Vol. II.; bound in 2 vols. half-morocco, price £3. 13s. 6d.

DIVISION C to G. making Vol. III., and completing the work, comprising Stationary Engines, Pumping Engines, Engines for Mills, and several examples of Boilers employed in the British Steam Navy; in half-morocco, price £3. 13s. 6d.

LIST OF PLATES.

DIVISION A.—LOCOMOTIVE ENGINES.

Elevation of the 8-wheeled locomotive engine and tender, the IRON DUKE, on the Great Western Railway.

Longitudinal section of ditto.

Plan, ditto.

Transverse sections, ditto.

Details of ditto: transverse section through working gear, transverse section and end view of tender; plan and section of feed-pump; plan and elevation of hand-pump; details of inside framing, centre axle, driving axle-box, regulation-valve, centre-beam stay, &c.

Elevation of Crampton's patent locomotive engine and tender.

Longitudinal section of ditto.

Plan of ditto.

Transverse sections of ditto.

Elevation of the PYRAMON 6-wheeled goods' engine on the Great Western Railway.

Half-plan of the working gear of ditto.

Elevation of a portion of the working gear of ditto.

Diagrams, by J. Sewell, L. E., of resistances per ton of the train; and portion of engines of the class of the

GREAT BRITAIN locomotive, including tender, with various loads and at various velocities; also of the additional resistance in lbs. per ton of the train, when the engine is loaded, to be added to the resistance per ton of the engine and tender when unloaded.

Side and front elevation of an express carriage engine, introduced on the Eastern Counties Railway by James Samuel, C. E., Resident Engineer.

Longitudinal and cross section of ditto.

Plan of ditto; with plan and section of cylinders, details and sections, piston full size.

Elevation of the outside-cylinder tank engine made by Sharpe Brothers & Co., of Manchester, for the Manchester and Birmingham Railway.

Section of cylinder and other parts, and part elevation of ditto.

Longitudinal section of ditto.

Plan of ditto.

Transverse sections of both ends, with sectional parts.

Mr. Edward Woods' experiments on the several sections of old and modern

TREDGOLD ON THE STEAM ENGINE.

valves of locomotive engines,—viz. fig. 1. stroke commences; fig. 2, steam-port open; fig. 3, steam-port open; fig. 4, steam-port open; fig. 5, stroke completed, steam cut off, exhaustion commences; fig. 6, stroke commences; fig. 7, steam-port full open; fig. 8, steam cut off; fig. 9, exhaustion commences; fig. 10, steam completed.

Ditto, drawn and engraved to half-size: fig. 1, old valve, $\frac{1}{16}$ -inch lap; fig. 2, $\frac{3}{8}$ -inch lap; fig. 3, $\frac{3}{4}$ -inch lap; fig. 4, $\frac{3}{4}$ -inch lap, Gray's patent; fig. 5, 1-inch lap.

Elevation of a six-wheeled locomotive engine and tender, No. 15, constructed by Messrs. Tayleur, Vulcan Foundry, Warrington, for the Caledonian Railway.

Longitudinal section of ditto.

Plan of ditto, engine and tender, with cylindrical part of boiler removed.

Elevations of fire-box, section of fire-box, section of smoke-box, of ditto.

Elevations and sectional parts of ditto.

Sectional parts, half-plan of working gear, ditto.

Elevation of Messrs. Robert Stephenson

and Co.'s six-wheeled patent locomotive engine and tender.

Longitudinal section of ditto.

Plan and details of Stephenson's patent engine.

Section of fire-box, section of smoke-box, front and back elevations of the same.

Plan of a six-wheeled engine on the Birmingham and Shrewsbury Railway, constructed by Messrs. Bury, Curtis, and Kennedy, Liverpool.

Longitudinal section of ditto.

Sectional elevation of the smoke-box, & Sectional elevation of the fire-box of ditto.

Elevation of the locomotive engine and tender, *PLEWS*, adapted for high speeds, constructed by Messrs. R. & W. Hawthorn, of Newcastle-upon-Tyne, for the York, Newcastle, and Berwick Railway Company.

Longitudinal section of ditto. This section is through the fire-box, boiler, and smoke-box, showing the tubes, safety-valve, whistles, steam and blast pipes, &c.

Plan of ditto.

Plan of the working gear, details, &c.

Forty-one plates and fifty-five wood engravings.

DIVISION B.—MARINE ENGINES, &C.

Two plates, comprising figures 1, 2, and 3, Properties of Steam.

Plan of H. M. screw steam frigate *DAUNTLESS*, constructed by Robert Napier, Esq.

Longitudinal elevation and transverse section of ditto.

Longitudinal section at A B on plan, longitudinal section at C D on plan of ditto.

Engines of H. M. steam ship *TERRIBLE*, constructed by Messrs. Maudslay, Sons, and Field, on the double-cylinder principle. Longitudinal sections of engines.

Transverse section and end view of ditto.

Transverse section through boilers of ditto.

Plan of engines, showing also bunkers, paddles, &c.

Oscillating engines of the Peninsular and Oriental Company's steam vessel *ARIEL*, constructed by John Penn and Sons. Longitudinal section.

Section at engines of ditto.

Section at boiler of ditto.

Plan at boiler of ditto.

Section at air-pump, and at cylinder.

Annular cylinder engines of the iron steam vessels *PRINCESS MARY* and *PRINCESS MAUDE*, constructed by Maudslay, Sons, and Field. Longitudinal section.

Transverse section at engines of ditto.

Section at boilers of ditto.

Plan of engines of ditto, showing bunkers, paddles, &c.

Plan of engines of H. M. steam vessel *SIMOOM*, constructed by James Watt & Co., of London and Soho.

TREDGOLD ON THE STEAM ENGINE.

- Longitudinal section of the **SIMOOM**.
 Cross section of ditto.
 Engine of the **RED ROVER**, side view and plan.
 Longitudinal section of ditto.
 Cross sections of ditto.
 Sheer draught and plans of vessel.
 Plan of the engine of H. M. steam frigate **PHOENIX**.
 Longitudinal section of engine of ditto.
 Cross section of ditto.
 Engine of the **RUBY** steam vessel, elevation and plan.
 Sheer draught and plan of vessel.
 Plan of engine of the **WILBERFORCE**, Hull and London packet.
 Cross section of ditto and vessel.
 Longitudinal section of engines of ditto.
 Elevation of engines of ditto.
 Engines of the **BERENICE**, Hon. E. I. Co.'s steam vessel.
 Section of ditto.
 Sheer draught and plan, stern view, and body plan of vessel.
 View of the **BERENICE**, whilst at sea.
 Boilers of H. M. ships **HERMES**, **SPITFIRE**, and **FIREFLY**.
 Kingston's valves, as fitted on board sea-going vessels for blow-off injection, and hand-pump sea valves.
 Boilers of H. M. steam vessel **AFRICAN**.
 Morgan's paddle-wheels, as fitted in H. M. S. **MEDea**.
 Side elevation of ditto.
 Plans of upper and lower decks of ditto.
 Sheer draught and profile of ditto.
 Morgan and Seaward's paddle-wheels, comparatively.
 Positions of a float of a radiating paddle-wheel in a vessel in motion, and positions of a float of a vertically acting wheel in a vessel in motion.
 Cycloidal paddle-wheels.
 Sailing of steamers in five points from courses.
 Experimental steaming and sailing of the **CALEDONIA**, **VANGUARD**, **ASIA**, and **MEDea**.
 Engines of H. M. steam ship **MEGARA**.
 Engine of the steam boat **NEW WORLD**, T. F. Secor & Co., Engineers, New York. Elevation and section.
 Elevations of cylinder and crank ends.
 Steam cylinders, plans, and sections.
 Details.
 Several sections of details.
 Details and sections.
 Details of parts.
 Plans and sections of condenser, bed-plates, air-pump bucket, &c.
 Details and sections, injection valves.
 Details, plan and elevation of beams, &c.
 Details, sections of parts, boilers, &c. of the steam boat **NEW WORLD**.
 Sections, details, and paddles.
 Engines of the U. S. mail steamers **OHIO** and **GEORGIA**. Longitudinal section.
 Elevations and cross sections of ditto.
 Details of steam-chests, side-pipes, valves, and valve gear of ditto.
 Section of valves, and plan of piston of ditto.
 Boilers of ditto, sections of ditto.
 Engine of the U. S. steamer **WATERWITCH**. Sectional elevation.
 Steam-chests and cylinders of ditto.
 Boilers, sections, &c. of ditto.
 Boilers of the U. S. steamer **POWHATAN**. Front view and sections of ditto.
 Elevation of the Pittsburg and Cincinnati American packet **BUCKEYE STATE**.
 Bow view, stern view.
 Plan of the **BUCKEYE STATE**.
 Model, &c. of ditto, wheel-house frame, cross section at wheel-house, and body plan.
 Plan and side elevation of ditto.
 Sheer draught and plan, with the body plan, of the U. S. steam frigate **SARANAC**.
 Longitudinal section of ditto, cross section.
 Engines of the U. S. steamer **SUSQUEHANNA**.
 Elevation of the U. S. **PACIFIC** steam packet engine.
 Plan of ditto.
 Boilers of ditto, end views.
 Ditto ditto.

Eighty-five engravings and fifty-one wood-cuts.

TREDGOLD ON THE STEAM ENGINE.

DIVISION C. TO G., FORMING VOL. III.

STATIONARY ENGINES, PUMPING ENGINES, MARINE BOILERS, &C.

Side elevation of pumping engine, U. S. dock, New York.

End elevation of ditto.

Elevation and section of the pumps, ditto.—2 plates.

Boilers of pumping engines, ditto.

Boilers, Details, &c. of pumping engines, ditto.

Plan of the boilers, ditto.

Isometrical projection of a rectangular boiler.

Plan and two sections of a cylindrical boiler.

Brunton's apparatus for feeding furnace-fires by means of machinery.

Parts of a high-pressure engine with a 4-passaged cock.

Section of a double-acting condensing engine.

Section of a common atmospheric engine.

On the construction of pistons.

Section of steam pipes and valves.

Apparatus for opening and closing steam passages.

Parallel motions.—2 plates.

Plan and elevation of an atmospheric engine.

Elevation of a single-acting Boulton and Watt engine.

Double-acting engine for raising water.

Double-acting engine for impelling machinery.

Maudslay's portable condensing engine for impelling machinery.

Indicator for measuring the force of steam in the cylinder, and diagrams of forms of vessels.

Section of a steam vessel with its boiler, in two parts—diagrams showing fire-places—longitudinal section through boiler and fire-places.

Isometrical projection of a steam-boat engine.

Plan and section of a steam-boat engine.

Ten horse-power engine, constructed by W. Fairbairn and Co.—4 plates.

Forty-five horse-power engine, constructed by W. Fairbairn & Co.—3 plates.

Plan and section of boiler for a 20-horse

engine, at the manufactory of Whitworth & Co., Manchester.

Messrs. Hague's double-acting cylinder, with slides, &c.

Sixty-five-inch cylinder, erected by Maudslay, Sons, and Field, at the Chelsea Water-works.—5 plates.

Beale's patented rotary engine.

Double-story boilers of H.M.S. DEVASTATION, 400 H. P.

Refrigerator feed and brine pumps.

Feed and brine apparatus, as fitted on board the West India Royal Mail Company's ships.

Boilers of H. M. steam sloop BASILISK, 400 H. P.

Boilers of the SINGAPORE, 470 H. P., Peninsular and Oriental Company.

Original double-story boilers of the GREAT WESTERN.

Telescopic chimney, or sliding funnel, of H. M. ship HYDRA, 220 H. P.

Seaward's patent brine and feed valves.

Boilers of H. M. mail packet UNDIKE, (Miller, Ravenhill, & Co.) 100 H. P.

Cross sections of engines of H. M. mail packet UNDIKE.

Longitudinal elevation of ditto.

Brine-pumps as fitted on board H. M. S. MEDEA, 220 H. P. (Maudslay, Sons, and Field.)

Boilers of H. M. S. HYDRA, 220 H. P.

Plan of the four boilers, with the supplementary steam-chests and shut-off valves, of the AVENGER.

Boilers of H. M. steam ship NIGER, 400 H. P., fitted by Maudslay, Sons, and Field.

Experimental boiler, Woolwich Yard.

Boilers of H. M. S. TERRIBLE, 800 H. P. (Maudslay, Sons, and Field.)

Boilers of the MINK and TEASER, 100 H. P. (transferred to WASP.)

Boilers of the SAMSON, 450 H. P.

Daniel's pyrometer, full size.

Boilers of the DESPERATE, 400 H. P. (Maudslay, Sons, and Field.)

Boilers of the NIGER (2nd plate).

Boilers of H. M. S. BASILISK (2nd plate).

Boilers of the UNDIKE.

TREDGOLD ON THE STEAM ENGINE.

- Boilers of the Royal Mail steam ships ASIA and AFRICA, 768 H. P., constructed by R. Napier, Glasgow.
- Longitudinal and midship sections of ditto.
- Boilers of H. M. S. LA HOGUE, 450 H. P. (Seaward & Co.)
- H. M. S. SIDON, 560 H. P. Plan of telescope funnel.
- Boilers of H. M. S. BRISK, 250 H. P.
- Copper boilers for H. M. S. SANS-PAREIL, 350 H. P. (James Watt & Co.)
- American marine boilers, designed and executed by C. W. Copeland, Esq., of New York, as fitted on board the American packets.
- Midship section of the hull of the steam packet PACIFIC, New York and Liverpool line.
- Elevation of pumping engines of the New Orleans Water-works, U. S., arranged and drawn by E. W. Smith, Engineer, constructed at the Allaire Works, New York.
- Elevation of pumps and valves, chests, gearing, &c.
- Elevation at steam cylinder end.
- General plan of a turbine water-wheel in operation at Lowell, Massachusetts, U. S., by J. B. Francis, C. E.
- Elevation of ditto.—Section of ditto.
- Plan of the floats and guide curves, ditto.
- Large self-acting surfacing and screw-propeller lathe, by Joseph Whitworth & Co., Manchester.
- Longitudinal section, showing arrangement of engine-room for disc engine applied to a screw propeller, and Bishop's disc engine, by G. & J. Rennie, with details.
- Arrangement of engine-room foreengines of 60 horse-power, for driving propellers of H. M. steam vessels REYNARD and CRUISER, constructed by Messrs. Rennie. Longitudinal section and engine-room.
- Ditto. Transverse section at boilers and at engines.
- Very elaborate diagrams showing experiments and results of various paddle-wheels.—8 plates.
- Steam flour-mills at Smyrna, constructed by Messrs. Joyce & Co. Double cylinder pendulous condensing engine, side elevation.
- Side elevation, horizontal plan, ditto.
- Longitudinal section.
- Horizontal plan of mill-house and boilers.
- Transverse section through engine-house and mill.
- Boilers, longitudinal and transverse sections, front view.
- Section through mill-stones, elevation of upper part, section of lower part, plan of hopper, &c.

SUMMARY OF THE ILLUSTRATIONS.

| | Plates. | Wood-cuts, |
|--|---------|------------|
| Vol. I. Locomotive Engines | 41 | 55 |
| II. Marine Engines | 85 | 51 |
| III. Stationary Engines, Pumping Engines, Engines for Flour-Mills, Examples of Boilers, &c., &c. . | 100 | 58 |
| Total | 226 | 164 |

FULL-LENGTH PORTRAIT OF
HENRY CAVENDISH, F.R.S.

Some few India paper proofs, before the letters, of this celebrated
Philosopher and Chemist, to be had, price 2s. 6d.

HINTS TO YOUNG ARCHITECTS:

COMPRISING

ADVICE TO THOSE WHO, WHILE YET AT SCHOOL ARE DESTINED
TO THE PROFESSION;
TO SUCH AS, HAVING PASSED THEIR PUPILAGE, ARE ABOUT TO TRAVEL
AND TO THOSE WHO, HAVING COMPLETED THEIR EDUCATION,
ARE ABOUT TO PRACTISE:

TOGETHER WITH

A MODEL SPECIFICATION:

INVOLVING A GREAT VARIETY OF INSTRUCTIVE AND SUGGESTIVE MATTER,
CALCULATED TO FACILITATE THEIR PRACTICAL OPERATIONS;
AND TO DIRECT THEM IN THEIR CONDUCT, AS THE RESPONSIBLE
AGENTS OF THEIR EMPLOYERS,
AND AS THE RIGHTFUL JUDGES OF A CONTRACTOR'S DUTY.

BY GEORGE WIGHTWICK, ARCHITECT.

CONTENTS :—

| | |
|--|---|
| <p>Preliminary Hints to Young Architects on the Knowledge of Drawing. On Serving his Time. On Travelling. His Plate on the Door. Orders, Plan-drawing. On his Taste, Study of Interiors. Interior Arrangements. Warming and Ventilating. House Building, Stabling. Cottages and Villas. Model Specification :— General Clauses. Foundations. Well. Artificial Foundations. Brickwork. Rubble Masonry with Brick Mingled.</p> | <p>Model Specification : Stone-cutting. —, Grecian or Italian only. —, Gothic only. Miscellaneous. Slating. Tiling. Plaster and Cement-work. Carpenters' Work. Joiners' Work. Iron and Metal-work. Plumbers' Work. Drainage. Well-digging. Artificial Levels, Concrete, Foundations, Piling and Planking, Paving, Vaulting, Bell-hanging, Plumbing, and Building generally.</p> |
|--|---|

Extra cloth boards. price 8s.

THE ENGINEER'S AND CONTRACTOR'S POCKET BOOK,

WITH AN ASTRONOMICAL ALMANACK,

REVISED FOR 1855-6. In morocco tuck, price 6s.

CONTENTS.

- AIR**, Air in motion (or wind), and wind-mills.
Alloys for bronze; Miscellaneous alloys and compositions; Table of alloys; Alloys of copper and zinc, and of copper and tin.
Almanack for 1855 and 1856.
American railroads; steam vessels.
Areas of the segments of a circle.
Armstrong (R.), his experiment on boilers.
Astronomical phenomena.
Ballasting.
Barlow's (Mr.) experiments.
Barrel drains and culverts.
Bell-hanger's prices.
Blowing a blast engine.
Boilers and engines, proportions of; Furnaces and chimneys; Marine.
Bossut's experiments on the discharge of water by horizontal conduit or conducting pipes.
Brass, weight of a lineal foot of, round and square.
Breen (Hugh), his almanack.
Bricks.
Bridges and viaducts; Bridges of brick and stone; Iron bridges; Timber bridges.
Burt's (Mr.) agency for the sale of preserved timber.
Cask and malt gauging.
Cast-iron binders or joints; Columns, formulæ of; Columns or cylinders, Table of diameter of; Hollow columns, Table of the diameters and thickness of metal of; Girders, prices of; Stanchions, Table of, strength of.
Chairs, tables, weights, &c.
Chathurn limestone.
Chimneys, &c., dimensions of.
Circumferences, &c. of circles.
Coal, evaporating power of, and results of coking.
Columns, cast-iron, weight or pressure of, strength of.
Comparative values between the present and former measures of capacity.
Continuous bearing.
Copper pipes, Table of the weight of, Table of the bore and weight of cocks for.
Copper, weight of a lineal foot of, round and square.
Cornish pumping engines.
Cotton mill; Cotton press.
Current coin of the principal commercial countries, with their weight and relative value in British money.
Digging, well-sinking, &c.
Docks, dry, at Greenock.
Draining by steam power.
Dredging machinery.
Dwarf, Table of experiments with H. M. screw steam tender.
Earthwork and embankments, Tables of contents, &c.
Experiments on rectangular bars of malleable iron, by Mr. Barlow; on angle and T iron bars.
Fairbairn (Wm.), on the expansive action of steam, and a new construction of expansion valves for condensing steam engines.
Feet reduced to links and decimals.
Fire-proof flooring.
Flour-mills.
Fluids in motion.
Francis (J. B., of Lowell, Massachusetts), his water-wheel.
French measures.
Friction.
Fuel, boilers, furnaces, &c.
Furnaces and boilers.
Galvanized tin iron sheets in London or Liverpool, list of gauges and weights of.
Gas-tubing composition.
Glynn (Joseph), F. R. S., on turbine water-wheels.
Hawksby (Mr., of Nottingham), his experiments on pumping water.
Heat, Tables of the effects of.

THE ENGINEER'S AND CONTRACTOR'S POCKET BOOK.

- Hexagon heads and nuts for bolts, proportional sizes and weights of.
 Hick's rule for calculating the strength of shafts.
 Hodgkinson's (Eaton) experiments.
 Hungerford Bridge.
 Hydraulics.
 Hydrodynamics.
 Hydrostatic press.
 Hydrostatics.
 Imperial standard measures of Great Britain; Iron.
 Indian Navy, ships of war, and other vessels.
 Institution of Civil Engineers, List of Members of the, corrected to March 15, 1852.
 Iron balls, weight of cast; bars, angle and T, weight of; castings; experiments; hoop, weight of 10 lineal feet; lock gates; roofs; tubes for locomotive and marine boilers; weights of rolled iron.
 Ironmonger's prices.
 Just's analysis of Mr. Dixon Robinson's limestone.
 Latitudes and longitudes of the principal observatories.
 Lead pipes, Table of the weights of.
 Leslie (J.), C.E.
 Lime, mortar, cements, concrete, &c.
 Limestone, analysis of.
 Liquids in motion.
 Locomotive engines; Table showing the speed of an engine.
 Log for a sea-going steamer, form of.
 Machines and tools, prices of.
 Mahogany, experiments made on the strength of Honduras. [wheels.
 Mallet's experiments on overshot
 Marine boilers; engines.
 Masonry and stone-work.
 Massachusetts railroads.
 Mensuration, epitome of.
 Metals, lineal expansion of.
 Morin's (Col.) experiments.
 Motion; motion of water in rivers.
 Nails, weight and length.
 Navies — of the United States; Indian Navy; Oriental and Peninsular Company; British Navy; of Austria; Denmark; Naples; Spain; France; Germanic Confederation; Holland; Portugal; Prussia; Sardinia; Sweden and Norway; Turkey; Russia
 Royal West India Mail Company's fleet.
 New York, State of, railroads.
 Numbers, Table of the fourth and fifth power of.
 Paddle-wheel steamers.
 Pambour (Count de) and Mr. Parkes' experiments on boilers for the production of steam.
 Peacocke's (R. A.) hydraulic experiments.
 Pile-driving.
 Pitch of wheels. Table to find the diameter of a wheel for a given pitch of teeth.
 Plastering.
 Playfair (Dr. Lyon).
 Preserved timber.
 Prices for railways, paid by H. M. Office of Works; smith and founder's work.
 Prony's experiments.
 Proportions of steam engines and boilers.
 Pumping engines; pumping water by steam power.
 Rails, chairs, &c., Table of.
 Railway, American, statistics; railway and building contractor's prices; carriages.
 Rain, Tables of.
 Rammell's (T. W.) plan and estimate for a distributing apparatus by fixed pipes and hydrants.
 Rennie's (Mr. Geo.) experiments; (the late J.) estimate.
 Roads, experiments upon carriages travelling on ordinary roads; influence of the diameter of the wheels; Morin's experiments on the traction of carriages, and the destructive effects which they produce upon roads.
 Robinson (Dixon), his experiments and material.
 Roofs; covering of roofs.
 Ropes, Morin's recent experiments on the stiffness of ropes; tarred ropes; dry white ropes.
 Saw-mill.
 Screw steamers.
 Sewage manures.
 Sewers, castings for. their estimates, &c.

THE ENGINEER'S AND CONTRACTOR'S POCKET BOOK.

Signs and abbreviations used in arithmetic and mathematical expressions.

Slating.

Sleepers, quantity in cubic feet, &c.

Smeaton's experiments on wind-mills.

Smith and founder's prices.

Specific gravity, Table of.

Steam dredging; Navigation; Tables of the elastic force; Table of Vessels of war, of America; of England; of India; and of several other maritime nations.

Steel, weight of round steel.

Stone, per lb., stone, qr., cwt., and ton, &c., Table of the price.

Stones.

Strength of columns; Materials of construction,

Sugar-mill.

Suspension aqueduct over the Alleghany River; Bridges over ditto.

Table of experiments with H. M. screw steam tender DWARF; of gradients; iron roofs; latent heats; paddle-wheel steamers of H. M. Service and Post-Office Service; pressure of the wind moving at given velocities; prices of galvanized tinned iron tube; specific heats; the cohesive power of bodies; columns, posts, &c., of timber and iron; the comparative strength, size, weight, and price of iron-wire rope (A. Smith's), hempen rope, and iron chain; corresponding velocities with heads of water as high as 50 ft., in feet and decimals; dimensions of the principal parts of marine engines; effects of heat on different metals; elastic force of steam; expansion and density of water; expansion of solids by increasing the temperature; expansion of water by heat; heights corresponding to different velocities, in French metres; lineal expansion of metals; motion of water, and quantities discharged by pipes of different diameters; power of metals, &c.; pressure, &c., of wind-mill sails; principal dimensions of 28 merchant steamers with screw propellers; of steamers with paddle-wheels; proportion of dilatation of metals by heat

discharge through thin-lipped orifices; quantities of water, in cubic feet, discharged over a weir per minute, hour, &c.; relative weight and strength of ropes and chains; results of experiments on the friction of unctuous surfaces; scantlings of posts of oak; size and weight of iron laths; weight in lbs. required to crush $1\frac{1}{2}$ -inch cubes of stone, and other bodies; weight of a lineal foot of cast-iron pipes, in lbs.; weight of a lineal foot of flat bar iron, in lbs.; weight of a lineal foot of square and round bar iron; weight of a superficial foot of various metals, in lbs.; weight of modules of elasticity of various metals; velocities of paddle-wheels of different diameters, in feet per minute, and British statute miles, per hour; the dimensions, cost, and price per cubic yard, of ten of the principal bridges or viaducts built for railways; the height of the boiling point at different heights;—to find the diameter of a wheel for a given pitch of teeth, &c.

Tables of squares, cubes, square and cube roots.

Teeth of wheels.

Temperature, the relative indications of, by different thermometers.

Thermometers, Table of comparison of different.

Timber for carpentry and joinery purposes; Table of the properties of different kinds of.

Tin plates, Table of the weight of.

Tools and machines, prices of.

Traction, Morin's experiments on.

Tredgold's Rules for Hydraulics, from Eytelwein's Equation.

Turbines, Report on, by Joseph Glynn and others.

Values of different materials.

Water-wheels.

Watson's (H. H.) analysis of limestone from the quarries at Chatburn.

Weight of angle and T iron bars; of woods.

Weights and measures.

West India Royal Mail Company.

Whitelaw's experiments on turbine

THE ENGINEER'S AND CONTRACTOR'S POCKET BOOK.

White's (Mr., of Cowes) experiments on Honduras mahogany.

Wicksteed's (Thos.) experiments on the evaporating power of different kinds of coal.

Wind-mills; of air, air in motion, &c. Woods.

Wrought iron, prices of.

Zinc as a material for use in house-building.

In one Volume 8vo, extra cloth, bound, price 9s.

THE STUDENT'S GUIDE TO THE PRACTICE OF DESIGNING, MEASURING, AND VALUING ARTIFICERS' WORKS;

Containing Directions for taking Dimensions, abstracting the same, and bringing the Quantities into Bill; with Tables of Constants, and copious memoranda for the Valuation of Labour and Materials in the respective trades of Bricklayer and Slater, Carpenter and Joiner, Sawyer, Stonemason, Plasterer, Smith and Ironmonger, Plumber, Painter and Glazier, Paper-hanger. Thirty-eight plates and wood-cuts.

The Measuring, &c., edited by EDWARD DOBSON, Architect and Surveyor. Second Edition, with the additions on Design by E. LACY GARBETT, Architect.

CONTENTS.

PRELIMINARY OBSERVATIONS ON DESIGNING ARTIFICERS' WORKS.

Preliminary Observations on Measurement, Valuation, &c.—On measuring—On rotation therein—On abstracting quantities—On valuation—On the use of constants of labour.

BRICKLAYER AND SLATER.

DESIGN OF BRICKWORK—technical terms, &c.

Foundations—Arches, inverted and erect—Window and other aperture heads—Window jambs—Plates and internal cornices—String-courses—External cornices—Chimney shafts—On general improvement of brick architecture, especially fenestration.

MEASUREMENT.

Of diggers' work—Of brickwork, of facings, &c.

DESIGN OF TILING, and technical terms.

Measurement of Tiling—Example of the mode of keeping the measuring-

ABSTRACTING Bricklayers' and Tilers' work.

Example of bill of Bricklayers' and Tilers' work.

VALUATION of Bricklayers' work, Earthwork, Concrete, &c.

Table of sizes and weights of various articles—Tables of the numbers of bricks or tiles in various works—Valuation of Diggers' and Bricklayers' labour—Table of Constants for said labour.

EXAMPLES OF VALUING.

1. A yard of concrete.—2. A rod of brickwork.—3. A foot of facing.—4. A yard of paving.—5. A square of tiling.

DESIGN, MEASUREMENT, AND VALUATION OF SLATING.

CARPENTER AND JOINER.

DESIGN OF CARPENTRY—technical terms, &c.

Breastsummers, an abuse: substitutes for them—Joists, trimmers,

DESIGNING, MEASURING, AND VALUING ARTIFICERS' WORKS.

and right use—Substitutes for girders and quarter-partitions—Quarter-partitions—Roof-framing—Great waste in present common modes of roof-framing—To determine the right mode of subdividing the weight, and the right numbers of bearers for leaded roofs—The same for other roofs—Principle of the truss—Considerations that determine its right pitch—Internal filling or *tracery* of trusses—Collar-beam trusses—Connection of the parts of trusses—Variations on the truss; right limits thereto—To avoid fallacious trussing and roof-framing—Delorme's roofing; its economy on circular plans—Useful property of regular polygonal plans—On combinations of roofing, hips, and valleys—On gutters, their use and abuse—Mansarde or curb-roofs.

DESIGN OF JOINERY—technical terms, &c.

Modes of finishing and decorating panel-work—Design of doors.

MEASUREMENT of Carpenters' and Joiners' work—Abbreviations.

Modes of measuring Carpenters' work—Classification of labour when measured with the timber—Classification of labour and nails when measured separately from the timber.

EXAMPLES OF MEASUREMENT, arch centerings.

Bracketing to sham entablatures, gutters, sound-boarding, chimney-grounds, sham plinths, sham pilasters, floor-boarding, mouldings—Doorcases, doors, doorway linings—Dado or surbase, its best construction—Sashes and sash-frames (examples of measurement)—Shutters, boxings, and other window fittings—Staircases and their fittings.

ABSTRACTING Carpenters' and Joiners' work.

Example of Bill of Carpenters' and Joiners' work.

VALUATION of Carpenters' and Joiners' work, Memoranda.

Tables of numbers and weights.

TABLES OF CONSTANTS OF LABOUR.

Roofs, naked floors—Quarter-par-

titions—Labour on fir, per foot cube—Example of the valuation of deal; or battens—Constants of labour on deals, per foot superficial.

CONSTANTS OF LABOUR, and of nails, separately.

On battening, weather boarding—Rough boarding, deal floors, batten floors.

LABOUR AND NAILS together.

On grounds, skirtings, gutters, doorway-linings—Doors, framed partitions, mouldings—Window-fittings—Shutters, sashes and frames, staircases—Staircase fittings, wall-strings—Dados, sham columns and pilasters.

VALUATION OF SAWYERS' WORK.

MASON.

DESIGN OF STONEMASONS' WORK.

Dr. Robison on Greek and Gothic Architecture—Great fallacy in the Gothic ornamentation, which led also to the modern 'monkey styles'—'Restoration' and Preservation.

MEASUREMENT of Stonemason's work.

Example of measuring a spandril step, three methods—Allowance for labour not seen in finished stone—Abbreviations—Specimen of the measuring-book—Stairs—Landings—Steps—Coping—String-courses—Plinths, window-sills, curbs—Columns, entablatures, blockings—Cornices, renaissance niches.

ABSTRACTING AND VALUATION.

Table of weight of stone—Table of Constants of Labour—Example of Bill of Masons' work.

PLASTERER.

DESIGN OF PLASTER-WORK in real and mock Architecture.

Ceilings and their uses—Unnecessary disease and death traced to their misconstruction—Sanitary requirements for a right ceiling—Conditions to be observed to render domestic ceilings innocuous—Ditto, for ceilings of public buildings—Barbarous shifts necessitated by wrong ceiling—Technical terms in Plasterers' work.

MEASUREMENT of Plaster-work.

DESIGNING, MEASURING, AND VALUING ARTIFICERS' WORKS.

Abbreviations — Abstracting of Plasterers' work — Example of Bill of Plasterers' work.

VALUATION.

Memoranda of quantities of materials — Constants of Labour.

SMITH AND FOUNDER.**ON THE USE OF METAL-WORK IN ARCHITECTURE.**

Iron not rightly to be used much more now than in the middle ages — Substitutes for the present extravagant use of iron — Fire-proof (and sanitary) ceiling and flooring — Fire-proof roof-framing in brick and iron — Another method, applicable to hipped roofs — A mode of untrussed roof-framing in iron only — A principle for iron trussed roofing on any plan or scale — Another variation thereof — On the decoration of metallic architecture.

MEASUREMENT of Smiths' and Founders' work.

PLUMBER, PAINTER, GLAZIER, &c.**DESIGN, &c. OF LEAD-WORK.****MEASUREMENT OF PAINT-WORK — Abbreviations.**

Specimen of the measuring-book — Abstract of Paint-work — Example of Bill of Paint-work.

VALUATION of Paint-work.

Constants of Labour — Measurement and Valuation of GLAZING — Measurement and Valuation of PAPER-HANGING.

APPENDIX ON WARMING.

Modifications of sanitary construction to suit the English open fire — More economic modes of warming in public buildings — Ditto, for private ones — Warming by gas.

In 12mo., price 5s. bound and lettered,

THE OPERATIVE MECHANIC'S WORKSHOP COMPANION, AND THE SCIENTIFIC GENTLEMAN'S PRACTICAL ASSISTANT;

comprising a great variety of the most useful Rules in Mechanical Science, divested of mathematical complexity; with numerous Tables of Practical Data and Calculated Results, for facilitating Mechanical and Commercial Transactions.

BY W. TEMPLETON,

AUTHOR OF SEVERAL SCIENTIFIC WORKS.

Third edition, with the addition of Mechanical Tables for the use of Operative Smiths, Millwrights, and Engineers; and practical directions for the Smelting of Metallic Ores.

2 vols. 4to, price £ 2. 16s.,

CARPENTRY AND JOINERY;

containing 190 Plates; a work suitable to Carpenters and Builders, comprising Elementary and Practical Carpentry, useful to Artificers in the Colonies.

THE AIDE-MÉMOIRE TO THE MILITARY SCIENCES,

Framed from Contributions of Officers of the different Services, and edited by a Committee of the Corps of Royal Engineers. The work is now completed.

Sold in 3 vols. £4. 10s., extra cloth boards and lettered, or in 6 Parts, as follows :

| | | £. | s. | d. |
|---------|---------------------------------|---------|----|----|
| Part I. | A. to D., NEW EDITION | 0 | 14 | 0 |
| II. | D. to F. | 0 | 16 | 0 |
| III. | F. to M. | 0 | 16 | 0 |
| IV. | M. to P. | 0 | 14 | 0 |
| V. | P. to R. | 0 | 16 | 0 |
| VI. | R. to Z. | 1 | 0 | 0 |
| | | £4 16 0 | | |

In 1 large Volume, with numerous Tables, Engravings, and Cuts,

A TEXT BOOK

For Agents, Estate Agents, Stewards, and Private Gentlemen, generally, in connection with Valuing, Surveying, Building, Letting and Leasing, Setting out, disposing, and particularly describing all kinds of Property, whether it be Land or Personal Property. Useful to

| | | |
|----------------|---------------------|--------------------|
| Auctioneers | Assurance Companies | Landed Proprietors |
| Appraisers | Builders | Stewards |
| Agriculturists | Civil Engineers | Surveyors |
| Architects | Estate Agents | Valuers, &c. |

In 1 vol. large 8vo, with 13 Plates, price One Guinea, in half-morocco binding,

MATHEMATICS FOR PRACTICAL MEN :

Being a Common-Place Book of PURE AND MIXED MATHEMATICS; together with the Elementary Principles of Engineering; designed chiefly for the use of Civil Engineers, Architects, and Surveyors.

BY OLINTHUS GREGORY, LL.D., F.R.A.S.

Third Edition, revised and enlarged by HENRY LAW, Civil Engineer.

CONTENTS.

PART I.—PURE MATHEMATICS.

CHAPTER I.—ARITHMETIC.

| SECT. | | SECT. | |
|-------|----------------------------------|-------|--|
| 1. | Definitions and notation. | 5. | Division of whole numbers.— Proof of the first four rules of Arithmetic. |
| 2. | Addition of whole numbers. | 6. | Vulgar fractions.—Reduction of vulgar fractions.—Addition and |
| 3. | Subtraction of whole numbers. | | |
| 4. | Multiplication of whole numbers. | | |

MATHEMATICS FOR PRACTICAL MEN.

subtraction of vulgar fractions.
— Multiplication and division of vulgar fractions.

Decimal fractions.—Reduction of decimals.—Addition and subtraction of decimals.—Multiplication and division of decimals.

Complex fractions used in the arts and commerce.—Reduction.—Addition.—Subtraction and multiplication.—Division.—Duodecimals.

Powers and roots.—Evolution.

Proportion.—Rule of Three.—Determination of ratios.

Logarithmic arithmetic.—Use of the Tables.—Multiplication and division by logarithms.—Proportion, or the Rule of Three, by logarithms.—Evolution and involution by logarithms.

Properties of numbers.

CHAPTER II.—ALGEBRA.

Definitions and notation.—2. Addition and subtraction.—3. Multiplication.—4. Division.—5. Involution.—6. Evolution.—7. Surds.—Reduction.—Addition, subtraction, and multiplication.—Division, involution, and evolution.—8. Simple equations.—Extermination.—Solution of general problems.—9. Quadratic equations.—10. Equations in general.—11. Progression.—Arithmetical progression.—Geometrical progression.—12. Fractional and negative exponents.—13. Logarithms.—14. Computation of formulæ.

CHAPTER III.—GEOMETRY.

Definitions.—2. Of angles, and right lines, and their rectangles.

PART II.—MIXED MATHEMATICS.

CHAPTER I.—MECHANICS IN GENERAL.

CHAPTER II.—STATICS.

Statistical equilibrium.

Centre of gravity.

General application of the principles of statics to the equilibrium

SECT.

—3. Of triangles.—4. Of quadrilaterals and polygons.—5. Of the circle, and inscribed and circumscribed figures.—6. Of plans and solids.—7. Practical geometry.

CHAPTER IV.—MENSURATION.

1. Weights and measures.—1. Measures of length.—2. Measures of surface.—3. Measures of solidity and capacity.—4. Measures of weight.—5. Angular measure.—6. Measure of time.—Comparison of English and French weights and measures.

2. Mensuration of superficies.

3. Mensuration of solids.

CHAPTER V.—TRIGONOMETRY.

1. Definitions and trigonometrical formulæ.—2. Trigonometrical Tables.—3. General propositions.—4. Solution of the cases of plane triangles.—Right-angled plane triangles.—5. On the application of trigonometry to measuring heights and distances.—Determination of heights and distances by approximate mechanical methods.

CHAPTER VI.—CONIC SECTIONS.

1. Definitions.—2. Properties of the ellipse.—3. Properties of the hyperbola.—4. Properties of the parabola.

CHAPTER VII.—PROPERTIES OF CURVES.

1. Definitions.—2. The conchoid.—3. The cissoid.—4. The cycloid and epicycloid.—5. The quadratrix.—6. The catenary.—Relations of Catenarian Curves.

of structures.—Equilibrium of piers or abutments.—Pressure of earth against walls.—Thickness of walls.—Equilibrium of polygons.—Stability of arches.—Equilibrium of suspension bridges.

24 JOHN WEALE'S NEW LIST OF WORKS.

MATHEMATICS FOR PRACTICAL MEN.

SECT.

CHAPTER III.—DYNAMICS.

1. General Definitions.
2. On the general laws of uniform and variable motion.—Motion uniformly accelerated.—Motion of bodies under the action of gravity.—Motion over a fixed pulley, and on inclined planes.
3. Motions about a fixed centre, or axis.—Centres of oscillation and percussion.—Simple and compound pendulums.—Centre of gyration, and the principles of rotation.—Central forces.
4. Percussion or collision of bodies in motion.
5. Mechanical powers. — Levers. — Wheel & axle. — Pulley. — Inclined plane.—Wedge and screw.

CHAPTER IV.—HYDROSTATICS.

1. General Definitions.—2. Pressure and equilibrium of Non-elastic Fluids.—3. Floating Bodies.—4. Specific gravities. — 5. On capillary attraction.

CHAPTER V.—HYDRODYNAMICS.

1. Motion and effluence of liquids.
2. Motion of water in conduit pipes and open canals, over weirs, &c.—Velocities of rivers.
3. Contrivances to measure the velocity of running waters.

CHAPTER VI.—PNEUMATICS.

1. Weight and equilibrium of air and elastic fluids.
2. Machines for raising water by the pressure of the atmosphere.
3. Force of the wind.

SECT.

CHAPTER VII.—MECHANICAL AGENTS.

1. Water as a mechanical agent.
2. Air as a mechanical agent.—Columb's experiments.
3. Mechanical agents depending on heat. The Steam Engine Table of Pressure and Temperature of Steam.—General description of the mode of action of the steam engine.—Theory of the same. — Description of various engines, and formulae calculating their power: practical application.
4. Animal strength as a mechanical agent.

CHAPTER VIII.—STRENGTH OF MATERIALS.

1. Results of experiments, and principles upon which they should be practically applied.
2. Strength of materials to resist tensile and crushing strain. Strength of columns.
3. Elasticity and elongation of bodies subjected to a crushing or compressive strain.
4. On the strength of materials subjected to a transverse strain. Longitudinal form of beams of uniform strength.—Transverse strength of other materials of cast iron. — The strength of beams according to the manner in which the load is distributed.
5. Elasticity of bodies subjected to transverse strain.
6. Strength of materials to resist torsion.

APPENDIX.

- I. Table of Logarithmic Differences.
- II. Table of Logarithms of Numbers, from 1 to 100.
- III. Table of Logarithms of Numbers, from 100 to 10,000.
- IV. Table of Logarithmic Sines, Tangents, Secants, &c.
- V. Table of Useful Factors, extending to several places of Decimals.
- VI. Table of various Useful Numbers, with their Logarithms.
- VII. Table of Diameters, Areas, and Circumferences of Circles, &c.
- VIII. Table of Relations of the Arc, Abscissa, Ordinate and Subnormal, in the Catenary.
- IX. Tables of the Lengths and Vibrations of Pendulums.
- X. Table of Specific Gravities.
- XI. Table of Weight of Materials frequently employed in Construction.
- XII. Principles of Chronometers.
- XIII. Select Mechanical Expedients.
- XIV. Observations on the Effect of Old London Bridge on the Tides, &c.
- XV. Professor Farish on Isometrical Perspective.

