660503

THE

GRIFFIN'S AID-DE-CAMP

BY

BLUNT SPURS.

EDITED BY A PROFESSIONAL MAN

WITH AN ADDENDUM

CONTAINING

HINTS ON THE EDUCATION OF THE HORSE

AND

THE VETERINARY PHARMACOPŒIA.

SIXTH EDITION.

MADRAS:
HIGGINBOTHAM AND CO.,
By Appointment in India to H. R. H. the Prince of Wales, and to the Madras University.
THACKER SPINK AND CO., CALCUTTA: THACKER AND CO., BOMBAY.
1882.

MADRAS:
PRINTED BY HIGGINBOTHAM AND CO.,
165, MOUNT ROAD.

TO

MAJOR OUTRAM, C.B.;

RESIDENT AT THE COURT OF HYDERABAD, IN SINDE,
ETC., ETC.

As an admirer of your public talents and private virtues, as well as of your justly celebrated character as one of the best hog-hunters on the Bombay side of India, but as no admirer of the neglect in your stable, this volume is dedicated by your much obliged friend,

BLUNT SPURS.

'The style of a man may be known by his horse."



THE IMPOSSIBLE HORSE.

PREFACE.

THE publication of a new edition of this very useful and greatly appreciated work calls for no remark, beyond noting that the size and style of printing the book have been altered to make it more compact and handy; and that "*The Veterinary Pharmacopœia*," taken from the *Complete American Farrier*, has been added.

AUTHOR'S PREFACE

(*To the first edition.*)

BROTHERLY love and friendly kindness being generally shown in an inverse ratio in the purchase of a horse to what they are in any other transaction, and the loss sustained from the high price being often ruinous and provokingly vexatious, I publish the following pages, in the hope that you, as well as your noble but ill-treated slave, may mutually derive benefit from the contents.

I have never fallen in with any little pamphlet entirely divested of technical language, such as a tyro would like from its simplicity, giving plain and succint directions how to purchase a horse for a racer, charger, and hunter, together with an easy account of putting him into condition, &c., and showing how far he might venture to treat him when diseased. What is here written has been gained by study from works of others,* and what professional men have

* This acknowledgment was the cause of my book, when in manuscript, being much depreciated, and by one of the very persons who, no doubt, will most benefit by it. "Compile!" said he, "any body can compile." Now, discoveries of this kind are of a *green*, and also of a jealous hue. How are our ideas confirmed, and how is our information got, if not from books? All amateur writings must of necessity be more or less compiled;

taught me, interspersed throughout with my own observations. Proceeding from an undiploma'd hand, these are not supposed to be infallible; but though errors may exist, they would not lead far out of the right path, and perhaps will aid a novice when buying, and afterwards tend to prevent him ruining his horse by injudicious treatment, more than if he pounced on a scientific veterinary volume. These learned books are but ill-adapted to a person who has neither had experience, nor given the subject any attention; and they are daily misunderstood, even in their most simple parts.

If I am occasionally a little censorious, remember, this is the best mode of impressing things on the memory; and if you are of opinion that the contents are very generally known, I can only say, that it is a great pity they are not more generally practised: at the same time, I may inform you, this is not written for those who do know, but for those who do not, so that for the future we may all start even.

I beg one favor—that you will not be so incon-

but it is only amateurs who have studied, and reflected, and had practice on such a subject as horse-flesh, that are capable of *translating* scientific and technical English for the understanding of their brethern. Nearly the whole of the last part of this work, to say nothing of much of the preceding parts, will be a compilation, then, if you like, from various professional and other authors; but as I have had some little experience, I reserve to myself the liberty of extracting from those whose opinions approach nearest my own.

sistent as to deride the contents of this little treatise, and at the same time adopt the precepts it contains; and moreover, if you do, take care you never invite me to inspect your stud, or I will "show up" more faults in each horse than you thought existed in the whole of the Bomb Proof.

To enable you, condescending reader, who may read for instruction, to purchase a casty, sound, and useful Arab; to keep him in condition always fit for work; to take your gallop out of him without injury, or to bring him on the turf; and to prevent your inflicting unnecessary pain when illness overtakes him, is the object of

THE GRIFFIN'S AID-DE-CAMP.

EDITOR'S PREFACE

(*To edition published, May* 1858.)

THE following pages have been revised and corrected by a PROFESSIONAL MAN. In a work, however, so generally correct from its author's hands, there has not been much to amend. The errors corrected were principally such as a non-professional man was likely to fall into. The Reviser has had a very extensive practice in India, extending over a period of eight and twenty years, and his position in the service of Government has afforded him the opportunity of making the peculiarities of Arabs and other horses his constant study during that long period :—while at the same time it enables him to say, that with but few exceptions in India, more horses have passed through his hands professionally than those of any other man.

He has, therefore, been able to make a few corrections in the text regarding the peculiarities of horses generally used in India, and their proper management; also to notice characteristics which might be unobserved, by even professional men, with less wide experience.

This little work is now strongly recommended by the Reviser, as richly meriting the attention of all possessors of horses, (private gentlemen as well the Profession) who take an interest in their comfort and health;—to promote and secure which many useful lessons will be found in its pages.

D. S.

PUBLISHERS' PREFACES.

(*To fourth edition, March* 1863.)

In issuing the Fourth Edition of the Griffin's Aid-de-camp, the publisher feels thankful that his attempts to draw attention to this valuable book, and to increase its usefulness by additional notes and an improvement in the text, have been most fully appreciated by the public, as has been proved by the rapid sale of more than one edition. In all the Presidencies it has been accepted as a trustworthy guide for the special treatment of horses in this country. What Youatt and other writers have done for England, Blunt Spurs, and his editors, have more than equally well done for India, so that the young Officer and Civilian may, immediately on his arrival, obtain a sure, and as far as such a thing can be, infallible mentor for the purchase and treatment of his horses.

The present edition has been carefully revised and brought down to the most scientific views of the present day, by one of the most experienced Veterinary Surgeons in India, and certainly the chief authority in Southern India. Where different modes of treatment have been recommended by various

authors, both are clearly exhibited. Those who possess copies of previous editions will find that the new matter, concerning the more common ailments of this noble animal, will more than repay the cost of another copy.

The editor has thrown into the text, between brackets, the more important additions. At page 10 for example, will be found a concise account of the causes, effects, and appearance of the various kind of spavin, illustrated by several diagrams. Splint is treated at considerable length on page 28, with a brief view of the necessary treatment. By the aid of the letter press and wood-cut, the merest tyro will be able easily to detect and obviate this common disease. The great folly of bleeding, the usual specific for all equine ailments, is clearly shown on and after page 85, together with the course of medicine that ought to take the place of the fleam and the lancet. Many other subjects are taken up with equal ability and clearness, and being usually illustrated by reference to cases that have occurred in Madras, are most valuable to the unprofessional man. Amongst these are castration, bishoping, diseases of the ligaments of the legs, shoeing, sandcrack, ring-bone, &c. At the end of the volume will be found an entirely new chapter on the various kinds of lameness, together with the simplest and best treatment to be adopted in each case. In an Appendix will be found a com-

plete Veterinary Materia Medica, with the appearance, properties, uses, and European and vernacular names (Hindostani and Tamil) in the Roman character, of the various drugs. In short, every thing has been done to make the volume a perfect handbook upon the subject, equally suitable both to the amateur and the tyro.

The Frontispiece is a Chromo-lithograph of the Horse's leg, foot, and hoof, showing every ligament, bone, and tendon, upon which the book treats. The publisher has no hesitation in asserting that this production of Mr. Barren is the finest piece of lithography ever performed in India.

(*To the fifth edition, April* 1868.)

In issuing this Edition the Publishers have introduced a chapter on Pegu Ponies, a desideratum long felt to make this work complete, and which has been very kindly contributed by an Officer of high standing in the Madras Army, possessing great experience on the subject, under the nomenclature of "Vagrant."

In lieu of the Chromo-lithograph of the "Horse's Leg," the Publishers have substituted in this Edition the "Muscles of the Horse," which they trust will prove acceptable.

CONTENTS.

PAGE.

PREFACE v
AUTHOR'S PREFACE TO THE FIRST EDITION vi
EDITOR'S PREFACE TO EDITION PUBLISHED, MAY 1858 ... ix
PUBLISHERS' PREFACE TO FOURTH EDITION, MARCH 1863. xi
PREFACE TO THE FIFTH EDITION, APRIL, 1868 xiv

PART I.

PURCHASING FROM THE STABLES, ETC. 1
The Caste ... ... *ib.*
The Build, commencing with the Quarters, Tail, Thighs, Stifle, &c. 6
The Hocks and the Hind Parts, proof of half the Breeding 8
Bone-spavin 9
Spavins 11
Bog-spavin—Thorough-pin 14
The Head, Nostrils, Mouth, Lips, Eyes, Ears, &c. ... 16
The Neck 18
The Withers and Back 19
The Shoulder and Chest 22
The Depth of Girth and Carcase 23
The Elbow and Forearm, &c. 24
The Knee and Back Sinews, and Suspensory Ligament, The Shank-bone, Fetlocks, &c. 25
The Shank 26
Splint 28
The Fetlock 29
Suspensory Ligament 30
Windgalls 31
The Pasterns, &c.... 32
The Ringbone 33
Position of theLegs 35
The Feet 36
The Testicles, &c.... 42

PAGE.

PURCHASING FROM THE STABLES.—(*continued.*)
Gelding 43
The Colour 44
Arriving Thin 46
Bad Condition on Landing, &c. 47
Manner of Going 48
Price ... ... *ib.*
Advice 51
The Griffin's Catechism 55
Racer 56
Charger 58
Hunter 60
Description of various Horses 62
Breeding 63
Premier 69
Perfection 70
Water Proof ... ... *ib.*
Outline 71

PART II.

THE AGE 72
The Colt's Grinders ... ... *ib.*
The Colt's Nippers 73
The Tushes 74
Marks that show the Age ... ... *ib.*
Remarks 75
Bishoping 78
THE PULSE 79
BLEEDING 81
PHYSICKING 87
To make Bran Mash, and Bran Tea 94
To make Fine Gram, or Wheat flour Gruel ... ... *ib.*
Muzzle 95
Drenching 96
Physic Ball and Drench ... ... *ib.*
Alteratives 97
General Alteratives 99

PAGE.

SHOEING 100
Cutting the Hoof 104
Three-quarter Shoes and Tips 107

PART III.

PUTTING INTO CONDITION 111
Stabling and Clothing... ... ... *ib.*
The Feet, Physic, &c. 114
Hours of Feeding and Watering 118
Grooming, Washing, &c. 120
Getting out of Order 123
Gora-Wallas Riding 124
Danger of Over-feeding 125
Mixed Boiled Food, Chaff, &c. 126
Native remedies for Improving, Fattening and giving a Horse an Appetite... 130
Exercise 131
How to keep Condition 132
Change from Stabling to the open Air, Hunting, &c. ... 135
Massallah Balls 140
Cordial Drinks ... ... *ib.*
Beer Tonic 141
Ball ... ... *ib.*
Native Massallah Ball... 143
TRAINING 144
Stable and Clothing 148
Stable Management 149
Difference of Form, &c. 150
Time required to Train ... ... *ib.*
A Day's Routine 151
Training Weight, Physic, Exercise, &c. 156
Trials 160
The Grain, Grass and Water 162
Quantity of Food, and Hours of Feeding ... 166
Muzzle, when required, Leather Dhool, &c. 169
Cordial Balls, Horse-bread, &c. 171

PAGE

TRAINING.—(*continued.*)

Bandaging 173
Sweating... 176
Physicking 183
Conclusion 184
Riding, Entering for a Race, and requiting a querulous Visitor 188
Stopping for the Feet 192
Ointment for the Heels, as a Preservative against Cracks *ib.*
Lotion for Tender Shanks 193
Sal Ammoniac Lotion ... ... *ib.*

PART IV.

TREATMENT OF A FEW DISEASES 194
Class I.—Sickness under Physic 196
Over-purging 198
Blister for the Belly 200
Class II.—Gripes 201
Treatment of Gripes 202
Class III.—Cold 207
Sore Throat 211
Influenza... 213
Class IV.—Strangles 215
Swelling of the Glands under the Jaw.. 218
Class V.—Fever ... ... *ib.*
Class VI.—Inflammation of the Lungs 221
Inflammation of the Bowels 226
Inflammation of the Liver, of Spleen... 228
Inflammation of the Stomach 230
Inflammation of the Heart ... ... *ib.*
Inflammation of the Kidneys, the neck of the Bladder, and Bladder 231
Inflammation of the Feet 233
Inflammation Undiscoverable 236
Class VII.—Red Urine 237
Bursautee 238

PAGE.
LAMENESS, &c. 242
Quittor—Thrush—Corns—Sandcrack 246
Pumice Foot 247
Bruises of the Foot ... ... *ib.*
Canker ... ... *ib.*
Navicular Lameness ... ... *ib.*
Injuries from Bad Shoeing... ... ... *ib.*
Punctured Wounds 248
Bog-Spavin, Thorough-pin and Wind-gall ... ... *ib.*
Curb 249
Strains of the Tendons or Back Sinews ... ... *ib.*
Shoulder Lameness ... ... *ib.*
Hip-joint Lameness 250
Dislocation of the Patella ... ... *ib.*
Lameness from Cutting ... ... *ib.*
Rheumatism ... ... *ib.*
WARRANTING 252
A Cheap Horse 261
BURMAH PONIES by "Vagrant" 263

ADDENDUM.

HINTS ON THE EDUCATION OF A HORSE ... ... PAGE. 288
Introduction ... PAGE. 288
The Longe... 290
Mounting 293
Mouthing and Bitting ... 296
Bending Lesson ... 297
"How to teach a Horse to obey the pressure of the Leg" 301
The use of the Spur ... 303
Reining Back 304
Shoulders in 306
Passaging ... ... *ib.*
Paces 307
To Stop a Horse ... 310
Leading ... ... *ib.*
Longeing Whip ... 311
Age and Work ... ... *ib.*
List of Medicines for Veterinary purpose ... 314
THE VETERINARY PHARMACOPŒIA FROM "FORREST'S FARRIER AND HORSE DOCTOR" 323

LIST OF ILLUSTRATIONS.

Chromo-lithograph of the Muscles of the Horse—FRONTISPIECE.

PAGE.

The impossible Horse ... iv
Spavin caused from external injury 11
Sound Hock, Spavined Hock 12
Splint, 3 Varieties 28
Ringbone, Healthy and diseased Bones 34
Outline of the Hoof 37
Premier 69
Perfection 70
Water Proof ... ... *ib.*
Outline of the Horse ... 71
Horse Nippers 78
Shoeing 100
Fore Shoe 101
Hind Shoe 103
Foot Bath 243
Sandcrack 247
Curb and Bandage ... 249
A Cheap Horse 261
Shan Ponies ... 265 and 268

THE

GRIFFIN'S AID-DE-CAMP.

PART I.

PURCHASING FROM THE STABLES, ETC.

The Caste.

High caste, or thorough blood, of course is the first thing to be observed in selecting an Arab for the turf, for without good caste it is all labour in vain; but as there are different breeds, each possessing high caste, it is sometimes difficult to decide which is the better bred out of three or four that happen also to be all well built; therefore, when it is an option between two of equally supposed good caste, do not hesitate, when buying for the turf, to choose that one shining most in running points in preference to the finer or handsomer horse; and these points are not so universally known, and very seldom remembered, if known, when in the act of purchasing. Be quite certain first, that you really are inspecting an Arab, for there are many that are taken

to be of good caste that never saw Arabia, nor sprang from Arab stock ; and though the speed which some few of these have occasionally shown, is proof that they must, by either sire or dam, have inherited good blood of some country, still, they are not to be trusted like the genuine Arab, whose pure descent and unsullied pedigree is generally graced with a shape that rarely or never deceives. There is a beautiful symmetry and harmony of proportion running through all the frame ; a superb quarter, with a high set on, well-carried tail ; a softness and thinness of skin ;* a brilliancy of eye, and an elegant contour of head ; qualities that mark the true blood, and which never exist in the numerous spurious brutes that so abound in the stables.

While these handsome points may be relied on as a sure criterion of caste, there are many that have good caste without showing it externally : these can only be proved on the day of running. Good build of itself in a great measure insures good caste,† never-

* Yet there are many that in the cold weather will wear a rather long coat, and be very sleek in the hot months.

† The three following extracts serve to corroborate this. I think they are also quite sufficient to refute the absurd saying, "No consequence what a horse's build is ; and if he has only blood he'll run." That thorough breeding in a bad formed horse, will generally beat half breeding in a good formed horse, must be granted ; but the slightest superiority of build in any part of the frame between two of equal blood, is known to every stable-boy to be of the greatest importance :—

Osmer, page 221. On form and performance. "For if a different for-

theless, many good caste horses are seen badly formed, and many half or three parts bred, well formed. High caste, united to a running form, will never fail; and some horses, when led out, show such superior blood, energy, and build, that if properly trained, and nothing constitutionally wrong, they would be sure to gallop: others showing *externally* only tolerable good caste, but good make; or good caste but indifferent make, may be called the doubtful ones: while there is a third, a numerous class, which one may with safety declare will never run, and many of these are handsome notwithstanding. But an ugly horse may have as much chance of running as a handsome one, provided the different essential running parts of his frame harmonize; that the joints are all equally corresponding with the size, and that the hind parts agree in relation to the fore parts, for

mation of the parts, &c., and the degrees thereof, be not the cause of difference in the performances, why then one of these horses of the right and true blood would act alike on all ground whatever, and be just as good though made like a hog and without joints." Again, page 229. "In the nature and elegance of their constituent parts, and the due formation thereof, consist the difference between horses of the same and different countries; or betwixt blood, and no blood." Again, page 230. "Conqueror and Othello were two full brothers, but one was a king and the other a beggar, with respect both to form and action. If then the difference in the performance of these brothers did not depend on their different formation of parts, &c., pray tell me on what it did depend, for the cause of it couldn't be in the blood, unless you will say this innate quality may appertain to one brother and not to another; and then, I apprehend, the bystanders will say you have proved nothing."

"ugliness, though the opposite of beauty, is not the opposite of proportion and fitness,"* and so it is with the horse.

To distinguish high caste appears a very complex affair at the imperial mart of the Bomb Proof; but, produce me an Arab with a straight spine, and a straight long quarter; a muscular and handsome dropped hind leg; a round barrel, swelling well out behind the elbows, with great depth of girth, and a moderately broad but flat chest; a very oblique and deep shoulder; a light neck; a well set on lean head, with a large brilliant eye, thin open nostril, deep mouth, wide clear jowl, and small silky ear;—give me this make, with the upper inner bone of the knees and hocks large, and hinder bone of each also large and the back sinews clear, powerful, and wide away from the suspensory ligament; the whole covered with a fine thin skin, and finishing with small, but well open, short, tough feet, and I'll take the caste for granted. If added to this, he is a long horse; five years old; perfectly fresh; short between the knee and fetlock; fourteen hands two inches high; higher at the croup than at the withers; fine action, and a good constitution, I'll take the speed besides for granted; for when these latter qualities of fine action, and a good constitution, are united to all the

* Burke on the Sublime.

above form, the only remaining desideratum, the nervous excitability,* will then always be found, more or less, associated with it.

[It is very many years since the Bomb Proof in the Fort ceased to be a Horse Mart. When I was in Bombay, five years ago, the following were the principal Horse-dealers :

Soorabjee Daddy Santook... Sonapoore.
Bazinjee Fakeerah ..... do.
Mahomed Bawker Khan ... Baindie Bazaar.
Nowrojee Nusseerwanjee ... Bycullah.

They are all very respectable men, understand thoroughly how to describe and show horses, and are, indeed, as accomplished hands at their profession as you will meet in any part of the world.

They get a commission of 17 Rupees from the purchaser, and 10 Rupees from the seller of each horse, without respect to price; so that, considering that they dispose of hundreds of horses annually, they make rather a good thing of it; and ought to erect better stables than the temporary and objectionable ones now in existence.

* Nervous excitability is used in contradistinction to dull temperament, yet there is a peculiar kind of docile laziness of temper characteristic of many high caste Arabs, but which, by dint of good training, put forth their energy, and shine on the turf as well as others of more choleric blood :—

J. Stewart, p. 55 : "It is to be remembered, that good conformation merely gives the power to perform extraordinary exertions; the will to exert that power depends on something else, which appears as necessary for great feats as the conformation. This energy, or disposition to work, may be too great for many purposes. The racer, indeed, can never have more than enough : it is a combination of energy with good conformation that commands eminence on the turf."

Soorabjee Daddy Santook's is, I believe, the oldest Firm. Daddy is a capital judge of a horse. Kind, obliging, and thoroughly honest and upright in his dealings, he appears well known to Gentlemen in all three Presidencies, who have such great confidence in him, that during each season (say from November to January) they make remittances in advance, and leave the selection of horses entirely to him; and I must say, from what I saw, that he studied their interests most faithfully, by procuring for them their money's worth.—Ed.]

The Build, commencing with the Quarters. Tail, Thighs, Stifle, etc.

Four of the most desirable points for the racer are, powerful well-shaped quarters, thighs, and stifles; and large, lean, bony hocks. The straighter and longer the spine is from the croup—the end of the loins—to the setting on of the tail, so much better is the quarter; for the most scientific writers tell us, that the straightness of the spine here, causing a slanting direction in the haunch and thigh-bones, enables the muscles to act to the greatest advantage, and that it is in the advantageous direction quite as much as the bulk of the muscle that the propelling power lies. Let that, therefore, be a *sine qua non*, that from the end of the loins to the setting on of the tail the spine must be straight, or nearly so, and it cannot be too long. A broad quarter may carry weight, but it must be long also to gallop, and the tail should be carried out straight, and carried well (it does not require always to be carried high to be carried well,

but it requires to be well set on, that is, set on high up). The tail of a blood Arab is likewise generally thin, of small circumference, tapering to a very small point, and the hair on it not two bushy. "Full mane, if you like, but thin tail." Carrying it on one side can be no sign of bad caste, although it certainly looks ugly.

[I do not think carrying the tail on one side indicates any particular caste, but depends upon weakness about the loins.—Ed.]

The thigh should be as broad as the haunch-bones; and as much as they swell out broader, they are better. A horse that is ragged-hipped, it is said, is not objectionable; but by this is meant, that, *notwithstanding* the great muscularity of the quarters and thighs, the protuberances of the haunch-bones still project a little wider; and not from the quarters and thighs being at all deficient in muscle, which is often also called ragged-hipped, and makes the haunch-bones look a little prominent.* Great thigh-muscle may recompense in some degree for a slight droop in the croup; but great thigh-muscle, added to a long and straight croup, is the make for the turf. A

* The side bones of the loins—the transverse processes of the lumber vertebræ—being short, are a cause of a horse being ragged-hipped; and these side bones being thus short, are also the cause of a horse being wall-sided; therefore it is that a wall-sided horse often looks ragged-hipped; but a deficiency of quarter and thigh-muscle, which causes the haunch-bones to stick out a little, is much oftener called ragged-hipped.

large hinder channel, from the anus downwards, caused by the muscles projecting, and being well asunder, is another good point, and much admired: it is a sign of great endurance.

The stifle and all that below it, must be broad and strikingly muscular: he must be what is termed well let down in the thighs, plenty of muscle inside the thigh as well, as outside, never hare-hammed; and the muscle just above the outside of the hock, in particular, should be large.* Some horses are accounted too long in the haunches, forcing them off at full speed, but this is not against their running well. These three points, the quarter, thigh, and stifle, show the blood of a horse quite as much as the head, eye, and jaw. A muscular and elegantly dropped hind leg, having a fine elastic spring, is far more difficult to find than a casty head.

The Hocks and the Hind parts proof of half the breeding.

The hock of a fresh unworked Arab is generally beautifully formed, and clean, and particularly excelling in the hind projecting bone at the point. You must never expect a runner in one who is glaringly deficient in this point. The hock should be broad also from side to side; the bone at the *upper inner* part prominent. As to whether the leg for a running horse

* The part a little below the stifle being narrow and wanting muscle. See Frontispiece.

should descend straight from the hock, or incline under the body, neither should be in extreme : there is a medium. The stride is said to be lengthened by its being well bent under, yet, that there are objections, arising from the greater wear of the hock, that greatly counterbalance this. Either for a racer, charger, or hunter, a rather straight drop for an Arab will be best and handsomest. In the high caste Arab, it invariably does descend rather straight, but whether the leg from the hock is bent under or straight, the hock itself must neither turn in nor out ; if the former, he will be cat-hammed, making his legs like an ill-shaped cow's, or, as the ladies say, donkey-legged ; if the latter, which is not so often met with, the toe will turn in, which is worse. For a running horse, it is an advantage if the toe is behind the stifle-joint.

A BONE SPAVIN in the clean hocks of an Arab, is generally visible enough if situated at the top of the splent-bone, that is, *on the inner side of the lowermost part of the hock.* If he should be spavined higher up, or the ligaments have been strained, it is not so easily perceived as the former, but in either case both the hocks will not look exactly alike ; and this will be quite sufficient to reject him. A small bone spavin is often a most difficult disease to detect ; and this is not extraordinary, considering that professional men


have in some instances been deceived, from not examining attentively. A spavin on the head of the splent-bone, on the inner side of the lowermost part of the hock, is often productive of no lameness if situated well forward, not even though it be as large as a marble ; therefore this, although the most frequent seat of spavin, is not so bad as that which lies a little higher up, for here is the principal motion of the hock, and the smallest ossification always causes more or less lameness. You must discover if there is the slightest difference either in the feel or appearance of the hocks in these two places (for a horse is seldom spavined in both hocks in exactly the same manner), and if there is, pause before you conclude the bargain. Should there be any difference on the outside of the hock, blows, kicks, or hurts have most probably produced it, not strains or over work, although spavin is occasionally found in this part likewise.

[Bone spavins may exist, and yet not be visible to the eye : they are generally formed by the union of the three little bones (cuneiform) which are situated on the front, and inner part of the hock, just above the shank bone. Between the cuneiform bones, are placed bursæ, containing an oily substance called "Synovia." From severe exercise they become ruptured, the bones are then brought into contact with each other, and acute inflammation is the result. During this state of the disease the horse is suffering great pain, and is very lame. Nature however soon comes to his relief by throwing out osseous matter, which in time unites the bones. After the union is complete, inflammation ceases, and consequently pain also. The animal however

has a stiff hock, but is fit for work, and may continue to be useful for years.

Spavin also arises from badly formed hocks, and it is not unusual to see Colts that were never crossed, spavined. Have nothing to do with such animals, for as the disease is only in its early stage, the chances are, that when they are put to work they will go lame, and remain so for months. In old horses the mischief has been completed by the union of the bones; and beyond a stiff hock, you have nothing to fear.

Spavins are also produced from external injuries, as is seen in the following sketch, from a specimen I have had by me since February 1859.



a. Where blow was inflicted on cannon bone.
b. Bony enlargement; but not spavin.
c. The three cuneiform bones united to cannon bones, causing spavin.


Spavin caused from external injury.

The Horse was a hack in Taylor and Co.'s, and received a blow from a stick just below the hock; after the inflammation and swelling had subsided, he returned to his work, when there was an enlargement observable just below the heads and between the metatarsal bones; (cannon bones large and small) this however did not interfere with the action of the hock joint, but within a couple of months' time he was again sent to my Infirmary for


treatment, and was so bad that he could not bend the hock, osseous matter having extended upwards till it united the cuneiform[c] with the metatarsal bones. This is a very unique specimen, showing how a bony tumour ("exostosis") may terminate in ("anchylosis") the union of several bones, and also, how an enlargement may exist, and be taken for a spavin, when such is not the case.

It must be apparent from this brief description of spavins, that they are incurable, and that firing and blistering the hocks are resorted to merely to increase the inflammation existing within, and thereby bring on a speedier union of the bones than would otherwise take place.

When osseous matter is produced to a greater extent than is required to unite the bones, all that is superfluous protrudes between them, and hence the enlargements called "Spavins" in every part of the world but the Cape, where those versed in horse flesh call them "Jacks." When there is no superfluity of osseous deposit, the bones are merely united without any prominence.



Sound Hock.
Showing the *three cuneiform bones distinct and unconnected with each other by osseous deposit.



Spavined Hock.
Showing the union of the † three cuneiform bones by osseous deposit.

Non-professional men who pride themselve in a thorough knowledge of bone spavin, in reality, know nothing whatever about it, at least, I never met one who did, although I have heard a great many talk very largely, and point out very knowingly particular parts of the hock where a spavin would be of serious consequence, and other places where it would not matter at all; little knowing that, in its mildest form, it must affect the action of the hock. I have shown some of them the bones forming the hock, and the difference between a sound and a spavined one, when they have admitted that they never knew that spavins were of such a complicated nature, and that they were always under the impression that a spavin arose from an *enlargement* of some bone, and *not from a union of several.* They pin their faith entirely on any prominence that may appear on the inside of the hock, be it normal, or abnormal, and deny the existence of spavin unless it is so.

The professional eye can however detect the existence of spavin from the gait of a horse, particularly whilst trotting. Entire flexion of the hock being impossible (for reasons already given,) the foot is imperfectly raised, and becomes worn away at the toe from being dragged on the ground; besides, when the animal is stopped short, or turned suddenly at the trot, the leg is jerked in such a stiff and peculiar manner as to leave no doubt on the subject.—Ed.]

If the least swelling or enlargement exists in any part of the back of the hock, below the point, which may be detected by carefully observing if they are both alike, you must reject him also; for this will denote a strain, perhaps a curb.

Capped-hocks are a great objection; they proceed from blows, hurts, or contusions from kicking, rarely, it is said, from a strain; but are apt to enlarge on

work. When the hocks are but very slightly capped (for if one is capped the other is generally found so too), they may give the appearance of the bones of the points being long. Don't be deceived in this way, but feel if there is anything like a tumour there.

The bog-spavin, a puffy swelling in front of the bend, and a little towards the inside of the joint, can never be mistaken if the finger is pressed on the large vein that runs over it, which will now sink in half an inch, and when the finger is taken away, bulge out again. If found in one hock, the other will generally have it also; and therefore, as far as this disease is concerned, though they may appear alike, they yet look full and fat, not lean and dry, as they should do. Bog-spavin generally looks larger in the middle of the day, the cold in the morning before going to exercise acting as a sedative; or the excited action of the absorbents during and immediately after exercise operating in keeping it down a little.

The thorough-pin, another small puffy swelling on the outside of the hock, nearly on a line with the point, and running through to the inside, may not prove any hindrance even in hunting; but for a racer there should be no doubt about the hinder extremities in any way, either as regards build, injury, or disease; they are the grand agents in progression, and a race-horse, to shine, must be nearly perfect in every thing

behind. You are not to infer from this, that a horse can run because he has no defect in his hind parts, only, that he will rarely run well if he has. Formed as described under these four heads of Quarters, Thighs, Stifles, and Hocks, they will look handsome both to the eye of the novice as well as the judge; and if, added to this, he has a decided casty head, you may take it for granted he is well bred; but the handsomest and most casty head that was ever on an Arab is only indicative of half the breeding, the hind parts must be formed as before laid down, to prove the other half.

All the good breeding in half or three-part bred horses will sometimes show itself in front, sometimes behind: for instance, put a thorough-bred horse perfect in make to a half or three-parts bred mare: one year a foal will probably be thrown badly formed behind, with a casty head and good forehand; another year one perfectly formed behind, with an ugly head and bad forehand. The latter of these, with the good hind parts, both being of equal good blood, would be the most promising for running. But again, a half or three-parts bred horse, even possessing all good running points, will, nineteen times out of twenty, be beaten in a long race by a thorough-bred horse not so well built as himself. This is well known in England, and the same law, of course, holds good with an Arab.

The Head, Nostrils, Mouth, Lips, Eye, Ears, etc.

The head ought to be lean and bony; the jaws wide, and not fleshy; the channel clean, and not filled up.

The nostrils must be open and thin, or he is useless for the turf; the best training in the world will never compensate for a thick shut nostril.

The mouth should be deep—a laughing mouth—which will allow more room for the opening of the nostril, and also give him a better mouth.

The lips should be thin and evenly closed, for the under lip hanging loose is a very unfavorable omen of his running.

The eye is never too large, and scarcely ever too prominent. A small eye should be objected to, as being more liable to disease, and symptomatic of cross blood. I warn you to look well into the eyes, so very many bear the milky marks, the result of inflammation. Whenever you find one with a cloudy appearance round the hedge and close to the white, or one with white marks streaked across the centre, or one which on a close examination appears a little smaller than the other, or puckered in the lids, depend upon it there has been disease, which is likely to return. Recollect a horse's eyes are not placed like a man's: if he is blind with one eye, he is unable, without turning his head round, to see any thing on

that side with the other. The slightest derangement in the eye may at times prevent him using his speed to the utmost. Should any speck of white be observable near the centre of the eye, it has been caused by some blow or hurt, and may interfere with the passage of the light through the pupil : you should therefore distrust it, for it is a very ugly blemish at all events, and reduces the value considerably. Some few others bear a resemblance to the green glass eye : there is as much difference between the clear, transparent, bright, healthy eye, and this glass eye (which is written as arising from the loss of power in the optic nerve, and generally incurable,) as there is between a polished diamond and a piece of window-glass. Shying is occasionally connected with some little alteration in the organs of vision, imperceptible to us who are not oculists.*

The ears of a genuine Arab are small, and the hair inside is silky : they never have that tremendous curling in of the points so proverbial among bad Persian and country-bred horses. Lop ears are quite as ugly as those that curl in, but a large lop ear, if only well shaped, rather peaked, and not large round at

* If you blow a soap-sub bladder between your thumb and fore-finger, and hold it up to the light, you will discover a kind of purple variegated streaks on the surface : these colours you will often see over the horse's eyes by narrowly inspecting them, keeping the head in a greater light than the body, as just within a stable door. I do not know their cause, but imagine they have no business there.

the points, with a thin skin, is no such sign of bad caste; it appears to belong to a peculiar breed.

The glands below the ears are sometimes found swollen; and if much so may be a serious detriment.

A small head on a large horse, or large head on a small horse, are of course equally out of proportion; a trifle either way is perhaps immaterial, if only lean, with the large eye, broad forehead, and hollow jowl; yet, a head possessing these and all the foregoing good points, may still, from some peculiarity of conformation, not altogether please the eye, or be a very handsome head; but one void of these requisites can never be termed a blood head. A "cut short" head, with a slight indentation about half foot above the nose, is a great favourite with some judges. I like it too.

The Neck.

The neck of an Arab I never knew too high or too long, though it is often quite long enough. Take care especially that it is light, and that the upper surface is not too thick: it should be very muscular towards the bottom, but enter the chest above the points of the shoulders; if it does not, you will have neither speed nor lightness of action: it must also curve a little where the head is set on, or he will pull against your arm, and feel heavy in hand, consequently this will never answer for a charger or hunter, and

it is nearly as bad for a racer. There must be no superfluous thickness on the upper part of the shoulder-blade : the division of the neck and shoulder must be distinctly marked; any extra bone at the upper part of the shoulder-blade, filling up the line of demarcation between it and the neck, always makes a horse go more or less stiff, and this is particularly observable in thick necks; yet, there must be sufficient muscle below to prevent the neck being a "loose neck." If he has a neck like a deer it is ugly, and does not afford so free a passage to his breath : the deer necks, however, with all the flesh set below, are not so bad as those where the upper surface is thick, and it is all set on above. Ewe necks, if only slightly ewed, and light, and not entering the chest low down, are rather favourable than objectionable for a racer.* The rounder a neck is the better; and if only shallow from mane to under surface, there is no fear then of its being too muscular from side to side. No thick neck is fit for the saddle, and a bull-neck is fit for nothing but a bullock-garree.

The Withers and Back.

High withers you will not always find, but they should undoubtedly rise a full inch higher than the

* Darvill, vol. ii, p. 10. I have a great aversion to a high-crested race-horse.

top of the shoulder-blade, or if you like the expression better, as high withers are not at all necessary for the turf, the top of the shoulder-blade must be a full inch below the top of the withers;* and if the withers are wide at top, he will generally carry weight better than if rather fine. After the fall of the withers the spine must run straight, without any ups or downs, or arches, to the croup at the end of the loins, that is, to the centre between the haunch bones; and from there, as mentioned before, it must be carried out straight to the tail. If the spine has any of these arches the back will be galled by the saddle, and if there is a depression at the back of the saddle, where the back and loins join, it shows weakness. A depression also where the loins and croup join, "the hind quarters looking separated from the back," is twenty times worse. After the fall of the withers, the spine is often straight, yet gradually ascending; this is an advantage, provided the ascent is continued to the end of the loins; and if from thence it is gradually carried on to the tail, he will, notwithstanding a good height of withers, look higher behind than before, which is favourable.† The straightness of spine is

* I once saw an Arab—a deformity—with the top of the shoulder-blades higher than the withers, yet he was hog-hunted.

† William Osmer. Fifth edition. Treatise on the Horse, p. 222. "If the foreband be more lofty than the croup, he cannot run worth a curse." How ridiculous it is to see a London horse-dealer placing the horse, when

essential: a horse at full speed, with his head and neck thrust out, gallops as horizontally as possible: a hollow backed horse scarcely ever runs well. Purchase none for the turf, therefore, but what may not be inaptly termed a horizontal horse.

It is actually requisite to have some length of back for the turf, and *provided* the quarter is long, the shoulder oblique, and the spine straight, there is not much fear of the back being too long. A long quarter, a moderately long back, and a rather long neck, must necessarily make a tolerably long horse, which is the form desired for a racer. A tall, short horse, is not found to keep the pace up so well as one that is longer. The length should therefore be great from the point of the shoulder to the hindermost point of the quarter; and then, if with this long quarter added to a good oblique slanting shoulder, the fore-legs are planted well forward, and the hind-legs properly dropped, there will be observable that "shortness above and length below" so much sought after; in other words, there will be a long horse with a short back. The reverse of this, a short horse with a long back, is shown in the frontispiece. So, the true

brought out for inspection, with the fore-legs on the highest ground, making him appear of that build as if he would slip his girth. A well-built horse should merely have a nice rise in the withers, but the above abominable position ruins the appearance, making the whole backbone from the withers to the tail gradually declining backward like the formation of a cameleopard.

meaning of a long horse being length from the point of the shoulder to the hindermost point of the quarter, it is very possible to have a long horse with a short neck also, which, if light, and with a curve where the head is set on, will do admirably for a racer: or to have a short horse with a long neck, which is more adapted for a charger or hunter.

The shoulder must run back with a good slant; and then, if the withers rise a full inch above the top of the shoulder-blade, it will not be found too heavily laden. It has been asserted, by both Europeans and Natives who have been best conversant with the Arab, that nothing denotes the superior cast more than that extreme obliquity of shoulder, and certainly nothing denotes the make more. When the shoulder is very oblique, how short the neck appears below from the back part of the channel to the chest; and how long above from between the ears to the withers! What superb conformation! An unexpected trip easily throws a horse with a straight shoulder down, and he generally hangs very heavy in hand as he tires. The straight-shouldered horse, if shining in other points, may pass handsomely for draught in a carriage or buggy, but will never answer well for the saddle. From the top of the blade to the point it should be long, very long, for that will compensate for a little deficiency in the obliqueness; and the space from the point to the fore-arm,

or leg, particularly short, in order that the legs may stand well forward; and the chest should be moderately broad, in order that these forward legs may not be too close together, but while it is broad it must also be flat; not concave and hollow, but flat, and not overloaded in front. A round full chested horse will do very well as a carriage-wheeler, but not for the turf. The great, broad, rounded and projecting hang-over chests, with the fore-legs often inclining under the belly, are continually called fine chests, but they are as bad as bull-necks.

The depth of Girth, and Carcase.

The depth of girth cannot be too great, and the carcase, which should resemble a barrel, and not be flat-sided, should swell out well under the elbows; then, with the moderately broad chest, you will be sure to have what are indispensable, good large lungs. A carcase that is flat-sided is equally bad for running as for appearance. The depth of girth is a point that admits of measurement, but then the chest must sink deep between the fore-legs, or else very high withers alone may contribute more than their proper share to make a large girth: sixty-four to sixty-five inches will be very good for an Arab of fourteen hands to fourteen hands one inch; but if you have an eye for a horse, you will tell at once if it is good, or not. There should also be some distance between the last

false rib and the haunch-bone, that the hind legs in the gallop may be thrown well under: this is allowed to be an excellency in the racer, though it detracts a little from the strength.

Horses, whose carcases are light, and legs rather long, "showing too much daylight," have been run down for more than is necessary: they are only objectionable for heavy people. When there is good build otherwise, a small round carcase, if it only swells on behind the elbows, need never be objected to by a light weight, for it rather adds than detracts from the beauty, and they are often splendid runners. A large carcase; a large circular barrel, with deep ribs, deep in the fellers, must have large broad flat legs to carry it, or they will quickly fail.

The Elbow and Forearm, etc.

The elbow, the bone at the top of the forearm, must be large and not turn in: the top of the forearm must be decidedly swelling and muscular: the length of the forearm from the elbow to the knee cannot be too great, in order that the distance between the knee and fetlock may be proportionably short: these are three indispensable points, but the two first are seldom sufficiently scrutinized. A horse with a thin forearm and corresponding lanky thigh, cannot possibly run, nor stand work either.

The Knee, and Back Sinews, and Suspensory Ligament. The Shank-bone, Fetlocks, &c.

The knee must be broad and flat: *the upper inner part* should present a striking width, or it will look round, which is ugly, and does not betoken strength. If there is the slightest bony excrescence in front, it may interfere materially with his running, and which firing and blistering will often fail to remove; and if there is any puffy swelling, it is worse than the bog-spavin. The hinder bone of the knee cannot be too large, so that the leg may not be tied in, that is that the back sinews may be wide away from the shank-bone; and these back sinews should feel smooth, strong, and well braced, like a piece of catgut tightly covered with fawnskin. The back sinews cannot always be told by the feel; they must be carefully inspected at each side, for sometimes, while they feel smooth, strong, and wiry, there will be an evident difference in the size about the centre; a very slight bow—a sure sign of having been injured. The suspensory ligament, the centre rope, must also feel and look, like the back sinews, fine and wiry. There is however, a distinction between smallness immediately below the knee, and that lying in which deforms the legs, and it consists in this. Some of the best Arabs, even while the hinder bone is of good size, are yet small below the knee, but equally small at the fetlock; this is no imperfection, but merely renders such

horses a little slight, and in this sense is not very good for the turf, and a great objection to weight certainly; but one that is small below the knee, or if not very small, yet gradually inclining broader towards the fetlock, is not only most unsightly tied in, but is one of the worst faults a racer, or any horse can have; for when the back sinews are thus tied in, the legs are very liable to become crooked, and fail in hard work; besides which, this last description of tying in is, like the small eye, a strong sign of mongrel breeding. A horse that is slight, slight as before described, all the way down, and short between the knee and fetlock, will be quite as strong as one that is broader below the knee, yet longer between the knee and fetlock; but great breadth immediately below the knee, in addition to shortness between the knee and fetlock, is needed to make this part perfect for a racer. An English groom of some experience that I knew, used to say, "Always be mindful of the scientific maxim: when purchasing or betting, if you feel a doubt, and have an option between two horses, always decide in favour of that one with the largest three essential bones; the hinder bone of the knee, the elbow bone, and the bone at the point of the hock."

The shank, when viewed in front, must look fine, and of small circumference. There is no surer sign of hard work than the shank, when viewed in front,

looking round and large. A curb or spavin may come by a strain, but the roundness of shank I here allude to—a thickening of the skin and cellular stuff beneath, quite independent of the bone—is produced by nothing but sheer hard work. Some few blood horses, exceptions, are born rather rounder in the shank-bone itself than others, yet it is bad ; and even this roundness in the bone itself is very ugly, and looks like work ; but even one whose shank-bones are naturally rather round, will become rounder still much sooner than one whose shank-bones are naturally flatter. Plenty of muscle above, large back sinews below, but small flat shank-bones, solid like ivory, is the blood horse, and horse for endurance or speed.

A splint, if situated about the middle of the shank-bone, is of no great harm, save that the price of a horse, if bought for appearance, is much diminished by the unsightliness of a piece of bone on the side of the leg, a lump there, from its prominent situation, offending the eye more than a large spavin. If found just below the knee, it may interfere with the motion of the joint, and if on the inside, it may be the cause of a speedy cut. If near the suspensory ligament, it also sometimes affects the free action of the leg ; and if extending on to the back sinews, it too often causes lameness. In all clean fresh blood legs, there are three different parts distinctly visible : the small, round, flat-looking shank-bone, the suspensory liga-

ment, and the back sinews; and these are clearly free from all lumps, and bumps, and bony excrescences whatever; the only natural prominence being the end of the splint bone, about the size of a pea, two inches above the fetlock.

[Splint—is a bony deposit in the inside of the forelegs, as described in the annexed diagram.* The splint bones are united to the cannon bones by a Cartilaginous substance which becomes osseous after the horse is put to work. When the deposit is greater in quantity than is required for the union of the bones, the superfluous portion is what forms a splint. The inflammation and pain are very considerable during this change of structure, and the Horse is very lame.



1. A splint, involving the bones of the knee joint.
2. A splint, interfering with the action of the back sinews.
3. A small splint situated under the tendon of an extensor muscle.

It often occurs that when young horses are first put to work, they are constantly blundering from splints just forming. I have known very valuable horses sold for a trifle because their owners looked upon them as "Chest foundered," and consequently confirmed stumblers. Whenever you find a horse trip more than usual, particularly a young one, examine his legs carefully, and press upon the part where the inner splint bone is united to the cannon; and you will very likely discover a splint so small, as not to be noticed by the eye, and yet so painful, as to cause a horse to drop under his rider as if he were shot.

* Taken from Mayhew's Horse Doctor.

Splints are found in the outside of the forelegs, but this is of rare occurrence.—Ed.]

The hair about the sides of the fetlock is sometimes a little ruffled: if you feel carefully, you will probably find he has been slightly fired. At other times the hair will look of a lighter colour; when, if you get him in a proper light, you will probably find he has been blistered. Ask the reason, and you will be told, he was fired for nothing, just as a preventive; or if blistered, merely for fun. Look on these as screws, and deduct at least seventy per cent. in consequence; for a man who is cruel enough to fire for nothing, or to blister for fun, will not hesitate to palm off on you a bad and patched-up leg.*

The small scars, or shortened hair from cutting, either behind or at the sides of the fetlocks, are easily seen. If cutting arises from the toes turning too much out, it is, of course, incurable, for "a goose will always go like a goose," and it renders a horse much

* Some sporting novices say, a fired leg is worth two others; it never goes. A fired leg, it is true, will often stand training when its fellow fails; the reason is plain: the horse favours the fired leg, and the other one has to stand the extra work; consequently, the good leg goes first, but the bad one has caused it; besides, there is good authority on record, that a leg never moves so freely after having been fired. It is astonishing how much a horse can favour a leg, without its being noticed by ordinary observers; and it has been remarked that ladies' horses generally fail in the near fore-leg first: these dear creatures always must have the horse lead with the off-leg, therefore it does not get its fair share of the stress, though the foot may get more battering.

more unsaleable than a loss of hair from girthgall or sore back, both of which are looked on as most serious objections, as, the hair once off, the spot is so easily galled again, that the horse has frequently to be laid up every other month.

At the inner side of the fetlock, as often as the outside, there is sometimes, in the otherwise-cleanest legs, an ossification. I have seen this twice passed over by a judge, though it was as large as an acorn: the feel and the difference of appearance of the two fetlocks, on nearly a front view, is sure to distinguish it. Ossifications at the sides of the fetlock, like spavins, are sometimes large, without causing lameness; but they are invariably productive of some slight stiffness, though not discernible to every eye. If the enlargement is just above the inside of the fetlock, and on the suspensory ligament, lameness will generally quickly ensue on work.

[Suspensory ligament, so called from the two little bones (sesamoid) forming the back part of the fetlock joint being suspended from it. Like the tendons (back sinews) it is often sprained from hard work and causes great lameness. Few men who profess to know the treatment of horses, look here for the seat of lameness, and finding the back sinews uninjured, the shoulder is safe to be considered at fault, and blister after blister is applied to it without producing the anticipated result.

If you are examining a lame horse, and are in doubt as to the part affected, take up his leg and press the ligament with your fore-finger and thumb above the fetlock. If it be injured, he

will wince under the pain you have inflicted and struggle to get away from you.—ED.]

Immediately above, and in front of the fetlocks, there is also frequently seen a kind of dent, the fetlock looking as if there had been too much stress on it; and so there has been. This may be produced in a slight horse by a single month's riding of too heavy a man, though it will take many months, with rest and bandages, before it resumes its primitive cleanness. When it accompanies the large rounded shank-bone, the chances are, he has been long shamefully overweighted, as well as overworked; and horses in this way, though neither standing with their fetlocks knuckling over, ready to "bite the dust," nor having windgalls, are often more or less groggy, and you run the risk of some latent internal disorganization about the joints of the foot, which sooner or later will produce one of those lamenesses that "come of themselves." You cannot reasonably expect to dispose of a horse again with these large rounded shank-bones and dents, except as a baggage-horse.

Windgalls are a most annoying eye-sore. As unfair or overwork produced them; so they will increase on work. If they are not large, and no roundness of the shank-bone accompanies them, and you are not particular as to appearance, he may be well worth half price: never purchase, however, a horse for the

turf, that shows by such evident signs as all these, the combined effects of bad legs, mismanagement, and overwork. To the former sagacious maxim of the three essential bones, my stable acquaintance impressed on me another injunction, which I hand to you, with advice to keep it uppermost in your memory:—"Never fall in love with thick ankles." A fresh horse, he said, like a French danseuse, is always delicately clean at this much-admired part.

From the fetlock to the hoof should be rather long, but not over-slanting: the pasterns of many Arabs, when long, are sometimes too sloping, the fetlock nearly touching the ground. The short pasterns, even, I have seen too oblique; and then, though the limb should be otherwise powerful, the beautiful springy action will be lost in a month by the riding of the least heavy weight; yet, on the other hand, if too upright, the fault is far worse, for the knuckling over will then assuredly soon commence on work, and when it does, he will often evince an evident anxiety to make his nose a substitute for a fifth leg. The pasterns of a horse, for a heavy weight, should be rather short; but for a light rider they can hardly be too long, *provided* they are properly placed. They should look small, round, and smooth; if there is the slightest enlargement, you may suspect a ring-bone.


The easiest and surest way of detecting a ring-


bone, is to place a couple of fresh young colts (and they are quickly got at the stables) besides the one you are buying; then feel each all up the centre of the pastern; and also at the side close to the hoof, about two inches from the heel. If ringbone is formed in either of these places, there will be an evident difference in the feel, which should make you sceptical as to the soundness, for ringbone is not uncommon at the stables among the newly arrived horses, both old and young. There are two ligaments that run down on either side the pastern, and it is the centre of the pasterns between these ligaments* that will show a fulness and hardness; or, if the ringbone should be at the side of the coronet, about two inches from the heel, you must discover it by the difference, which you will if you search narrowly, first feeling one horse, then another. The centre of the pastern between these ligaments about two inches above the coronet, quite on the seat of ringbone, is sometimes a little swollen, the consequence of having been fastened with a rope there; but this, though common enough among country horses, seldom arises from this cause with Arabs. I knew a cunning fellow, who purchased a new arrival from the boats with one of these swellings on the off fore pastern, which unluckily turning out a ringbone; he cleverly tied a rope

* These ligaments, in some flat-shanked, wiry-limbed, clean-pasterned horses, occasionally stand out very prominent, like a piece of thick cord.

round it, so as to take the hair off a little, and searching out his cousin, a great connoisseur of country-breds, sold it to him for nearly double what he had paid. Be cautious then of your friends, for ringbone often produces no lameness till it spreads on the joint.

Healthy Bones.





Diseased Bones.

b. Large pastern.
c. Small pastern.
d. Coffin bone.

1. Union of the small pastern with Coffin bone.
2. Union of large with small pastern bone.

[Ringbone is of the same nature as bone spavin. Osseous matter is thrown out, and unites the lower portion of the large with the upper portion of the small pastern bone, producing a fulness in the middle of the pastern. This is most frequently the part affected, but sometimes you will observe a swelling just above the coronet, when the lower end of the small pastern bone becomes united with the coffin bone as shown in the cut. Ringbones are more frequent amongst draught horses, that have to drag heavy loads, and are seen oftener in the hind, than in

the fore pasterns. They may involve bones partially or encircle them completely, and are as serious in their consequences as bone spavins.—ED.]

Position of the Legs.


The forelegs must stand straight, as you may naturally suppose : you will not be griffin enough to purchase crooked legs. They should look moderately wide at the chest, gradually approaching each other at the fetlock. If they stand fixed wide apart, "pinned," like a horse laboring under inflamed lungs, it is almost as faulty as if they were crooked from the knee downwards; but in the latter case, the knees look a trifle too forward, in consequence of the legs being a little bent backward ; he does not in fact stand straight on his forelegs, and therefore you should take the hint here, as well as in the capital crime, the knuckling over of the fetlock joint; for they always become more crooked on work. These unlicensed pins are sometimes defended on the plea of the horse having been born so, but one that was born crooked is surely quite as bad as one that has been worked so. The leg at other times, from the finish of the knee, has a slant forward instead of backward, the knee appearing too much straightened, bent a little back, calf-kneed. This is a malformation, but never proceeds from work, like the former, neither is it of a hundredth part the consequence ; indeed, some persons prefer a slight slant forwards, so as to be sure

there is none backwards. If your own eyes are straight, and not askew, you ought, when standing about a yard distant at the side, to discover either of these defects at a glance, however slight they may be.

The legs are sometimes bowed, and a bow-legged horse generally dishes, and one that dishes will never run. Dishing action should not be called bad action, but rather floundering, or crooked action : bad action is when the legs are not lifted sufficiently high, nor propped sufficiently forward.

The feet are not to turn out, nor in : if the former, he will be liable to cut ; if the latter, it is equally bad, for the weight either way is most unevenly distributed. The point of the toe should be found exactly under the point of the shoulder : dropping your stick perpendicularly from the point of the shoulder will tell you this in a second, if your eye does not. As much as it is behind that, so much is the weight thrown too much forward ; and as in this case the chest will look either like the chest of a dray-horse, called, a remarkably fine chest ; or the legs will slant backward under his belly, which is as bad ; so will he be more likely, especially down hill, to fall. This you may likewise naturally suppose ; for when standing upright on your own legs, and inclining your head and body forward, a push behind will easily upset you.

The hoofs of a genuine Arab are rather small. You will seldom find them too small, *provided* the hinder part of the quarters are the broadest part.



A A, the hinder part of the quarters; B B, the bars; C C, the frog; D, the cleft; E, the toe; four F's, the sole; six G's, all round the crust; H, a line drawn across the centre, showing the outside bottom of the hoof, is more circular than the inside. If the hinder part of the quarters is not the broadest part, then the foot was either never perfectly formed, or has become contracted from want of paring, bad shoeing, or disease. If the hinder part of the quarters should be as broad as the forepart of the foot; the frog a good width; the bottom of the foot properly concave; and the hoof in front, from the coronet down to the toe, rather upright, this will be the next most desir-

a long and horizontal spine is very fine; but have nothing to do with a long and horizontal foot. The sole, however, is often flat without any lengthening out of the foot, and so thin that it bends like a piece of whalebone. One or two months' gentle riding, is generally all such a foot will stand before it descends quite on a level with the crust. When it sinks lower still, down to the convex pumiced foot, it is, of course, worse; but this is merely like a difference of choice between a wooden leg or crutches.

Do not neglect to see that both feet are exactly the same size, for this is as often heedlessly passed over as the difference in the size of the eyes. The two hind feet must correspond likewise; but not being so liable to contraction, they seldom get out of order. If after all this you are still at a loss to comprehend what constitutes a good foot, keep the heels as open and the foot as hollow as the hind ones; then you will not be far off the proper form.

White feet are considered objectionable; but recollect, a white foot, properly formed and shaped, is far superior to a black one badly formed, and that white hind feet are not of much consequence.*

Examine attentively the warmth of the feet; but

* Horses with white feet are more subject to cracked heels in wet weather and if not properly dried after washing. The crust also is generally more brittle.—Ed.

this must be done early in the morning, by eight o'clock, before the horse has been exercised, to be of any avail. If one is warm whilst the others are cool, there has been a strain of the coffin joint, or there is something else wrong internally; but the lameness, in some of these cases, is often so very slight, that you will not discover it till after one or two ridings.

You should observe also, whilst eating grass in his loose stall, if he places one foot stretched out before the other: if he stands decidedly in this position, one foot too much under the body, and the other pointed out nearly a yard in front, you may suspect there is something wrong, either a recent hurt, or, more probably, some old-standing tenderness. A dealer in England will prevent your seeing this, by not placing such a horse in a loose stall; but in India it is generally boldly exhibited; and I once heard the cause equally as boldly accounted for. The buyer asked, "what makes him stand in that awkward position, with one leg pointed forward?"—"Oh," replied the seller, "that is merely in consequence of the shortness of his neck; he could not reach his grass easily otherwise, many of the Arabs have short necks." The innocent victim, who was a native of the Scilly Isles, appeared quite satisfied with this answer, and giving another look at the neck, said, it appeared to his eye of very good proportion, and concluded the

bargain. This was one of the worst kind of the long, horizontal, flat-soled feet.*

[When a horse is eating grass, or is standing still in his stall, with one foot placed much in advance of the other, you must look upon him with suspicion, as this position generally indicates navicular lameness. This is a dreadful disease, and inflicts constant and intense pain. The advanced foot is the one affected, and being thrown forward is relieved from all weight, and consequently diminishes in size, the sound one has now to bear the whole weight of the four part of the body, and in course of time becomes diseased also.—ED.]

Look up at the testicles : they should be small and closely hung up. Swollen testicles are very common and may prove a considerable drawback to his speed. The testicles, even when not enlarged, are sometimes diseased. I don't know if this can be detected by the feel, or appearance of them, but a horse's value in India is diminished fully one-third by being gelt.

[When the Griffin's Aid-de-Camp was first published, horses were never gelt unless they were vicious, and hence the valu-

* I have myself had two horses with perfect feet; one had been shod for two or three years; the other was a colt, and never shod. The feet of both these were small, but very wide at the heels; the horn was of the proper depth, neither too high nor too low at this part, and of that fine greyish black, Oxford-mixture color. Neither of them without shoes, over hard ground, moderately ridden, ever went tender or made a false step.

Some persons, who have paid dearly for purchasing a horse with contracted heels, instead of running into the proper opposite, one with open heels, search for nothing but a large spreading foot; hence their second bargain is nearly as bad as their first; for a large spreading foot is almost always a weak and a flat foot: moreover, a large spreading foot is most unfavourable for speed.

ation put upon Geldings by "Blunt Spurs." For the turf, entire horses are to be preferred, but for the Army, and private purposes, give me Geldings. In this country, up to 15 years ago, it was considered great cruelty to geld a horse, and owing to several of them having covered, before they were purchased by Government, they were very troublesome in the ranks, and serious injuries were constantly occurring to their riders.

In the year 1847, a squadron of the horses of the 4th Madras Light Cavalry were gelt, and after recovering from the effects of the operation, they, in company with a squadron of entire horses from the 2nd Madras Light Cavalry, proceeded from Bangalore, on a trial trip, to Arcot and back again, (about 250 miles) travelling at the rate of 30 miles a day. The squadron of geldings came into cantonment with as much regularity, as they would have turned out for exercise on their own parade ground. The entires were just the reverse, and came straggling in, several hours after the squadrons had arrived.

Such a result was never anticipated, and a report was sent in by a Committee of experienced Artillery and Cavalry Officers to the effect, that geldings were to be preferred to stallions, for the following reasons:—

1st. For greater powers of endurance.
2nd. For being more docile.
3rd. For keeping their condition better.
4th. For being less subject to sore back.

And this report doomed every horse (Officers' horses excepted) in the Madras Army under 12 years old to emasculation.

It is said that a gelding never carries a good crest, unless he is operated on after he attains his full growth. I am not of this opinion. I have for many years compared horses gelt when foals, with those gelt when five or six years old, and invariably found the former better crested and bolder Animals.

Take away the testes from a foal about 6 months old, and in a

few days the incisions heal, and he grows up a well formed lively animal. But not so when the operation is performed at the age of five or six years. The horse feels that he no longer enjoys those animating and pleasing sensations his testes afforded him. He becomes timid, and so listless, that he will pass by unnoticed objects that would have caused him whilst entire to throw up his head and look about him. The muscles of the neck have not now the exercise they used to have, and hence in my opinion the deficiency of crest.—ED.]

"A small yard in a horse," says the native adage, "is an infallible characteristic of high blood; the foal he gets, will certainly be good." There is more truth in the first part of this sentence than in most of their sayings. As to the latter, it must depend, of course, on the kind of mare he is put to.

If you are purchasing in the highways and by-ways, you may inspect the vein of the neck, and also the chest, to see if he has been often bled or rowelled (lower the neck to the ground to detect the small swelling from bleeding): these are operations that are generally performed for disease, though the former too often when there is no need for it.

The Colour.

The best colour for an Arab is the grey, roan, or white,* and next to this the lighter kind of chestnut,

* A writer from Edinburgh says, "It is a remarkable fact, that an Arabian of a dark grey colour was never known in India as a winner. Bays, chestnuts, and silver greys are always to be depended on. I am, &c. &c.,

but the former are rightly preferred ; for caste, bone, shape, freedom from disease, hardiness of constitution, and good feet, are certainly, generally, more conspicuous in something of the grey colour. In the Persian less so, the grey, chestnut, and bay, having the good qualities about equally distributed between them. Some of the Arabs say, the highest caste horses are generally bay, and they ought to know best. I speak of what come to India. "A good horse cannot be of a bad colour;" but there is a great deal of fancy in colour, some *outrê* people preferring a dirty white, or dun, piebald, or tiger-marked.

Many of the horses arrive in the boats quite skeletons, and badly hide-bound. You must not be deterred from purchasing these. The head, jaws, channel, nostrils, mouth, lips, eye and ears ; the breadth of the haunches ; the straightness of the spine from the fall of the withers to the setting on of the tail ; the position of the legs ; the three essential bones ; the large square knee, wiry suspensory ligament, and clean back sinews ; none of these are effected by a horse being thin. The belly may hang like a cow's, the ribs may stick out, the neck may have lost its crest, and the quarters may have sunk down to a

Thomas Brown." This must have been my old friend, Mr. Green, whom I am about to introduce you to, at my last page ; for a letter signed Zeal replies, Mercury, Pyramus, Renegade, Emilius, Bundaola, Sackcloth, and Harmonica, were all iron-greys.

frightful hollow: the thigh, stifle, and forearm will, however, yet show a little muscle, if there ever was any; and all that frightful hollow at the quarters will fill up to its proper blood-form in a very few months, if the spine bone here is only straight, and not drooping; the neck also will regain its crest, and the belly draw up, as the horse gets into condition: but if, added to this thinness, there should be any dropsical-like swelling under the chest or belly, the breath foul, the flanks perceptibly moving with a rise and fall, and the pulse above fifty, you must consider well the risk you run: it may be only slight fever, or temporary derangement caused by the boat, bad feeding, &c.; but there may be, also, some radically bad disease lurking.

[It is quite a sight to see a Buggalow, (arab boat,) laden with Horses, arrive at Bombay. The Horses are as closely packed as they possibly can be, in two rows facing each other, throughout the whole length of the deck. There is not the least attempt made towards cleanliness; grooming in the passage being quite out of the question, and it is a wonder that more casualties do not occur amongst them. They are kept on very scanty fare, and from exposure on deck, land with catarrh and strangles, and are generally speaking very much out of condition.

The day horses are landing is one of great excitement amongst those who have racing stables, and owing to great competition, he most promising of the batch are purchased at once, at very high prices.—Ed.]

Most horses from the stables cough a little at first,

but the whole catalogue of diseases of the lungs and air-passages, excepting chronic cough, are comparatively rare in India. If the cough is very short, like that of an asthmatic person, or very loud and stertorous, either will denote some chronic derangement, and it will be on the safe side to reject him : (I have seldom passed over a valuable horse myself in consequence of this, but merely deducted something for the risk ;) and if when one nostril is closed, the breathing is not clear in the other, it will denote an ulcer in this passage, and he should be rejected also. Pinching the windwipe is the only ready mode of exciting a cough, if the walk and trot fail ; but all dealers will not stand much of this pinching, for it is very possible to bring on a cough that never existed. It must be the last thing resorted to when purchasing : you should previously make up your mind to take the horse, if satisfied with the sound of the cough.

Having finished your examination, view him in a canter. He should go wide behind* and close before, skimming the ground with his two fore feet, like a true daisy-cutter ; but whether in the gallop, trot, or walk, he must on no account step short ; the feet must be lifted with a kind of spring, and brought

* If the hocks turn out, and the toes turn in, described under hocks as most faulty make, he will of course go wide with his hocks ; that is, if he goes at all, which is doubtful : but hang such wide-going as this !

firm and flat to the ground. There never was a truer passage in a book, than that the safety of a horse depends a great deal more on the manner in which he brings his feet to the ground than on that in which he lifts them up.

You should not conclude your bargain yet, till you have mounted. The dealers, if you are a light weight, will always allow you a five minutes' walk, trot, and canter, in front of the stables, and that is as much as any man ought to ask, or get. If the fore-feet are not lifted light, quick and airy, but feel to stick or dwell long on the ground, he has been overweighted, and his action ruined, or there is something wrong in the chest, or feet.

Price.

If you are now purchasing for the turf, you will pay from eight to fifteen hundred rupees, according to age, caste, shape, &c., and your own knowledge of making a bargain. You should never exceed fifteen hundred for an untried horse, however promising he may look. Many of the best racers on this side of India, that were not particularly handsome, or if handsome, not high, have been bought by judges, and some by luck, for there is a great deal of that, for from six to twelve hundred.

If you are purchasing for the parade, and get all the points described under the following head of

Charger, with fine high action, fourteen hands two inches high, high caste, handsome, and stylish, four or five years old, and perfectly fresh and unblemished, you get him cheap at fifteen hundred. I should be happy to obtain such a one for that sum, even if he were slight.

If you are not so very particular, and can excuse an ugly head, a small eye, a filled up narrow channel, a bent fore-leg, a small hock, a drooping quarter, a straight shoulder, a bad neck, indifferent action, or any other defect, you ought to suit yourself from eight hundred to a thousand ; and if you can excuse blood, or take a Persian with one or two of these defects, you will have no difficulty in procuring it for five hundred.

If you go still further down, and take a screw of high caste appearance, with a gummy leg, a blind eye, a wapping spavin, a neatly fired ringbone, or a foundered foot, there are plenty to be had for two hundred, though they are often priced at two thousand, in the hopes of your making an offer. You will find, however, many of tolerable good caste and make, five years old, fresh and free of wind-galls, fourteen hands high, well adapted for hunting, from four to seven hundred rupees, but you cannot expect a fresh sound horse of height, caste, and appearance, for so small a sum.

The price of a high caste, well built, fresh Arab,

of five years old, is enormously increased by a little height; for every inch you will have to pay five hundred rupees. If fourteen hands one inch of this description can be bought for one thousand rupees, fourteen hands two inches, with the same qualifications *in every respect*, would be fifteen hundred rupees; fourteen hands three inches with ditto, would be two thousand rupees. And the increased height is well worth the extra money; for a large high caste, faultlessly formed, and fresh Arab of this age, is as difficult to find as a large diamond.

Two horses, very nearly resembling each other, may be very differently priced: take for instance, two of valuable caste and build, one having rather a straight shoulder, the other an oblique one; or one rather a thick neck, the other a light one; the latter, all other points equal, would for the saddle be worth fully double the former. So also a blemish, though not detracting from the actual utility, will often reduce a handsome horse from fifteen hundred rupees down to one thousand rupees; in the same manner as a dab of grease or paint will ruin the value of a dress-coat: the coat may be still as useful, but not for appearance.

Recollect, a horse's price is not according to the vulgar error, "what he'll fetch," for many a ruined horse, with a good name, often fetches more than his

original worth, when fresh; and *vice versa;* but the real intrinsic value of a horse consists in what you would have on a fair average to pay for such another, as good and unblemished in every respect, in the country, in which you may be.

Notwithstanding my endeavours to put the foregoing before you clearly and simply, I much fear, five or six days after your purchase, you will discover you have overpaid some twenty or thirty per cent. The brass, and plausibility of some rogues, in selling a deformed, useless, or diseased animal is incredible. If made like a cow, and showing the ill-breeding of a country horse, they will name to you some known good running Arab that had an ugly exterior, which is nothing at all to the purpose. A horse in his *tout ensemble* will often have an ugly appearance, and yet evidently show blood, and possess most of the best running points; one, however, whether handsome or ugly, that does not possess them, will never be a first-rate. If the legs are crooked, or the feet long, another runner will be named that had these faults, and they will affirm positively, it was in consequence of these defects, not in spite of them, and from their other very superior qualities that they happened to run well. If the shank is as large and round as a pint bottle, they will put their hand down to demonstrate to you the circumference, say-

ing, "There's strength of limb!" A bunged, wind-galled leg will become clean and wiry; a bull-neck will fine away; a herring-gut will turn into a beautiful barrel; a heavy shoulder into a light one; and a contracted and long foot will both open and shorten, when he is properly shod; in short, a brute will in a month be perfection, if you will only hand over your rupees. All this spluttering balderdash proves, either that they know nothing at all about a horse, or else that they are trying to deceive you: the former is the most charitable supposition, and, no doubt, the correct one, though a very large share of the latter is always mixed up with it. When, therefore, you hear arguments so truly nonsensical, and thinly-veiled as these, the sooner you unfold your knowledge of human nature, if you have none of a horse's, the richer you will find yourself in pocket.

Another deceptive mode of talking, shallow enough certainly, but peculiar to the Bomb Proof, is in this way;—When the stride is found shortened and clumsy, from having been overweighted by riding or carrying heavy burdens—a very common occurrence—the answer is, "Oh, that was done in Arabia;" or, "he was always so." Well, if over-weighted in Arabia when he was young, and the action injured in consequence, the chances are a hundred to one against his ever recovering it, unless a very powerful horse, and always after ridden by a light weight of seven or eight

stone: and if he was by nature always so, why, then of course, it is incurable. When the eyes are dim, or cataracts forming: "Oh! that's only from the heat of the stables." When the legs are injured, and the back sinews all bowed: "O! that's only from the heel-ropes, or a tent-peg:" just as if it made any difference to the purchaser whether the eyesight was injured by the heat of the stables or a pitchfork; or the legs ruined by a tent-peg, or a sledge hammer. Again, if a colt is produced that is badly formed; "Oh! he's only a colt, he'll alter in another year." To be sure, all colts alter, but if he is well-formed at three years old, he will be well-formed at twenty; and if badly formed, and want of due proportion at three years old, he will be the same as long as he lives. Now, unless you have been educated at Doncaster, or dubbed yourself a judge, self-constituted, as I have done, I strongly recommend, especially if you are a New Arrival, and have any regard for "your order," (letter of credit, I mean,) to ask some friend to accompany you in your rambles through these stables. Should you purchase from "gentlemen judges," the only chance of escape you have, is candidly to confess that you are only in the elementary instructions of the "Griffin's Aid-de-Camp," and then it would be ungenerous and unfair indeed to deceive and take advantage of you; but if you are a judge, (a question I always like to

ask, if I can do so without giving offence,) then you have no right to complain of being outwitted: you dubbed yourself; the more shame for you to acknowledge being so quickly dismounted.

Remember, however, that though a good Arab may be found without all the points I have described, a bad Arab will never be found with them; and also, that a horse may be of the highest caste, and yet have no great speed, in consequence of some faultiness of build, or being more adapted for the purposes of parade. A charger, racer, and carriage-horse, cannot be all of perfect build in one and the same skin: you cannot expect to obtain all these qualities in one horse, though they are much more nearly procurable in the Arab than the English breed. A draught horse seldom makes a good riding horse; you must make up your mind what you want, and be thankful if you get it.

For any purpose except the turf, always choose make before blood. A three-part bred Arab, of the proper form, will be preferable for either charger or hunter, to better caste faultily built; and for the carriage, half-bread, if well made, will also be better than more blood with less shape. An unexceptionably formed Persian or half-bred, will always bring a fair price, but a well bred, if faultily formed or deficient in appearance, will often be objected to, unless he has

speed.* Yet, let no man presume to quote from my book who has not well learnt by heart.

The Griffin's Catechism.

What gives a horse endurance ?—Blood.

What gives a horse speed ?—Ditto.

What gives a horse beauty and symmetry ?—Ditto.

What gives a horse a thin skin ?—Ditto.

What gives a horse a straight croup ?—Ditto.

What gives a horse large thighs ?—Ditto.

What gives a horse a well-formed large hock ?—Do.

What gives a horse a light neck ?—Ditto.

What gives a horse a large eye ?—Ditto.

What gives a horse a wide jowl ?—Ditto.

What gives a horse a thin open nostril ?—Ditto.

What gives a horse a deep chest ?—Ditto.

What gives a horse a fine forearm ?—Ditto.

What gives a horse flat shank-bone ?—Ditto.

What gives a horse a wiry limb, and large back sinews ?—Ditto.

What is the real meaning of blood ?—Blood.

Blood is shown in Premier and Perfection, pp. 69 and 70 ; Half-bred, in Water Proof, p. 70 ; and no

* Of late years, many horses with an English caste about them have found their way into the stables, and are called Arabs when well built, with a fine quarter, open jowl, and large eye; and Persian when otherwise. The opinion seems to be, they are Arabs, or Persians, or some mixed Eastern breed; but some few certainly look as if they had a dash of English blood in their veins.

Blood, in the Frontispiece. If you are so dull of comprehension that you cannot understand the meaning of it from all these drawings, assisted by all the foregoing explanation, confine yourself to a tattoo or screw; you were never born to display Perfection.

To decide upon the exact quantity of breeding every individual horse may possess, is another affair. The numerous shades of difference from the lowest caste up through the half and three-quarter bred to the highest caste, complicated as they are rendered by a greater or less degree of good and indifferent build attached to each, makes that impossible; it can only be known to a certainty by the breeder, where the pedigrees have been accurately noted down from generation to generation; and in India, so long as all pedigrees are taken upon trust, these intermediate shades of distinction must always remain a matter of opinion; but I do not hesitate to say, that if you are unbiassed, one month's study and reflection of the first part of this book will, as far as this blood or caste, and the build are concerned, enable you to purchase for yourself without disappointment.

A high-caste Arab, for a Racer, Charger, or Hunter, should therefore possess the following points:

Long quarter; straight spine; well set-on tail, and not too bushy.

Broad across the haunch-bones, but broader at the thighs.

Large and muscular stifle.

Large and broad lean hock, with the bone at the point long.

Lean head; broad forehead; large, full, bright eye; open thin nostril; deep mouth; evenly-closed lips; wide jaws; clean channel; and small erect ears.

Neck, rather long and light.

Withers, whether high or low, should rise above the top of the shoulder-blade.

Back, straight, rather long.

Long and slanting shoulder.

Short arm, and legs well forward.

Moderately broad and flat chest.

Great depth of girth.

Some distance between the last false rib and the haunch-bone.

Round barrelled carcase.

Large elbow-bone.

Long and muscular forearm, that there may be shortness between the knee and fetlock.

Broad square knee and large hinder bone.

Suspensory ligament and back sinews strong, wiry, and far apart.

Shank-bone flat; small round in front.

Pasterns long, but not too oblique. The point of the toe under the point of the shoulder.

Sound, well-shaped, black fore-feet; rather upright in front of the crust; never lengthening out more than the angle of forty-five degrees; open heels; round frog; and a concave sole.

Charger.

Long quarter; straight spine; well set on tail, and not too bushy; but, for a handsome charger, the tail should be carried rather high also.

Broad across the haunch-bones, but broader at the thighs.

Large and muscular stifle.

Large and broad lean hock, with the bone at the point long.

Lean head; broad forehead; large, full, bright eye; open thin nostril; deep mouth; evenly-closed lips; wide jaws; clean channel; and small erect ears.

Neck rather long and light; but a fine crest, and well-carried head, are indispensable.

Withers should be high for a charger.

Back must not be too long, or he will never pull up properly on his haunches, without which he cannot be a good charger.

Long and slanting shoulder; but it should have a little more substance than the racer's, so that, with

high withers, added to a good girth and fine chest, the saddle may keep back in its place; for the charger that does not carry his saddle well is a nuisance.

Short arm, and legs well forward.

Moderately broad and flat chest.

Depth of girth not actually requisite to the same extent as in the racer. A good caste Arab is never very deficient in this.

Not much distance between the last false rib and the haunch-bone. A charger should be well ribbed home. Being well "coupled up" is a sign of strength and endurance.

Round barrelled carcase.

Large elbow bone.

Forearm quite as muscular as the racer, but no harm if there is a trifle more length between the knee and fetlock, for a charger must lift his fore-feet well up, and not move like a daisy-cutter, or over stony ground he will soon come head over heels.

Broad square knee, and large hinder bone.

Suspensory ligament and back sinews strong, wiry, and far apart.

Shank-bone flat, small round in front.

Pasterns should not be so long, nor quite so oblique as the racer's. The point of the toe in its proper place is most essential.

Sound, well-shaped, black fore-feet; rather upright in front of the crust; never lengthening out more than the angle of forty-five degrees; open heels; sound frog; and a concave sole, are indispensable. A racer on soft turf might get on better with an indifferent shaped foot: have any fault in preference to an indifferent one for a charger. A bad foot can be worth nothing to either.

Hunter.

The nearer he comes to the description of a Charger the better. There is no necessity for his being so high or so handsome. Hog-hunters generally prefer small horses. Great strength, however, in all his limbs, with a moderately broad and flat chest, and circular carcase, is essential.

The neck must have the curve where the head is set on; the mouth deep, and the jaws wide, that he may be light in hand.

The back should be short (yet not a too short horse), but the spine straight.

The withers must be as high, and shoulder as slanting, with as much substance (not thick and cloddy) as the charger's. These two points, with a fine chest, are most requisite, both in the charger and hunter, that the saddle may be carried in its proper place. The charger is always kicking, and his tail becoming

galled by the crupper, when the saddle presses forward, and the hunter, in a leap, not having a crupper, may have the saddle and driver thrown on the withers, and fall topsy-turvy in consequence.

The distance between the knee and fetlock need not be so short as in the racer; for the hunter (though not required to have the high action of a charger) must lift his legs well. This is a point, however, that is seldom found too short, and in a racer never. As high caste and excellence of build can very seldom be obtained, and quite as seldom afforded, attend.

In the Hunter, first to the fore parts: the well set-on head and neck, slanting shoulders, well placed fore legs, and good, open, tough feet.

In the Racer, first to the cast and the hind parts.

In the Charger, as you value your neck and comfort, attend to every part. Money here is well laid out: he should be faultless: his name perfection, and his motto, *Fier, mais sensible.*

The annexed drawings will serve as a good index to the foregoing description. The second is a beautiful model of the proper form of a Light Dragoon Charger, and an excellent cut as a general runner of all races. The Racer and Hunter may look very different to the same class of horse in England; but though the points should be the same, the Arab,

in consequence of his stature, which in some of the best runners has not exceeded fourteen hands one and a half inch, and from his having inherited nothing but his native blood, assumes a very different appearance.

A Cavalry horse must partake of something of this second form; for a charger with a long back and straight shoulder, that cannot be brought on his haunches, is as ridiculous as a racer with a bull-neck and a camel's hind quarter, that cannot be got into a gallop; both being about as useful to answer their trades as a Newmarket carrier-pigeon without wings; and yet, such are continually purchased by Mr. Green and his brother; and the former not unfrequently sent to the light cavalry. Another description chosen for troopers, is the low Persian, or Gulph horse, often without any breeding at all, sometimes having a roach back, and consequently deformed quarter, drooping to the elegant angle of fifty degrees; such a brute can neither be useful nor ornamental for any purpose. A half-bred Arab, or well-formed Gulph horse, of fourteen hands one inch high, may do very well as a charger for a light weight of eight stone; but a good Kattywar, with his handsome crest and high action, is worth a hundred Gulph fourteen hands one inch tattoos. Size, strength, and activity, with fresh strong legs, tough feet, and open heels, are the grand desiderata for the cavalry.

Breeding the cavalry horse is foreign to my present purpose, but no man should be entrusted to pair horses and mares, until he understands something of the proper build of both, and is also fully aware of the undoubted truism, that "Like will produce Like." The keeping a bad horse costs the Honorable Company's Government quite as much as a good one, indeed, more; for the curious-shaped animals that are sometimes sent as colts for troopers take five times as long to break in as proper formed ones would; and even then, they are always laming themselves, or annoying their riders. There is no more difficulty in breeding good shaped horses than bad ones, if the breeder has only a little knowledge how to pair: blood is not required; half to three parts bred is abundance of blood, but make is indispensable. Some few *lusus naturæ*, under the most scientific management, will, of course, be thrown; it is beyond the power of man to prevent these occasional freaks of nature; but nine times out of ten, if the horse and mare are adapted to each other, four-fifths at least of the good form and good qualities will be inherited. A bull-neck will run through a dozen generations, and a stallion with a bull-neck should never be bred from. A thick neck, in an otherwise fine horse, may be bred from solely for draught for the artillery, but a down-right bull-neck should be shot. A straight-shouldered stallion, if otherwise good, may be bred from, for straight

shoulders are required for draught; and if the shoulder has only good depth, and crossed with an oblique-shouldered mare, this point may occasionally alter sufficiently for the cavalry; but not so certainly as if the obliquity was on the side of the stallion. A wall-sided stallion, if not very flat-sided, and otherwise good, may be bred from, for if put to a well circular-barrelled mare, the carcase will always improve: the Godolphin Arabian was wall-sided. A drooping hind-quarter, if only muscular, may be bred from, for if crossed with a straight-crouped mare the quarter will never be bad. No small carcased or narrow-chested, or lanky-thighed horse or mare should be bred from; and no horse or mare with a very long back or badly set-on head should be bred from. I am speaking as to breeding for the cavalry. Regarding the head itself, I should not care what it was like: the less brains a troop-horse has the better; but I would not breed from a stallion with a small eye, neither would I ever breed from one with a very large yard. Every stallion must have harmony of proportion united to general substance, and never be over laden at the top of the shoulder blade-bone; and every brood-mare must also possess these qualifications, besides being particularly broad in the haunches. Another great error that is committed, is in the difference of the size of the horse and mare. Not more than an inch and a half difference in the height should ever be allowed, and

even that is too much, unless the manager has proper discrimination with regard to the form : for instance, put a very fine, substantial, well-built stallion of fifteen hands to a rather slight but yet well-formed mare of fourteen hands two inches and a half ; the chances here, with make on both sides, or ten times as much against a symmetrical produce as if the height had been the other way, and a very fine, substantial, well-built stallion of fourteen hands two and half inches had been put to a rather slight but yet well-formed mare of fifteen hands : these last may, with some little propriety, be termed nearly even-sized horses. The only true method of increasing the size of the Indian horse, and at the same time insuring symmetry, is never to allow more than an inch and a half difference in the height, and only that under the restrictions just mentioned. An indifferent point in the stallion must also always be met by a very superior one in the mare, and *vice versâ ;* but the cardinal points of both must be perfect, or there will be no improvment in the breed worth speaking of. This is attended with no difficulty, and very little expense, when choosing half and three parts bred horses. There are hundreds to be had : good feeding and proper care of the colts and fillies would then in a very short time amply repay an establishment. English, Arab, and Kattywar horses and mares, judiciously chosen and crossed, would, in five generations, or thirty years, yield a breed that

would pay a hundred per cent.; but when a huge, faulty stallion, whether thorough-bred, passing for thorough-bred, or half-bred, is put to a coarse, country, drooping-quartered, and perhaps crooked-legged mare, and this mare, two or three inches smaller than the stallion,* what can be expected? It would be contrary to nature to find a good produce.

A half, or three parts bred horse, may often be found with all the good points of a thorough-bred, save three or four, and the want of these three or four shall indubitably prove him no thorough-bred. In lieu of the large brilliant eye, thin skin, small, flat shank-bone, and large back sinews, substitute a smaller eye, a thicker skin, a larger round shank-bone, and small tied-in back-sinews; who will then believe him thorough-bred? Or, to take other points, in lieu of the clean, wide jowl, thin open nostril, deep mouth, and large muscular hind quarters, substitute a closed fleshy jowl, a thick shut nostril, a heavy-lipped mouth and hind quarters, deficient of thigh muscle; and who will then believe him thorough-bred? With faults like these, he must gallop a mile and a half in very good time before you will obtain thorough-bred price. Even one single faulty point will frequently enable a


* Professor Coleman stated in his lectures (I have not seen it in print). that trying to increase the size of the Indian horses by crossing with the large English stallions was ruination to the breed, and that no well-proportioned foals could be expected.

good judge to detect a flaw in the blood. And this brings me to a question I have often heard mooted. When two horses are brought out and sold, each showing externally every point of thorough breeding and good build in exact equal proportion, and both of equal energy and equal supposed good constitution, how is it that one turns out far superior to the other, and how is it to be distinguished? To distinguish at the time of purchase between two so exactly equal is impossible; but supposing every point was allowed by the best judges to be exactly equal, and we could positively know that the constitutions of each were exactly equally good and strong, still, in India we are ignorant of their blood—of their line of descent: and even if we knew that as correctly as they do in England, still the solution of the question is impossible; but there may exist a closeness of fibre about the one, and a finer cementing together of the different parts of the frame, which scientific dissection might detect. The living proof in these cases, although we must wait a short time for it, is nevertheless a true proof, viz., the greater speed and *endurance* shown on the turf after each has had six months of proper training. Where you have the running build and every other external visible point in equal good proportion, and the training, riding, &c., equally good, then the turf, and the turf alone, furnishes the only living proof of the better blood, and this blood, this


latent energy for work, is comprehended with very good accuracy by persons who unite experience to a knowledge of the structure of the horse.

Another abstruse question is, why this superior blood loses itself in a greater ratio when crossed with inferior blood in good build, than with good blood in inferior build, when build itself is one of the best proofs of good blood? This does hold true, but only to a certain extent: that highly-finished interior mechanism of some internal parts, in whatever it may consist, does, beyond a doubt, in many horses with external indifferent conformation, excel in point of endurance that of inferior blood in a better external conformation. But some parts of external conformation denote blood more than others; for instance, take one horse with every point perfect, and then change his large brilliant eye to a small one; take another with every point perfect, and then change his straight forelegs to crooked ones, or make the chest too narrow: the former proves more a loss of blood, the latter more that of build; and the odds would be three to one in favour of the latter for a race. All these little minutiæ are of material consequence to breeders of horses, or men on the turf; but practice, added to study, and the possession of a good "eye for a horse," can alone give them. As beginners; do not devote too much time to any speculative points,


but first pay attention and learn the *externals:* the foal with any bad blood, will, to a certainty, when grown up, show some external visible flaw, such as a small eye, thick closed nostril, bad ear, small back sinews, &c., or very probably three or four of these. The points that denote pure caste, or thorough blood of long descent, are given at p. 3, together with the other requisites that enable a horse to gallop; but every good point therein enumerated is rarely attainable.




PREMIER.



PERFECTION.
'Fier, mais sensible.'



WATER PROOF.



OUTLINE.

Showing different parts of the horse, also the seats of disease.

A to B, the back; B to C, the loins; C to D, the croup; C to E, the quarter; E to F, the thigh; G, the stifle-joint; G to H, all that below the stifle; C to H, taking in the thigh, stifle, and all that below it, is sometimes called the whole quarter; I, the bone at the point of the hock; J, the shoulder blade-bone; K, the point of the shoulder; K to L, the arm; L to M, the fore-arm; N, the elbow-bone.

1, the place of the side, or lateral cartilages; when they ossify they lose their springy feel, and become hard as granite; they are then called side-bones; by some, ringbones; 2, the seat of ringbone in front, just above the coronet, seen in the fore as often as the hind pasterns; 3, the place enlarged, from being tied round the pasterns, close on the seat of ringbone; 4, the place often seen ossified on the outside of the pastern; 5, the place often seen ossified on the inside of the pastern; 6, the place that enlarges from sprain of the back sinews; 7, the seat of bone-spavin on the inner side of the lowermost part of the hock; 8, the seat of bone-spavin, a little higher up, and more towards the back part of the hock; 9, the seat of bog-spavin; 10, the seat of thorough-pin; 11, the seat of curb; 12, the place that enlarges from sprain of the back-sinews of the hind leg, seldom met with.

PART II.

THE AGE.

The teeth are forty in number: twenty-four grinders, or double teeth; twelve nippers, or single teeth; and four tushes.

The Colt's grinders.

The colt, at his birth, has eight colt's grinders, four above and four below; that is, two above and two below, on each side of his mouth.

At a month old he has four more colt's grinders, two above and two below; that is, one more above and one more below, on each side of his mouth.

These are subsequently shed, in the same order as the nippers.

At one year old four more grinders come, two above and two below; that is, one more above and one more below, on each side of his mouth.—Not shed.

At a little before two years old four more grinders come, two above and two below; that is, one more above and one more below, on each side of his mouth. —Not shed.

At about four years old four more grinders come, two above and two below; that is, one more above and one more below, on each side of his mouth.—Not shed.

Change of the Colt's Grinders for Horse Grinders.—At two and a half years old four colt's grinders fall out, one above and one below, on each side of his mouth, and four horse grinders come in their stead.

At three and a half years old four more colt's grinders fall out, one above and one below, on each side of his mouth, and four more horse grinders come in their stead.

At four years and a half old four more colt's grinders fall out, one above and one below, on each side of his mouth, and four more horse grinders come in their stead.

The Colt's Nippers.

The colt, at his birth sometimes, but always at a fortnight old, has four front colt's nippers, two above and two below, in the front of his mouth.

At a month and a half old, he has four middle colt's nippers, two above and two below, one on each side of each front nipper.

At eight or ten months old,* he has four corner

* Generally at nine months.

colt's nippers, two above and two below, one on each side of each middle nipper.

Change of the Colt's Nippers for Horse Nippers.—At two years and a half old, the four front colt's nippers fall out, and four front horse nippers come in their stead. When they are well up, he is three.

At three years and a half old, the four middle colt's nippers fall out, and four middle horse nippers come in their stead. When they are well up, he is four.

At four years and a half old, the four corner colt's nippers fall out, and four corner horse nippers come in their stead. When they are well up, he is five.

These colt's nippers take about four months falling out, commencing two months before the half year, and finishing two months after it. From the day the last corner nipper falls out the colt becomes a horse, and the filly a mare.

The Tushes.

From three and a half years to four and a half years old, the four tushes come, two above and two below; that is, one above and one below on each side of his mouth, and never change.

The mare has seldom any tushes.

Marks that show the Age.

At six years old, the marks in the two front horse nippers in the lower jaw are worn out.

At seven years old, the marks in the two middle horse nippers in the lower jaw are worn out.

At eight years old, the marks in the two corner horse nippers in the lower jaw are worn out.

Remarks.

As the teeth of horses grow very differently, and the marks are retained much longer in some than in others, it requires a little careful inspection of a few different mouths to enable you to decide at all correctly. The common rule of concluding that a horse is six, when the marks in the two front horse nippers in the lower jaw are worn out; that he is seven, when the marks in the two middle horse nippers in the lower jaw are worn out; that he is eight, when the marks in the two corner horse nippers in the lower jaw are worn out, will occasionally deceive you, if taken solely as a guide; for a horse that is always fed on dry grain and dry grass, wears out the marks quicker than one that is always fed on soft grain and green grass. The groove in the inside of the tushes that gradually fills up as he advances in years, varies so much in different horses, that it is no better a guide; for in some colts they come before three and a half years old, and in others not till after four and a half: in some horses they are very little blunted at the points at twelve, and in others they are blunted

at eight. The length of the teeth is no surer a test, for they are sometimes found short at twelve, and long at ten.

If a horse that is known to be ten or eleven years old, has the marks remaining in the two corner horse nippers in the lower jaw, it is sometimes concluded from this that he has been bishoped. If at this age the tushes should happen to be sharp at the points, and the teeth also short, like a young horse, it no doubt would be a good opportunity to endeavour to deceive us by bishoping a couple of marks in these two corner horse nippers, if they should be worn out; but the stain on such a tooth, somewhat similar to that burnt on the bone handle of a knife to give it the resemblance of tortoiseshell, will often expose the cheat of itself. A little round hole, made with a pointed iron, as well as the mark of a file, with occasionally a very small particle of the outer side of the tooth chipped off, is sometimes distinguishable at one of the "on my honour" stables* at Bombay. I have, however, seen a troop horse of eleven years old with the marks remaining in the corner horse nippers, the tushes rather sharp, and the teeth rather short, giving to him the appearance of a seven-year old horse; but the teeth were not quite so straight down in the mouth as they ought to have been; they projected


* Figging is also very common here.

a little obliquely forward, and they were very yellow: this furnishes a better criterion in these doubtful mouths.

To discover the age, therefore, examine carefully first, the marks in the teeth; secondly, if the tushes are small and sharp, or grown round and dumpy; thirdly, the length of the teeth; and, lastly, the position and colour of the teeth.

Up to six years old, there is not much difficulty in deciding; and from six until eight, if you balance all these four rules, not taking the marks solely as a guide, you will generally tell within six months. After eight, it is all guess; yet, when the tush is round and blunted, and the marks in the two upper front nippers are gone, you may conclude he is turned off nine, though marks should remain in the lower jaw; but the upper front nippers are, now-a-days, also sometimes bishoped, and I have seen a horse of sixteen sold for a seven-year old. At ten, the marks in the two front horse nippers in the upper jaw; at twelve, the marks in the two middle horse-nippers in the upper jaw; and at fourteen, the marks in the two corner horse-nippers in the upper jaw will sometimes, at these respective ages, also wear out regularly; though these cannot be depended on like the lower jaw: but a horse whose teeth are at all long, and at the same time yellow, and projecting a little

obliquely forward, you may be sure is no chicken. At ten years old, the lower horse-nippers also begin to lose their oblong shape, so that at fifteen they are nearly triangular from the front of the mouth backward. *(See sketches below.)*



[Bishoping is extensively and successfully practised by Arabs, Persians, Caboolees, Candaharees and the Mahomedans of this country. I have known several who prided themselves on being able to tell a horse's age correctly, sadly taken in by pronouncing horses that were 10 or 12 years old, to be no more than 6 or 7, from taking artificial, for natural marks. Independent of the shape and length of the teeth in old horses, the imposition of bishoping is easily detected by those who understand the construction of them.

The crown of a permanent tooth just cut, is hollow and covered with enamel, which, from attrition, gradually wears away, and in about 3 years' time the mark becomes obliterated, when the tooth presents a smooth surface, the centre of which is yellow, and composed of bone. The artificial mark is now made in the centre of the tooth, and has not a particle of enamel about it.

In order to tell a horse's age accurately, it is necessary that mouths of different ages should be examined and compared with each other; and I would recommend such as are interested in this subject to provide themselves with a copy of "Mayhew on the Horse's teeth."—ED.]

As a horse grows old, the pit above the eye deepens; but this will also be found sunken in one that is much

emaciated, and it is thought to be observable in those that have been got by old stallions.

The comparative ages between horse and man have been estimated at about the following comparison: one year of the horse to every four of man being reckoned from the age of two up to that of ten, and after that less. Thus,

A horse at	2 years	is as a boy of	8 years.
„	3	as	12.
„	4	as	16.
„	5	as a man of	20.
„	6	as	24.
„	7	as	28.
„	8	as	32.
„	9	as	36.
„	10	as	40.
„	15	as	50.
„	20	as	60.
„	25	as	70.
„	30	as	80.
„	35	as	90.

THE PULSE.

It really is unpardonable in some freshmen taking on themselves to decide on the complaint of a horse, and then recommending a treatment for him, without ever feeling the pulse. It is cruel to abuse, and half

destroy a horse's constitution, in the dark, in this way. There can be no difficulty in "catching" the pulse, if you put your two forefingers under the upper part of the lower jaw, and press the artery very gently against the bone. If the channel is not clean, but much filled up and fleshy, a hard pulse may feel softer than it actually is; but taking out your watch will always tell you if it is too quick; and, after feeling one or two horses under disease, you may be able in some degree to allow for this thickness of skin and flesh.

Thirty-five to near forty is the standard pulse* (young colts two or three quicker); and when it beats about this number in the minute, going off with a full bounding feel, it is the pulse of strong health in a strong constitution. Forty-five, is too quick; thirty, too slow. When the pulse is under forty-five, there can be no active inflammation that will not quickly show itself.

When it is weak and not full, it bespeaks more or less of debility, according to other symptoms that may be present.

When it rises to sixty, there may be fever, or some local inflammation, according as other symptoms are present.

* This is correct as regards English and other large horses, but in the little Arab, the healthy pulse is from forty to forty-five, and in young colts two or three quicker.—Ed.

When at the number of seventy, or eighty, or more, it is hard, and yet small and wiry. It denotes inflammation of the bowels, if the other symptoms are present.

When it is oppressed and indistinct, feeling as if there was too much blood in the artery, it denotes inflammation of the lungs, if the other symptoms are present.

When the pulse is irregular or intermittent, stopping a second, then going on again, it denotes great danger under disease, if other bad symptoms are present. The intermittent pulse is sometimes produced intentionally, by giving digitalis, or other poisonous medicines.

BLEEDING

should never be resorted to till the disease is clearly ascertained.

When you bleed from the neck vein, the first six inches near the head must not be punctured, nor lower down the neck than a foot. Any spot between six and twelve inches will do. The vein branches off about six inches below the jaw, and the best spot for bleeding is two inches below that.

If you have never bled a horse before, or the vein does not show sufficiently full, smooth it down and press upon it below with the fingers, then tie the cord round the neck and lay the fleam along the course of

the vein, never crossways, gently fixing it without penetrating the skin. Having blindfolded him, do not shut your eyes and jerk the fleam when you strike, like a boy pulling the trigger of a gun, or you have no more chance of hitting the vein than he has the bird; but, keeping your eyes fixed on the fleam, give it a sharp but not a severe knock with a blood-stick or tent-peg.

Some people cannot bleed, which is unfortunate in case of inflammation. Perhaps you may be more successful with a lancet. Having first tied a piece of cloth round the bottom, or holding it tight between your forefinger and thumb, that it may not go in deeper than the fleam would, insert the point gently into the vein, and ram it in a little upwards, so as to make an orifice full half an inch in length.

The beauty, and the great effect, of bleeding a horse in the neck vein, particularly under inflammation, consists in taking a large quantity of blood in a short time; therefore both veins in these cases should be opened at once, and the larger orifice the fleam makes the better. No fleams from Mr. Long, or any good maker, are made too broad-shouldered; if made too deep, the artery underneath may be penetrated, and the horse lost. Having got the quantity of blood you want, do not let the flow stop quite suddenly, or air may get into the vein and half kill him; but

loosen the cord, if one has been on, gradually; then close the wound immediately, and that without pulling the skin from the neck, by which blood is apt to get underneath and cause a bad swelling; pin the two edges together, taking care not to include any of the hair between the lips of the wound, with the smallest possible pin, or needle, which is better, not an inch and a half skewer; and do not forget to tie the head up for an hour, and to see that he does not rub the part afterwards, or inflammation and loss of the vein may follow. The pin should be gently drawn out two days after, first cutting the horse-hair or thread tied round it with a pair of scissors. If the blood does not flow sufficiently free and quick, keep his mouth in motion by putting your fingers in at the corners, and throw hot water over the loins, if needed, or put a sheep-skin or half-a-dozen jhools over them.

Take care that your fleam and lancet are clean, and not rusty, and always ascertain how much the koondee holds that is to receive the blood, or you may take away a gallon and a half for a gallon, which I have often seen done.

The blood should fall in a stream into the middle of the koondee, or nothing can be judged from the colour of it afterwards; and it must rest undisturbed in the shade for half an hour, which is about the time

it takes to coagulate. The appearance, however, the blood puts on after its coagulation, and the coagulation being slow or quick, is so exceedingly complicated an affair, and influenced by so many causes, that no criterion as to the degree of inflammation can be gained by it, at least by us; the great thickness of the buffy or sizy coat at top being present in a state of health as well as under disease; and often none at all existing, though the blood may be darker, when the inflammation is greatest. When, however, the blood has become solid, after being drawn in a full stream, and is found with or without this sizy coat, yet at the same time thin or watery, it will, perhaps, more correctly denote that it ought not to have been taken. When bleeding is necessary, you will find it under the proper head; so, if you take on yourself to bleed for imaginary purposes, you deserve to suffer for it.

In taking a large quantity of blood from a horse labouring under any inflammation, always take care there is a good soft bedding in case he should faint. The pulse will first falter; and he will generally begin to droop and stagger a few seconds before he falls, which, of course, must be the signal to desist, and then ease him down gently, if possible. The faintness will go off in a few minutes; but he must be left to rise at his own pleasure.

Bleeding from the forearm vein, inside the forearm and from the thigh vein above the hock, should be done with the lancet, it not being driven in so far as in the neck vein. Bleeding from the eye vein, two inches below the inner corner of the eye, must be done with a very slight puncture, or the point of the lancet will be broken against the bone.

To bleed from the toe, you have only to cut away the sole underneath at the toe, where the crust and sole join, until blood comes. When the sole is cut away underneath with the drawing-knife, take a fresh sharper one, or a clean "searcher" to cut through the vessels. If you cut deep enough, you will generally get the quantity of blood you want. A little tow or cotton is then to be stuffed into the hole, and tar over it, and a bandage wrapped over all for four or five days.

[Blood-letting, fortunately for man and beast is not carried out to that extent it used to be; indeed many practitioners have very properly discarded it altogether, and now depend upon sedative medicines to control the heart's action which they do most effectually, without prostrating the system as bleeding does. I have tried all the medicines in the Veterinary Pharmacopœia, under the head of "Sedatives," but find none so potent in their effect as Tincture of Aconite (Fleming's Strength). I have used it for all cases of inflammation with marked success, for the last five years, in doses varying from 20 to 30 drops, mixed up in a pint of gruel, and administered every 2 or 3 hours, till intermission of the pulse was produced.

A few years ago, depletion, to the extent recommended by

"Blunt Spurs," was resorted to by professional men, as well as empirics, without respect to the nature of the complaint, the animal was suffering from. The Lancet, Phleme and Blood-stick were then pocket companions and the vital stream was allowed to flow till water buckets were nearly filled, and the poor patient fell to the ground, unable to stand on his legs any longer. It was considered that unless such an effect was produced the operation could not be attended with any benefit, and many valuable horses have gone the way of all flesh, not from the diseases with which they were affected, but from the butchering treatment they were obliged to endure.

About 30 years ago, Dickson, in his lectures on the "Fallacy of the Faculty" raised his voice against the use of the lancet, but the practice was too popular to be put down at once, and a quarter of a century elapsed before the "Faculty" could appreciate one of the greatest benefits conferred upon mankind. In his lecture on Blood-letting he states—

"Out of upwards of twelve thousand cases of disease, that have, within the last few years, been under my treatment, I have not been compelled to use it once. Resorted to under the most favorable circumstances, its success is anything but sure, and its failure involves consequences which the untoward administration of other means may not so certainly produce. Have we not seen that all diseases have remissions, and exacerbations—that Mania, Asthma, Apoplexy, and Inflammation, are all remittent disorders! From the agony or intensity of each of these developments of fever, you may obtain a temporary relief by the use of the lancet; but what has it availed in averting the recurrence of the paroxysm? How often do you find the patient you have bled in the morning, ere night, with every symptom in aggravation! Again you resort to bleeding, but the relief is as transitory as before. True, you may repeat the operation, and repeat it until you bleed both the blood and the

life away. Venesection then in many cases may be a temporary, though two often a delusive relief. The general result is depression of vital energy, with diminution of corporeal force.

"I have not always had this horror of blood-letting. In many instances have I formerly used the lancet, where a cure, in my present state of knowledge, could have been effected without: but this was in my noviciate, influenced by others, and without sufficient or correct data to think for myself."

After enumerating several fatal results from blood-letting he concludes thus—

"Such instances, too numerous to escape my observation, naturally led me to ask—Can this be the proper practice? It was assuredly the practice of others—of all, could all be wrong? Reflection taught me that men seldom act for themselves, but take, for the most part, a tone or bias from some individual Master.

"By education most have been misled,
"So they believe, because they were so bred."

"I had the resolution to think for myself—ay, and to act; and my conviction, gained from much and extensive experience, is, that all diseases may not only be successfully treated without loss of blood; but that blood-letting, however put in practice, even where it gives a temporary relief, almost invariably injures the general health of the patient."]—Ed.

PHYSICKING

takes six days: two for preparation; the day it is given, and the day on which it operates; and two more to return to the former diet: but on an emergency, or when the dung appears softened, four days may be sufficient: one for preparation; the day it is given,

and the day it operates; and one more to increase to the former diet.

Whenever a dose of purgative physic is to be given to a horse in consequence of being out of condition, or previous to being put in training, or when in training, or coming out of training, it is most necessary he should have bran mashes, instead of his grain, and very little grass for the previous day, or two days, whether the physic is to be aloes or oil, a ball or a drench. If he will not eat bran mash, he must be well stinted of dry grass, and a little green (cut the previous day) given. There is a peculiarity of structure about a horse's inside, that, unless he is properly prepared and carefully looked after under physic, you not only will do no good, but, on the contrary, you will throw him some weeks back, and even run the risk of losing him; for it is surprising how suddenly a horse will sometimes "go out" under a dose of physic improperly given. You should endeavour to give your physic, for any of the above purposes, of that strength, that not more than ten extra evacuations will be produced. Some horses are easily operated on with a ball of four drachms of aloes; others require a large dose of seven. At one time in the year, a horse will sometimes purge with five drachms, that another time will not feel the effect of six. Four to six drachms of Mr. Sprague's or Treacher's aloes are

generally ample for any Arab horse, if properly prepared and properly exercised; but eight drachms may fail to purge, without proper precautions and exercise, and yet easily produce inflammation and death. Some horses, again, will purge in twelve hours;* others, not for thirty-six. The best and safest purge for a novice to give is a drench, and that drench composed of aloes and Epsom salts, or oil and Epsom salts: thus, if you think five drachms of aloes is required to purge, give three drachms, and five ounces of Epsom salts (two and a half ounces of salts for one drachm of aloes), dissolved in three-quarters of a pint of warm water, adding one ounce of salad-oil and a quarter of a drachm of ginger; or, if oil is to be given, a pint of linseed-oil and five ounces of the salts. These aloetic-balls prove occasionally most raking to the bowels of an Arab horse, and a large dose of them is always attended with risk. The above drench may be safely given, even when bran mashes are refused, provided the horse has been muzzled at night and the other directions duly attended to; and, with a clyster of salt and water on the same day, it will, with proper exercise, purge effectually, and rarely disagree. The following directions will guide you how to treat a horse under physic. Suppose it is to be given on Wednesday morning:—

* A little Arab I once had charge of used to purge in ten hours, though the physic was given in a ball; and in fifteen hours it had generally set.

Monday.—Give four mäps, that is, one bazaar seer of bran mash for his first-feed at eight in the morning, and only one pooly of grass after it. Four ditto at one o'clock and only one pooly of grass after it. Four ditto at evening's feed, and then allow him to eat grass until nine o'clock, when the muzzle is to be put on for the night. Give as much water as he likes to drink, but no more grass than this. He should have an hour and a half of walking exercise, both in the morning and evening, for which he may be ridden; or if strong, and in good health, may be gently trotted.

Tuesday.—Exactly the same as Monday, with another bran mash at nine o'clock, before being muzzled.

Wednesday.—At day-break let him have the physic, always washing it down with two drachms of common salt in half a pint of warm water, whether it is a ball or a drench; this facilitates the "working," and may prevent the horse feeling sick. Give five mäps of bran mash for his first feed at eight in the morning, but only a mouthful of grass after it. Five ditto at one o'clock, and half a pooly of grass after it. Five ditto at evening's feed, and one pooly of grass after it, when the muzzle is to be put on till nine o'clock, at which hour he may have another bran mash, before being muzzled for the night. The more water that is drunk the better, but on this day the chill must be taken off, whether it is the hot or the cold weather.

Walking, or very gentle trotting exercise, morning and evening, the same as on Monday and Tuesday.

Thursday Morning.—The physic, during exercise, will perhaps begin to work; if it does not,* you must feed exactly the same as the day before, for sometimes it will not operate for another ten or twelve hours. A mash and a little warm water, and another half hour's trot, will often bring the purging on; and, when it does come on, he may have bran mash again, a little dried green, or a pooly of grass, and water with the chill off. If only two or three evacuations take place, give more warm water and another trot, which will carry it off. But, though the medicine should fail entirely to act, you must not administer another dose, for fear of producing over-purging or inflammation. Some bran mashes must still be continued, with but little grain, for the following forty-eight hours; and then, after an interval of another four days, making exactly a week from the time the first dose was administered, a stronger may be given, preparing the animal the last two days as before.

Should the physic have been a drench,† precisely the same rules are to be followed throughout; but

* If it has not commenced operating by this time, give a clyster of warm water and Europe brown soap. This with exercise will generally cause it to operate freely.—ED.

† Have a drachm of aloes more than you are going to give, finely powdered so as to be ready in case of spilling.

the purging will in this case commence on the evening of the same day, or about sixteen hours after it is given. Whenever it does commence, he may have bran mash, grass, and water with the chill off. You can give these mashes warm, but there is no occasion for any extra clothing; and, if he has not been accustomed to any clothing at all, do not put any on, unless the weather is damp or cold. If he has been standing outside, he should, during these six days, be brought under cover; but never dream of putting him into a shut-up, close stable, like many of those in Bombay. Such is far more hurtful and dangerous than leaving him outside.

If the dose you have given was not a mild one, which it ought to have been if you were not aware of your horses's constitution, and he should purge more than ten times, give immediately a quart of lukewarm finest bajree, or wheat-flour gruel, with two drachms of gum arabic dissolved and put into it, and repeat this every two hours till the purging stops, or he has once dunged a little thicker. He is not to eat or drink a particle of anything else till this takes place, but remain quiet in a loose stall with a muzzle on, and allowed to lie down on his bed. If he will drink the gruel, let him take a couple of quarts each time; if not, drench him with one. A horse should not have a dozen watery evacuations, when the intention is merely to promote

condition or prevent illness, or, the object for which the physic is given will be defeated. This fine bajree, or wheat-flour gruel, and gum, will soothe the bowels, and in the course of six or eight hours generally stop the purging. Should it not be found sufficient, and he should still purge on to the number of twenty times and more, it will be time for you to look to OVER-PURGING, in a subsequent page of this volume.

When the physic has done working, whether it has been a ball or a drench, two days more are still required before he can resume his former grain, and be put to work; for the too rapidly filling of the stomach and bowels immediately after physic, often proves as dangerous as giving the physic without preparation. Two mäps of bran mash, and one of ground grain, can be given at each feed, and one pooly of grass after it, and at night he may eat away at grass till the muzzle is put on at nine o'clock: the water must have the chill off, as before. The next day the grain may be increased to two mäps at each feed, and the bran mash reduced to one; the water may now be given cold, and the muzzle left off at night.

I have particularized everything under this head, because I have been witness to numerous horses being greatly reduced in flesh, and also to severe gripes, and even inflammation and death taking place, solely from neglect of the above precautions. If you deviate one

iota from these directions, or do not see that they are minutely attended to, and your horse should sicken, and the prescription under Class I of Part IV fail to cure him, do not blame the Aide-de-Camp.

To make Bran Mash, and Bran Tea.

Mash.—Pour boiling water on sweet fresh bran, stirring it about; then cover it up until it is cold. Some horses dislike bran, especially if it is stale, or has been given them dirty at some former period. A seer mixed with their gram, at each feed, two or three days before you commence preparing for physic, is a good way of accustoming them to it. If he should refuse to eat the bran mash on either of the two first days, you may put half a mäp of soaked gram in it, put none on the day the physic is given.

Bran Tea.—One gallon of boiling water to be poured over every four mäps of bran, and strained off when cold. The bran is quite fit to be eaten afterwards.

To make fine Gram, or Wheat flour Gruel.

Mix two mäps of the finest flour with six quarts of boiling water in an earthen koondee: put it on the fire for eight or ten minutes, stirring it about. If merely mixed with water, it is liable to clot in the stomach and do harm, especially when given in over-purging; for this last case it must therefore always be made with the hot water, and properly boiled: or

keep at hand some previously made thus: boil two or three pounds of dry wheat bajree, or gram flour, tied up in a cloth like a pudding, for four hours: it will keep for months, and a piece being broken off, and mixed with boiling water, it is ready at once. No animal, and no man either, has so nice and delicate a taste for drink as a healthy horse; and if the water is the least smoked, or the gruel should get the least smoked when on the fire, or the koondee it is put into the least greasy or dirty, or even, perhaps, if the koondee is quite new, he will be sure to smell it, and most probably refuse to drink; and, if he has not occasionally had gruel before, will very likely never afterwards voluntarily swallow it: now, as this is of consequence, inasmuch as gruel is very necessary at times for a horse recovering from sickness, it behoves you to take care and offer it him clean. Many horses will turn their heads away even from clean gruel, when under physic, that at other times would drink a dhool full.

Muzzle.

Let your muzzle be at least eight inches deep, with two large holes, both full two inches in diameter at the sides close to the bottom, for each nostril; and the top of it large enough to put your hand in on each side. It is barbarous to see the way in which some horses are half suffocated with a little hard muzzle tightly drawn

over their mouths. Do not choke him either with the neck-strap: see that it is well and fast buckled, but not drawn tight round the throat. The neckstrap of the head-stall is also often tied or buckled so tight, that the poor animal is half throttled, and not being able to speak, loses his whole night's rest, in consequence.

Drenching.

In drenching a horse either with gruel or physic, a common soda-water bottle, or a thick English wine-bottle, with a piece of leather sewed tight round the neck, to prevent accident from the glass breaking, is cleaner than any thing else. Put the neck gently into the mouth just above the tush: if he is a quiet horse, and it is properly given, he will swallow down the most nauseous dose with as little trouble as a child takes its pap. If very restive, you can have it put into a thick leather bottle, which a Moochee in any bazaar will knock up for you in a few hours.

Physic ball, and drench.

Ball.—Finely powdered aloes, and a quarter the quantity of kidney mutton fat, that has been well boiled down. Make into a long ball, or two, if it should be large, adding a dozen drops of oil of anise-seed, and some calomel, if required. Wrap it up in tissue-paper, oiling the same with linseed-oil at the time of giving, for a ball cannot be too soft. A

hard ball takes double the time to dissolve, besides the danger of its sticking in the gullet, and choking the animal. If the aloes be soft and difficult to powder, mix in the mutton fat and warm them together over a fire.

Drench.—Put the aloes and Epsom salts into a pint bottle, pour three quarters of a pint of hot water over them, shake until *well dissolved,* and, when cool,

When calomel is required over night, previous to a ball or drench on the following morning, it is to be mixed up with a spoonful of linseed-meal and ghee, and given at five or six o'clock, on an empty stomach, before taking his walk. The bran mash and a pooly of grass may be eaten afterwards, at evenings feeding-time, and then the muzzle put on for the night.

Alteratives.

Nine to ten drachms of best aloes, one drachm of calomel, and thirty drops of oil of anise-seed, made up

* Some horses will hold a drench in their mouths for ten minutes without swallowing it.

† When aloes are given in solution, they should always be prepared as follows, aloes in small pieces one part, distilled water seven parts, proof spirit one part. Dissolve the aloes by means of the water bath, that is, put the aloes and distilled water in a bottle. Put the bottle up to the neck in boiling water in a pot or chattie and allow it to simmer over a slow fire for two hours, and when removed, add the spirit. The dose, as a purgative, will vary from four to six fluid ounces.—Ed.

with linseed-oil, and a tea-spoonful of linseed-meal; or, as under "PHYSIC BALL," and divided into eight balls, one to be given every morning at daybreak, and one also every evening about five, when he goes out for his walk, will answer this purpose, when they are required;* but an alterative drench morning and evening, of one drachm of aloes and two ounces of Epsom salts for each drench, is sometimes preferable: a drachm of Turkey rhubarb may be added to each drench for a thin horse. The same care is to be taken as under "PHYSICKING." He must have bran mashes instead of gram from the day the first ball is given, and be muzzled at night; the water always with the chill off, and only one pooly, or the same quantity of dried green grass after each feed, except in the evening, when he may eat grass until muzzled, at nine or ten o'clock. If he is thin, half a mäp of soaked or boiled ground gram may be added to each mash. Good walking exercise morning and evening, but no wet green grass, and no washing under physic. If they do not operate twelve hours after the last is given, and after you have given a trot, continue them on until they do; but clyster first, with one ounce of soap and

* I have always found the following an excellent alterative. Aloes one drachm, calomel ten grains, powdered gentian root two drachms, thirty drops of oil of aniseed, or tincture of ginger made into a ball with treacle or honey. One to be given morning and evening. When the bowels become relaxed, discontinue the ball for a day or two, and then go on with it again.—ED.

two drachms of aloes in a gallon of warm water, or the horse may be severely over-purged from the effect of the whole taking place at once.

When calomel is required in combination with alteratives to a greater amount than a drachm, it should be mixed in the four first balls, to prevent any chance of salivation.


If your horse will not eat bran mash, which is most probably your own fault, do not give laxative alteratives.*

General Alteratives.

Three drachms of black sulphuret of antimony, three drachms of sulphur, two drachms of nitre, and half a drachm of ginger, made into a ball with honey or ghoor, and given every night for a week or ten days, the horse eating partly-boiled food, and receiving only gentle exercise, is the best alterative for improving appearance when no particular complaint exists. It is also good for preventing plethora.

* A one-pound powder-canister once full, piled up, is about equal to a mäp, or kutcha seer; but whenever a mäp or seer of ground gram is mentioned, it does not mean a kutcha seer of whole grain taken and ground, that would be upwards of a kutcha seer and a quarter; for eight mäps, seers or powder-canisters full of unground gram, will make eleven when ground. A pooly of grass is about two pounds.

SHOEING.




A SOUND FOREFOOT PREPARED FOR THE SHOE.

A. The heel of the crust.
B. The toe.
CC. The quarters of the crust
DD. The bars as they should be left with frog between them.
EE. The angles between the heels and bars where corns appear.
FF. The concave surface.
GG. The bulbous heels.
H. Cleft.

Various kinds of shoes and nails cannot be constructed in India : there are neither proper forges, nor proper persons to superintend them. The pattern here given is easily made, and will be found to answer well for either a Racer, Charger, or a Hunter, and be

less productive of injury than a broader web at the toes, which, being never sufficiently beveled out, is consequently apt to press on the sole, besides being more liable to pick up stones. There is no occasion to discard it when it becomes old and thin at the toe; the older the better, if only strong; and wearing thin at the toe is no disadvantage.*

If this shoe is properly made; the crust that it rests upon evenly rasped; the sole properly pared;



FORE SHOE.


* Some of the French shoes are bent up at the toes when first put on, in the shape of men's wooden shoes; but to be able to judge of the comparative merits of different forms of shoes and nails, requires great study and great practice. Whether you adopt that which I have recommended or not,

and the foot, during the intervals of shoeing, always kept properly stopped and moist, you will most likely be able to avoid contraction; if you fail in any of these, you will have contraction.

It is five-eighths of an inch broad all the way round from toe to heel, and one-fifth of an inch thick all the way round from toe to heel. There is a groove sufficiently large and deep to receive the small oblong heads of the nails, which are to be driven in even with the shoe; and this groove is well away from the edge of the shoe, so that the nails shall be driven nearly as far in as where the crust and sole join—"just between wind and water"—and brought out about three quarters of an inch high up in the hoof. From the nail-holes on the inside, it is scooped out, beaten out, beveled off, or filed-away, to admit of a pricker being passed under to clear out all sand and gravel, and so that it shall not press on the sole. The inch at the heel that is not beveled off is to rest where the bar and

keep the heels open. In France, my horse's heels never contracted, and in England they never contracted: in the former country they were generally shod under the superintendence of a Government verterinary surgeon, and in the latter at the celebrated Mr. Turner's: neither in India have my horse's heels ever contracted, and here they have always been shod with this kind of shoe.

A pattern of the fore and hind shoe is lying at Dady's, in Bombay, which will show the groove, as also the form of the nail, and how the nail-holes are to be punched; and by which you will see if the nails are properly pointed; so that the Nolband will have some difficulty in pricking the horse even, if he tries.



HIND SHOE.

crust join, not quite touching the frog. There are six nail-holes, three on each side; and the nail-holes are to be punched with the Nolband's common punch, firstly, on the inside, lastly, on the outside; by which means they get a slanting direction outwards, and there will be no danger of the foot being pricked. These nail-holes must be punched while the shoe is hot, or the iron, which is rarely of the best quality, will split. Always see that the nails fit the holes before you put the shoe on; for if the heads of the nails are not small enough to go into the groove, or the

groove not large enough to receive them, they will be broken, and the shoe come off.

A horse with straight pasterns, on a very hollow foot, may have it made a little thinner at the heel than at the toe; and if the foot is black, strong, and well open, another nail may be put on the outer side for hunting, &c.; but, out of some hundreds of feet, I have found the outside very nearly as often contracted as the inside. The hind shoe has the nails rather farther back, and the toe is a little squared.

The weight of this shoe is about six ounces, and by reducing the thickness a little it will weigh about five ounces, which will be a proper training shoe. It must not project the least beyond the heel, but be there nicely rounded, or the hind foot in the gallop may possibly come in contact with it, and the horse be thrown, or the shoe wrenched off. The ground surface of the toe of the hind shoe should also be filed a little, to take off the edge, as a further precaution.

Cutting the Hoof.

The foot having been soaked in water to render it soft, the sung-turash is to be used very sparingly—not at all for a flat foot; for if too much of the bottom of the crust is first taken off, there may not be sufficient horn left in the sole to make it properly concave: then with the drawing-knife cut away the horn, till, by the

strong pressure of your own or the Nolband's thumb, you can feel the sole bend up towards the corners, as well as the fore-parts: this is the guide for all feet worth the expense of shoeing. If more is cut away, the horse will be tender, perhaps lamed: if less is taken off, he will require shoeing again in a fortnight. When the foot is not softened before going to the Nolband's, either by cow-dung stopping for the two previous days, or by being soaked in water two or three hours before shoeing, or by cloths wrapped round the coronet, and warm water poured on them for that time, a great deal too much horn may be cut away before the sole will yield. A fine flanky foot will often show of itself the proper quantity to be taken off; but others again would suffer; besides, the hoof being moistened, adds to the nails taking firm hold, and prevents the horn splintering; the foot, moreover, is shod in half the time, and the Nolband saved from a gash in his arm. The horn between the bars and the crust having been cut away with the drawing-knife, a full sixth of an inch of depth must be left up in the corner where the bar and crust join. Never cut away the horn between the bar* and the frog, but cut

* For mercy's sake! learn where the bars are, for the manner in which English farriers, as well as Nolband's, cut them away at this part is disgraceful. The English farrier calls it opening the heels; and I have seen them retrograde into this most destructive method of cutting, a few months after losing the supervision of a veterinary surgeon. The bars are the finish of the crust, of harder horn than the sole, and a different shade of

away a little of the very hindermost part of the heel, just the last quarter of an inch, so as to allow the heel of the frog, where the cleft is, to expand, and prevent the horn bending in there. Neither cut the frog unless it is ragged, excepting towards the toe; and only there, if it should be hard and higher than the heel.* Lastly, file till the crust and bars are even, leaving the frog, if possible, the eighth of an inch higher so that it shall be just within the level of the shoe. The toe should always be shortened as much, as it will admit of, and any unevenness in the wall rasped smooth. There should be room enough for the edge of a knife-blade to play between the shoe and the crust at the heel when on; and the shoe at the heel should extend the eighth of an inch beyond the crust on the inside, as well as the outside. You need not fear that this will be a cause of cutting: a horse cuts with the quarter, sometimes with the coronet; his legs must be very faultily placed to cut with the heel. The natural well-formed foot, also, is a smaller half circle on the inside than the outside, as shown at page 33; and this form is, of course, to be preserved.

colour; they never run down straight, but curve, as shown at p. 37 in the description of feet; and the place for opening the heels is not in the middle of the bars, where the Nolband sideways scoops them half way, but opposite the heel of the shoe *behind* where the bar and crust join.

* If the frog is deficient in height, after the heels have been lowered, tar must be daily laid over it. If naturally too high, which is not so often the case, it may be slightly pared.

Three-quarter Shoes and Tips.

Three-quarter shoes I dislike, for I think they lame as many horses as they cure of contraction: they lame, because they are rarely put on good feet, but on those that have contracted heels, in the hope that the heels will open without the horse being taken out of work; but it is only strong black heels that are tough enough to stand hunting, or battering on a hard road, when the shoe does not come well home. They seldom have much beneficial effect on contraction, because the heels are never sufficiently lowered when a three-quarter shoe is worn; and if the heels, when the pasterns are long, were to be sufficiently lowered, too great stress would be thrown on the back sinews, unless more horn at the toe is pared away, as I am about to explain, under Tips.

Tips are equally objectionable on this principle, and more so if the horse is to be worked (though not quite so liable to get imbedded in the foot.) But when a brittle hoof is laid up from contraction, a tip with two nails on each side, close to the toe, will save the crust without any impediment to the heels opening; yet there are two errors universally committed in India in putting on a tip. The heel of the tip is always too thick; it should slope down to as fine an edge as a dinner-knife blade, and the horn towards the toe should have as much extra pared away, after the foot is finished cutting, as directed under "CUTTING

THE HOOF," as the tip is thick ; then the horse is not thrown in the slightest degree more than natural on the heel, which is of great consequence with an oblique pastern. Secondly, a tip should always be narrow. A horse on soft turf, or Bombay sands, may be trotted, or gently cantered, or put in the lounge, with benefit to contraction, and no fear of strain, when tips are made strictly in this way ; and if put on good strong black feet, with open heels, they will prevent contraction taking place, and answer very well for ordinary riding, or even for hunting, in some of the sandy soils of India.

[In Madras the Native Smiths who make horses' shoes, can do nothing else, not even to the making of a door nail; but are generally speaking very expert at their work. The only exception, to my knowledge, is a man at Vellore who is a very superior Black Smith, and under the able instruction of a Gentleman understanding the horse thoroughly in health, as well as in disease, can forge and fit a shoe exceedingly well; but I believe the Gentleman in question had many difficulties to overcome before he could prevail upon his pupil to break through caste prejudices, and shoe a horse, as he is a Hindoo.

A Smith, on a salary of 14, with the assistance of a hammerman on 7, and a bellows boy on 3 Rupees a month, will in about 8 hours time forge 50 shoes for ordinary purposes, and you have only to give him a pattern, and he will produce any description of shoe you may require.

I endeavoured to get a couple of Natives to forge under the superintendence of a European smith, but it was of no use; they very reluctantly tried the work for a couple of days, complained

of pains in their legs, and said their fore-fathers worked sitting, and they did not see any advantage in standing up. I then got two East Indian lads out of the Male Asylum, they set to in right good earnest and in about a year's time proved themselves very fair workmen; but I must admit, that, contrary to my expectations, they could not turn out work in quantity or quality equal to the Natives, and as they suffered a great deal during the hot season, I withdrew them from the forge altogether.

The Nolbands are Mahomedans and when instructed by Europeans to use no other than drawing knives, they are really smart fellows at their work. I have two men, one 11 and the other 8 years in my employ, who think nothing of shoeing 8 horses each, and when there has been an unusual press of business, each has shod as many as 12 a day.

Very few people who possess horses attach the importance to shoeing that it deserves; they see that horses' feet are composed of horn, and conclude that as long as a horse is shod, it does not matter how it is done. They would soon alter their opinion, if they saw all that is contained within the horny part. There is nothing so beautiful nor so delicate in the structure of the horse, as the sensitive part of his foot. It is one mass of small arteries, veins, and nerves, and mismanagement on the part of the shoer, even to pressure on the parts, causes great mischief, by producing lameness, tripping, and injury to tendons, as well as suspensory ligaments.

Nolbands go travelling about with their bags from house to house, and they get a Rupee for each horse shod, out of which 4 annas are given to the Horsekeeper, but should there be such a functionary as a Coachman in the establishment, the Horsekeeper shares his perquisite equally with him. Take Blunt Spurs advice and have your horses shod under the best superintendence you can procure. A European or East Indian Farrier, who has gained his experience in the army is better than a Nolband.

Your servants will of course lose their fee, but your horses will be properly shod. They will play all sorts of little tricks, such as loosening a shoe, raising one or two of the clenches, so as to cause the horse to cut, they will even go the length of pulling off a shoe the day after the horse has been shod, and unless you can make them see that you are quite up to them, by levying a fine, whenever you are satisfied that they are deceiving you, they will have their pet Nolband back, and their fee into the bargain.—ED.]

PART III.

PUTTING INTO CONDITION.

Stabling and Clothing.

Having made your purchase from the stables, whether as a Racer, Charger, or Hunter, the same system is to be pursued in order to put him into strong healthy condition, externally and internally; and, from the day he comes out of the Bomb Proof, never let him enter a shut-up stable, but picket him in an open pendal, or under a tent. You are not in England now, and you will reap the benefit of this by-and-by. From about the 1st November to the 1st of March, one unlined cumly is necessary at night, when picketed in either of the above; but never more than a dungaree one, if he is, where he has no business to be, in a close shut-up stable. There is no medium with some people; they either advocate shutting their horses up in a dark and hot stable, thereby doing their best to introduce half a dozen English diseases; or else picket the poor brutes out in the open mydan the whole year long, caring nothing about the thermometer for six months being at 130 degrees in the sun,

whose rays penetrate their very brains; while, at other seasons, the rains pelt down on their unlucky backs unceasingly for twelve hours, and sometimes for twenty-four, without their being exercised during this time; thereby producing fevers, rheumatisms, bad surfeits, and inveterate mange. The man who pays fifteen hundred rupees for his nag, generally adopts the first of these, the hot and dark stable, by way of taking care of him; the man who pays five hundred, the open mydan, by way of allowing the poor brute to take care of himself; either way impairing their constitutions, and making them miserable.* A stable in all the warm latitudes of India should be nothing more than an open thatched or tiled pendal, fifteen feet high, and made into loose stalls of twelve feet square each. During the hot and rainy months also if the ground is not damp, make his bed outside; the insects in some places may be more numerous, but it is cooler. If at a cold station, and one side of the stable closed up, let it be the north-east side, and open to the south-west. Should you be anxious to ascertain whether it is the proper temperature, and clean or foul, sleep in it yourself for a couple of nights; and if you find it close and uncomfortable, so will your horse.

* The Bombay Cavalry are always picketed outside, and this is called a *state of nature*.

A small jhool, merely covering the back, and not even meeting under the belly, is but half a jhool. Jhools, whether to keep the horse warm, or to keep the rain, the sun, or the flies off, should all be made on the same principle. A fourth-inch broad band should tie or buckle across the chest, to prevent it slipping backward; the hinder part should cover the thighs nearly down to the hocks, a large hole being cut for the tail; this in the cold weather prevents the wind from rattling along the belly; but, when the jhool is made of thick numbda, it is better to have the hinder part of lighter stuff. In the hot weather, when made of dungaree, this form alone keeps the flies off the inside of the thighs and testicles, and it never interferes with the motion of the legs. Whether made of heavy or light material, it should double well under the belly, and have two ties of inch and a half broad tape, one piece a little behind the withers, and one in the middle of the back, both fastening on the near side, just under the ribs. The breastpiece must be separate, or the horse can never have the free use of his legs; and it should always be worn with the body jhool. An extra piece, a foot square, should also be put on the back, when a roller is worn. The neck and head piece in one, when required, must fasten with tape over the withers on to the body-jhool, but not too tight, or it may tear from off the head when lowered to lie down or to eat.

Fly-flappers should always be worn from the 1st of March to the 1st of November.

When you use a tobra, let the mouth be the smallest part and not the bottom, or a lot of the grain is lost on the ground by the horse throwing up his head.

The Feet, Physic, etc.

Having picketed your horse where he can breathe fresh air, and at the same time be safe from a stroke of the sun, send for the Nolband, take off his horrible shoes, have the soles properly pared, the horn rasped smooth, and let him be walked on the sands unshod. Give a little bran mash in each of his feeds of ground gram; as much water as he likes to drink; but not much grass. Continue this for a week or ten days and then prepare him for physic. This is the only time I shall urge you, without a very good reason, to give physic; but on first coming out of the stables (if he has not been physicked since he came out of the boat, which you should inquire for many of the dealers are great physickers) he will be safer; and, if thin, get into condition quicker by a mild dose.* Prepare him,

* J. Stewart, p. 48. "Hundreds of horses have been ruined or destroyed by the first journey they have made after being purchased. The buyer has been ignorant, that, to command even moderate work from a horse, he must be prepared for it very gradually, and by a systematic course of treatment."

Horses never thrive well in Bombay during the rains. If purchased from the stables in April or May, they should be physicked and sent off to the Deccan by the 1st of June: the benefit will be great, *if properly* looked after.

therefore, as directed, and give over night a drachm of calomel, and at daybreak the following morning a drench, deducting a drachm of aloes for the drachm of calomel.* If your medicine is not made up at Mr.

A short time since I heard the following dialogue between two of my brother amateurs, one a civilian, the other a militaire.

Civilian.—Are you fond of physicking horses?

Militaire.—No, I never yet physicked a horse, and never would. I never knew any good come from physic.

Civilian.—Well, I differ from you. I never had a horse come into my possession, I did not immediately physic; and never would. If I had fifty, I would physic them all to-morrow.

Now, there are some individuals in almost every camp holding exactly these opinions: they never give a thought, nor care a button, whether the horse requires physic, or not: they are persons who have never seen a stable properly managed; have most likely never had a valuable horse, and most certainly never ought to have one. One always physics, because it is his custom to do so: the other never physics, because it is his custom not to do so.

But worse than the above are some of those fresh-landed sportsmen, who, with pockets well lined, have, previous to embarking, obtained a slight insight into their grandfathers' or uncles' great studs, and therefore they would have us believe they must intuitively have great knowledge of horse-flesh, like the strolling boys that played the jews-harp, saying, they must be good musicians for their fathers belonged to the town band. These people commit more cruelty, havoc, and ruin in one year than their careless or less wealthy brethren do in a dozen; and I never knew one that could harden the flesh and get a horse a good wind, and at the same time preserve the legs clean, and the heels open.

* See "PHYSICKING." Never venture on more than one dose in order to put your nag into condition. If that does not effect all that is required for this purpose, when no ailing exists, twenty will not; there is something else needed besides purgative physic. Eight drachms of purified bazaar aloes, such as you may get and purify yourself, I find equal to about six of those procured from Mr. Treacher or Mr. Sprague. Twelve drachms of unpurified bazaar aloes I have given without effect: they are very apt to bring on gripes, bloody urine, and inflammation of the bowels.

Treacher's or Mr. Sprague's, at least learn that the aloes is good; as that brought from the bazaar contains from a third to one half of dirt. Never attempt to allow for this by giving a larger quantity, or you may be much deceived, since two drachms too much may irreparably injure the bowels of a weak horse, perhaps send him out of this world. Two days after the physic has set, if he should have been purchased thin, hide-bound, having the lampas, a staring coat, and looking dull, as these are all merely the symptoms of a deranged inside, take ten drachms of emetic tartar, five drachms of ginger, and five ounces of nitre; mix these up with honey or ghoor, divide into ten balls, and give one every morning at daybreak. Continue a little bran mash at each feed with his ground gram, and let the gram be steeped in hot water ten minutes to soften it, or change the food to boiled oorud, which is an excellent fattening grain on these occasions; or give some boiled barley at each feed with his bran and soaked gram.* There is no necessity to muzzle him every night you give these balls, unless he is a foul feeder and eats his bed, and then it should never be omitted until his stomach regains its natural tone. When they are finished, put a drachm of black salt, finely pounded, in his gram at each feed,

* In boiling or steeping grain, never put more water than it will soak up, or a large portion of the strength of the grain goes into the water that is thrown away.

and in a few days double it, and then treble it; nothing being more palatable, or better suited to recall the appetite of a horse recovering from sickness than this. Green grass, or lucern, but never with the dew on it, (it should be dried twenty minutes in the sun after cutting,) should also be given in moderate quantities, if procurable; if not, sliced carrots; and he should be walked, or ridden at a walk, morning and evening, according to his strength. It is want of care during the first month with these thin horses; giving them eight seers of hard grain when they cannot digest half; putting them into a canter when they are scarce fitted to be mounted for a walk; omitting physic, or giving it too strong; and allowing them to continue gorging on all day and night at dry grass, that disarranges their inside, and renders it so difficult to bring them round again into any thing like order. If you have attended to the foregoing, and evident improvement has not taken place in a month, but he still looks dull, and feeds badly, and you can discover no rheumatism, no fever, no worms, (examine well the pulse, mouth, teeth, &c.) persist strictly in the above way of feeding, and give every morning at daybreak, for a fortnight, some of the beer-tonic, p. 141. Let him, also, always now drink some warm gram gruel, sweetened with a lump of ghoor, when he comes in from his walk in the morning, and a little also on return from his walk in the evening. This simple

treatment will benefit him greatly ; and when the ribs are well covered, the flesh on the quarter, and the belly of moderate size, he may be called in condition externally—a sort of dealers' condition, having an appearance ready for sale, though, of course, not ready to be violently galloped—one, however, whch undoubtedly is far preferable when making a purchase, to taking a skeleton, inasmuch as some few horses really do exist that never can be made fat, and you would be so far certain you had not got hold of one of these ; besides, it is half the way to real condition, and is a state you should always advance your horse to before you attempt to put him in condition internally ; for, by commencing work too quickly before you have ascertained whether there is any latent ailing about him requiring a different course to be pursued, you may be thrown back three or four months.

Hours of Feeding and Watering.

The usual feeding hours in India are morning, noon, and evening ; and as that is a good system by which horses get even quantities at regular hours, and go out with their stomachs empty, (riding times being morning and evening,) it cannot be advantageously altered : for noon, however, substitute one. The hours of nine, one, and seven are a better division than nine twelve, and seven ; and do not allow him to eat grass at nine and one for more than an hour and a half afte

each feed, when it should be invariably all swept away. The use of salt for preserving health ; two or three drachms in each feed is granted, and should therefore continually be given. If the black salt is used, a smaller quantity will suffice.

Water is usually given twice a-day ; first, after morning's feed, and again between three and four o'clock. They will often thrive very well on this, and if offered it at other times, even in the hot weather, will frequently refuse to sip any ; but a horse that is out of condition, or one that is to be trained after he is got into condition, will be better if water is allowed oftener. Give a dhool, about three gallons, half an hour before the morning's feed, and another half an hour after it ; a dhool at half past twelve, half an hour before the one o'clock feed ; and another at half past three. In the evening, when he comes in from walking, and is cool, offer half a dhool more, but do not let him drink more than this at any one time. Horses that are only watered twice a day will often drink a bhæstees large dhool twice filled, six gallons each time ; some will drink more, which, with their bellies full of gram and grass, I have seen produce most violent gripes.* Never allow the dhool to be

* Watering immediately after feeding on barley is said too often bring on gripes and inflammation : this is far truer of wheat. A too sudden change of food ; from gram to barley, and then watering immediately after, may bring on gripes and inflammation ; but then the sudden change of food is as

used for any other purpose than drinking, and then the older it is the better.

Grooming, Washing, etc.

A good gora-walla, well looked after, will clean your horse quite as well as an English groom; and twice in the day, at eight in the morning and half-past three, with a rub down and a slight brushing when he comes in from exercise in the evening, are the proper times. During the months of March and October, or when he is changing his coat, dispense with the curry-comb, if thin skinned, and use only the hand-rubber and brush. Moulting, in all hot latitudes, is very slight, yet some horses are a little weak in consequence: a bran mash should be added to the gram every evening if this is the case, and a little sweet gram-flour gruel given in the morning. The hair of the mane and tail is often greatly disfigured by the curry-comb and hand-rubber, making it all scraggy, and giving it the ugly and dirty appearance of mange. Never permit the curry-comb or hand-rubber to go within three inches of either the mane or the tail, and then the hairs of the mane from the near side, that hang over to the off side, will be

much the cause of the disease as the watering after. Watering immediately before or immediately after any grain is unwholesome, and to some horses dangerous, and after gram quite as much as after barley. Horses of delicate stomachs, indeed all horses, should be watered half an hour after morning's feeding, to prevent the food swelling in the stomach too much.

equally long with the rest; and those nearest the root of the tail will be also long: the hair destroyed in one day by a slovenly mode of cleaning, will take a whole twelve-month to grow again to a proper length. The comb is often in fault too, from the numerous teeth in it: have it of thick, strong bone, but file away every other tooth, not leaving more than six, including the corner ones; and then the hair will neither be broken nor pulled out. The mane and tail being in nice order, add much to the appearance, and often materially facilitate the sale of a horse. The manes and tails of all horses, excepting the cavalry, are combed to the off side.

When brought home in the morning, after his walk or canter, at whatever season of the year it may be, never allow him to be taken into the pendal till he is quite cool. The wisping with grass, the instant the saddle is taken off, must then be liberally performed; first over the saddle-place, till every hair is dry, and then all down the legs to the hoofs, until the fetlocks feel warm from the rubbing; this will often prevent swelling of the legs: but, if the legs are begun with, the saddle must not be removed; or, the back being neglected, the first minute it is taken off may give cold, or produce warbles. At this grass-rubbing, it is always better to have a couple of gora-wallas employed, if at hand, and which is imperatively necessary if the horse has been in the rain; but, that

over, he may be left as safe till the gora-walla is ready to begin with the curry-comb. During this cleaning leave the legs from the knee downwards alone, till he is watered; or, if you like, till half an hour after that, when the grain is given; then lay on the hand-rubbers, and afterwards brush all smooth. Wash his feet and heels lastly, if they must be washed; but, unless they are dirty, it is quite unnecessary. That abominable English custom of washing a horse's legs when he first comes in, and that often in cold water, tends to produce windgalls, cracks in the heels, and contraction. In the cold weather, it is true, it is generally done with warm water; but the rubbing and scrubbing to dry the heels, and then the draught of cold wind they are exposed to in the stable, is apt to produce cracks as well. In India, once every other day, when he is finished cleaning, at four o'clock in the afternoon, is quite often enough to wash his legs: the chill should be taken off the water in the cold weather.

Should you be lately from England, you perhaps have as great an aversion to wash your horse's body as you have a predilection for washing his legs. Some grooms have, I know, a great dislike, even in the height of summer (the only time it is asked,) to wash a blood-horse all over: you think it looks strange and fancy it spoils his coat. The poor post-horse gets the advantage here, only that he frequently goes in

hot. Do not be prejudiced: you are now in India, and, for three or four of the hot months in the year, your horse, especially if he is a high-caste one, should be taken to the river three times a week, about noon, before he has his one-o'clock feed, and washed all over: he enjoys a lie-down in the water; it refreshes and invigorates him greatly. In May, and the very hot months, if he perspires about twelve o'clock, half an hour before sunset should be chosen, and let him have his walk afterwards. Always take care that he does not remain in above ten minutes, and that he is rubbed down dry, and moved off in as short a time. Independent of this, the mane and tail should be washed once a week in lukewarm water and soap, (trotting well afterwards in the cold or rainy weather:) it keeps the scurf away, and makes the hair grow. The inside of the ears is a part so little thought of, that large scabs sometimes form there from the sticking of clots of dirt. Every morning and afternoon, when the cleaning is finished, these should always be nicely wiped out with a wet towel, taking care that no water drops into the ear.

Getting out of Order.

If you have purchased your horse fat, in good external condition, and during the time he is being put into internal condition, by stronger exercise, &c., he should unexpectedly get stale, and cease to improve

if his skin should become somewhat hide-bound; his coat lie rough; small surfeit-bumps arise on his body; his mouth have the lampas, &c.; should he, in short, get into that lubberly state, like a "ship in irons,' evidently wrong, and neither go backwards nor forwards, you have been keeping him in a hot, shut-up stable, or, you have not duly attended to all the directions here laid down. Whatever has been the cause, treat according to "PUTTING INTO CONDITION," commencing with the mild drench of aloes and salts, followed by a week of the beer-tonic, p. 141, if rather fleshy, three or four of the ALTERATIVES, p. 97, should succeed to the physic, previous to giving the beer-tonic; but never think of bleeding; no, not if he is as fat as a hog.

Gora-wallas Riding.

Some persons have great objection to gora-wallas riding. If you have any kind of a decent gora-walla, and not above seven and a half stone weight, or eight and a half for a strong horse, he will, by one month's instruction, do as well to ride, at a walk, as if you had Jem Robinson. Give an old saddle, and see that it is placed well backward, clear of the shoulder, and kept there by being lined with plush; or, if necessary, by a crupper; and take care that the stuffing is quite free of the back bone, so that looking under the pommel you can see daylight right through, or the back may be galled, and you will accuse your gora-

walla of your own negligence, and be crying out, "this comes of gora-walla riding!" Let him be ridden also with a small light mouthing-bit, and make the gora-walla keep the head well up, not by pulling at the bridle, but by pressing the heels to his sides, and occasionally giving him a touch behind with the bagdoor, one end of which is to be fastened to the ring of the mouthing-bit, the other end being in his hand, to guard against the horse running away if by accident he should fall off. This will preserve his mouth, prevent his ambling, make him put out his fore-legs, and go light in hand, and be proper walking exercise; while the gora-walla will not be too tired to clean him well when he comes in. There is not half so much danger of your horse falling down when the gora-walla is on his back, riding in this way, as when he is slouching by his side, allowing the head every minute to come within a foot of the ground. A horse should only be led (except in training) when he is too sick, or too weak, or too tired to be ridden; and then, if it is the cold weather, the jhool in the morning's walk should be kept on; but walking exercise at other seasons of the year, with clothing on, when the thermometer is ninety, or more, is enough to put any horse "in irons."

Danger of Overfeeding.

Condition is not the work of a week; it will often take half a year for a horse that is thin, or fresh

from the stables, and if you attempt to fatten too quickly by grain, before the strength will admit of your giving proportionate exercise, the stomach, unable to digest a large quantity, will be weakened; thus the grain will often pass whole, and the horse fall off instead of improving, or he will grow dull, heavy, and gross. And even should he be able to digest all you give, this raising of the condition too quickly is very apt to produce fever, or inflammation of the bowels. It is always dangerous to allow an idle horse to overfeed on grain, it brings on scouring; besides, flesh gained at this risk adds but little to strength. There are some horses also that will keep plump on eight seers, and yet not thrive on twelve.

Continual increasing and decreasing of grain is likewise very bad. Feeding a thin horse on ten seers a day, then reducing it suddenly to half the quantity, and then increasing it again to ten—all this irregularity is destructive to getting into condition; yet I have seen a man, when training, act not very differently from this, but I never heard of one winning, unless it was a donkey race.

Mixed boiled Food, Chaff, etc.

Mixed boiled food is often absolutely necessary to recover lost flesh; and, when given only once a day, the evening is the proper time. One mäp of coltee, one of oorud, one of barley, one of bran, half a one

of linseed-meal, with three drachms of salt, or six ounces of ghoor, according to the horse's taste, all put into hot water, and boiled for twenty minutes, with one mäp of dry ground gram, and one of sliced carrots or turnips afterwards added, make an excellent evening's feed, and one which is very refreshing after a long day's hunt.* This kind of supper, however, is often refused, unless the horse has been a little habituated to it, or if the cooking-pot it was boiled in was the least greasy or dirty, or perhaps from dislike to one of the articles. Any change of diet, any thing new, must always be introduced by little and little, or the sensitive and dainty appetite of a horse will be sure to take disgust. Potatoes, or yams, should only be steamed, and turnips only parboiled; but these had better not be given in large quantities raw. Carrots

* Coltee, oorud, moong, and methee are the grains for boiling, and sometimes barley also. Coltee requires more boiling than either of the other three, but none of these should ever be given raw. Bajree and mhut are not so good for boiling; they are better in their natural state, and ground gram also, unless particularly hard, or the horse sick. Bajree and mhut are always mixed, as bajree alone is too heating; and mhut alone gives the gripes, especially when it is mixed with rats' dung.

Wheat is the most nutritious of all grain, but at the same time the most liable to disagree. The strongest-stomached horse could not bear a change to an entire wheat diet under four or five months, and then it is very unwholesome. Barley is the next most nutritious grain, then oats, then peas, and lastly, beans. It appears singular that beans, which in England are deemed absolutely requisite to a horse at hard work (not to train on), and which are known to put on the hardest flesh, should contain less nutriments than the other grain; and it appears equally singular that lucern, the value of which we all know in India in fattening a horse, should be the least nutritious of all the grasses; but so it is.

and beet may be given with impunity, and very wholesome they are; begin with a little. Turnips, carrots, and linseed-meal, when continued for any length of time, improve the skin and coat considerably. In the hot weather, two or three of the large radishes, called moolee, may be also given daily, stalks and all; but the natives term them cold and injurious, if water is drunk immediately after them. A seer of methee, boiled, and given at each feed, in lieu of one of the seers of gram, is also very good in some cases; for this grain has a double advantage, being very nourishing, and its adhesive quality preventing its being swallowed too quickly. Giving large quantities of ghee, ghoor, and trash of this kind, is disapproved of; and very properly so: but to a horse displaying all his ribs, a seer of bajree flour, six ounces of ghoor, and a wine-glass of ghee, baked up together into a large thick ap, with half a drachm of ginger and two drachms of anise-seed, and given after the morning's feed and the same again after the evening's feed, if he can only be induced to eat it willingly, will rapidly help to cover them. Ghee is often very old and rancid; so, if disrelished, try it without the ghee, or with nice fresh butter. Sugar-cane, sliced, is also most nutritious and wholesome; but too many things are as bad as too few; you must select those that best suit his taste. Chaff will sometimes be found very useful. A pound of dry grass, a pound of barley or wheat-straw, boosa,

and four pounds of lucern, chopped up small, and given after each feed; or his feed of boiled grain, bran mash, sliced carrots, &c., may all be mixed in the trough with half the quantity of this chaff; and the other half given afterwards.

After a horse has picked up in flesh, and is capable of being ridden for morning's and evening's exercise, he must not be allowed to get grass. A horse, after four years old, should never be too fat, any more than he should be too thin. Feeding also too much or too long on boiled food, chaff, sugar, &c., is bad; it must always be discontinued as improvement takes place, for it makes the flesh too soft, and may possibly have something to do with engendering worms; and the same benefit will not be derived from it as if given only during the times of thinness or sickness.

A horse that has been long thin and out of condition, (if the emaciation is not caused solely by starvation or overwork,) always needs a *mild* dose of physic, previous to being fed on boiled mixed food. The "ALTERATIVES," p. 97, are generally best adapted to this case, provided the owner will only see his horse properly mashed, to prevent the chance of over-purging, and thereby doing more harm than good.*

* Caveat Emptor, p. 19. "I fed him for a month on chopped clover, bran, and malt, fermented by a little yeast. This is the way to pickle a horse for a friend!"

The above kind of feeding, continued incessantly for eight or nine weeks

Native remedies for improving, fattening, and giving a horse an appetite.

FOR IMPROVING.

The Ralib, called Hulwaee Udruk, to be given only in the cold season.

Take of ghee, pounded turmeric, green ginger, and methee, two seers and a half each; put the ghee on the fire first, fry the turmeric well in it; then throw in the methee and green ginger, and let them fry thoroughly: then mix five seers of brown sugar in that, and add ten seers of milk. Work the whole up into a confection, and give a quarter of a seer once a day after watering. Increase it by degrees to a seer. It is a perfect elixir and, if given throughout the cold season, the horse will be a world of beauty

FOR FATTENING.

Give turmeric, for six weeks, that has been steeped for twenty-four hours thus: mix one-eighth of a seer of turmeric with milk in which it has soaked for twenty-four hours, and give it once a day, fasting. Increase the quantity gradually to a quarter of a seer.

The seer for the above, means the native puckah-seer, which is nearly two pounds. The cutcha seer is nearly a one-pound measure.

FOR GIVING AN APPETITE.

Take equal parts of mustard, bhang, salt, and kichree, (kichree is rice and dal,) with flour just sufficient to mix the whole in. Weigh, and give about ten drachms once a day, after he has finished his feed, keeping him reined up for a quarter of an hour afterwards. This is also good when the bowels are flatulent, or when at exercise they make a rumbling noise, like a mussuk of water in the inside.

All these are useful at times, provided they are given in moderation, and nothing whatever ails the horse, but only simply wanting flesh, or appetite. If taken willingly, of course the better.

lies on the fat in an extraordinary manner. The belly does not become large; on the contrary, there will generally be a nice, roundlooking carcase, from the great nutriment contained in the food, and its easiness of digestion; but the whole barrel, as well as the neck, become thickly laden with gross fat. In England, where the generality of horses are geldings, and the neck consequently thin, the bellies being also often of large size, a little of this benefits the appearance; but to Arabs, who are almost all entire, and whose necks, with the exception of the high caste, are seldom over-light, too much feeding on such fattening stuff ruins, instead of improving, the appearance.

The system to be adopted with regard to exercise must be regular; and it is to be gradually and never suddenly increased. Having now improved and become strong, he should have a canter during his morning's exercise three or four times a week, and any green meat that he has been feeding on changed to dry grass; but three weeks must be allowed before the change is wholly completed. Increase one mäp of ground grain every fortnight, and let the exercise be proportioned to this gradual increase of grain, and not the grain to the exercise: he is now only being put in condition, so, if he gets thin, decrease the exercise; but do not increase the grain too quickly. Some horses will get fat and keep in condition on very little grain; so much the better. By the time his appetite and digestion can contend with ten, eleven, twelve, or thirteen seers of ground grain *per diem*, (thirteen seers of ground grain is about ten pounds,) according as he may be a large or small horse, and a large or small feeder, he ought to be in strong healthy condition, externally and internally, which is to be thus defined: when he is fine in coat, and that fineness has been gained by the free use of hand-rubbers and brushes, and not by warm jhools, or hot stables; when he is firm in flesh, the flesh well up on the quarter, and he carries a good carcase; neither tucked up under the flank, nor let down like a cow: when he is high in spirit, produced by kind-

ness, and regularity of feeding: when he is fresh on his legs, and they are as clean and unblemished as on the day that he was foaled: and when all this has been brought about while he has been kept in an open pendal, so that with one extra head and body covering put over the single blanket, he could go on a march, or be hunted, and sleep in the open mydan, in the cold weather, without showing or feeling the slightest bad effect from it. This is the acme of perfection of condition.

How to keep Condition.

As continual gentle exercise is the surest way to preserve health and prevent disease, so never omit, when unable to ride yourself, to let the gora-walla mount for an hour's walk, both morning and evening, giving some of the grain boiled every other night; this, when a horse is idle, or only at ordinary riding, being the chief part of the secret in keeping that plump appearance over the quarter: but, if even for upwards of a month he has nothing but walking exercise, reduce the grain, if he is fat, and feed partly on green food; and be cautious that the change from any soft feeding, to dry feeding, and from common exercise to hard work, is always gradual, or the digestive organs will be weakened, and the legs will swell. When a horse is suddenly put to work, after being fattened on boiled food, lucern, &c., three

moderate gallops will often take off all the flesh he has gained in as many months, making him ill besides.

Dry hard grain and dry grass are as injurious to a horse's body, when standing for weeks without exercise, as boiled food and green grass are to his legs whilst hunting. Turning out to grass, as they do in England, to eat nothing but green grass, or laying up altogether, in India, to eat nothing but boiled grain and green food, has long ago been proved destructive to condition. We have no opportunity in this country of doing the former; and the latter should never be resorted to, unless sickness or great poverty demands it. Too large a quantity of bran, such as bran mashes in every feed, is also very improper, and very lowering, if continued even for a fortnight only, to a horse in health. Bran mash is the diet of the sick, or lame horse, or occasionally to give at night mixed with the grain, when the dung is in small hard balls; but the dung in general is far too soft in India.

Never muzzle at night, unless you have some reason for it, such as preparing for physic; the horse being a bed-eater, &c.; and in the latter case something should always be given in the middle of the night, or an hour before daybreak. The system of fasting from nine at night till nine in the morning, to please the fancy of the owner, is ruinous to a horse's inside.

Never allow of stimulants or massallahs indiscrimi-

nately; and particularly avoid them whenever the least symptom of illness, or falling off in condition appears, for they are destructive at the commencement of many illnesses, although most salutary in other stages; therefore, give them only when out in cold or rainy weather, or during hard work; on which occasions the pint of Hodgson, sweetened with an ounce of ghoor, adding half a drachm of ginger, finely grated, and one drachm of anise-seed, whilst on the fire warming, is the finest cordial of any.

Never fail to have the feet pared once in every twenty-five days, if they grow quickly, or thirty days at most; and the first day you observe the least contracting of the heels, take off the shoes and treat for "CONTRACTION."

Never picket your horse with two tent-pegs six feet apart, unless he is a weaver, or you are about to shoot him; two close together are far more secure, and he is less liable to hurt himself: and if you must use heelropes, let them be very loose at night, or many an hour's sleep will be lost from fear to lie down. I have seen many horses thrown completely out of spirits, and some out of condition, solely from this cause.

Look after your horse yourself; make much of him always on mounting and on dismounting; and *see*

that he is made comfortable in his stall: by this means a bad tempered horse will grow fond of you.*

Change from Stabling to the Open Air, Hunting, etc.

As walled-in stables are now universal in every cantonment, and as little probability exists that you will alter those attached to your habitation to that form of structure most congenial to a horse's health, you may easily suppose, that when brought from one of these warm stalls to sleep in the open air in the cold weather, and no precautions taken, illness of some kind, as a matter of course, will follow. A warm bed-blanket, head and body piece, wrapping well under the belly, should always be put under the ordinary jhool on these occasions, and the horse kept close under the tent-walls, out of the wind: this, with half a massallah at night, and taking care the gora-walla mounts at day-break, or before, (but without taking off more than the upper jhool,) for a good hour's walk, will be found the best preventive against those colds and coughs, which bring on a staring coat, debility, loss of flesh, and general bad condition.

If unexpectedly taken out during the monsoon, and rain should fall during the night, change the jhool,

* Lawrence, vol. i, p. 279. "The tempers of horses, like those of their masters, are various, endowed with a greater or less proportion of intelligence, sagacity, and feeling; and it is but too often the beast evinces the greater degree of rationality.

give a small massallah, and trot him for quarter of an hour *immediately* the shower is over. A single light black Deccan jhool, *unlined*, (the best description of jhool there is next to the bed-blanket,) should always at nine o'clock during the monsoon be put on all horses, unaccustomed to the open air for though it may be close and hot, and no rain fall, yet it is generally damp. Keeping thick jhools over the body during the heat of the day to guard against the scorching sun is very pernicious, but a light dungaree one, padded half an inch thick with cotton, along the spine, is proper, and should always be worn when out in the sun, in the hot or rainy season. The shade of a tree, however, should be searched for: it is cooler and more agreeable to a horse than even stable or pendal; and hog-hunters should not forget this, for the continual exposure to the fiery sun throughout the day damps the spirits of many horses considerably. No one, but a native, can keep his hand upon a horse's back for ten minutes between the hours of twelve and three in those months, when picketed in the sun, without burning or blistering it: you may imagine, therefore, the effect on a high caste, thin-skinned horse, and tormented by the flies to boot.

Hunting a horse for two or three months on nothing but green food, only cut the day before for his forage, —not at all an uncommon occurrence,—or even half

green and half dry, is as much the cause of those rounded shank-bones and gummy ancles, as being taken out of an idle stable, and suddenly put into training exercise, without gradual inurement. Hunters are often fed too much and too long on green food; racers may also be included. Considering that a few days' green meat is often wanted to recruit the strength after a severe cold or other illness, no horse kept solely for racing or hunting should have more than ten weeks' regular soiling throughout the year, and that at two different periods. As training usually commences in October, the month of April, or early part of May, and that of June, or early part of July, will be judiciously chosen as the most proper season, during which time the grain should be decreased nearly one-half, and the exercise never exceed a trot. The hunter can have his in August, September, and October, when the grass is long and rich, and the boar allowed to rest; very little being given during the months he is hunted. Green grass at coming in, during the first fall of the rains, is very weak and washy; but lucern, kept half-buried in water, is equally good throughout the year.

To retain a horse, then, in hunting or hardworking condition, he must be kept chiefly on hard food: he should have a three-mile gentle canter every other day, and always be exercised mounted, every morn-

ing and evening. If wanted after breakfast, allow but little water after finishing his morning's grain and less grass, so that he may go out with both stomach and bowels tolerably empty. Feed and water in moderation (a gallon of water and a seer of ground grain, with a mouthful of grass, will be moderation,) if you halt to take your own tiffin; give first a small cordial drink, if the least fatigue is apparent; and, even whilst hunting, always allow two or three go-downs of water at any river or tank, if he thirst for it; this little will not hinder his galloping, but will prevent faintness. Invariably walk the last mile to cantonment, that he may return home cool, but never feed until he has been in an hour, and well cleaned above: then *see* that he has a cordial drink, massallah ball, or some gram flour gruel; likewise that boiled food according to his liking is made, to be given *after** he has eaten about two mäps of dry ground grain; that a small dhool of water is offered half an hour before the dry grain is put into the trough, and another half an hour afterwards: *see* also that his legs are well malished while eating it, and that a good bedding is placed under him, in a loose stall. These little attentions after severe work are a horse's rights: he

* The soft food at this late hour will produce a quick and wholesome distention of stomach, which will induce sleep, and prevent his standing two or three unnecessary hours to produce it by eating dry grass, which, at this late hour of night, is as bad for a tired as for a weak-stomached horse.

is sagacious enough to comprehend and appreciate them, and will amply repay you—if your ugly conscience does not—by cheerfulness of going the next time you require his services. When the fetlocks and feet are so dirty as to require washing, let it be done in warm (not too hot) water, and *see* that the legs are well dried and malished afterwards. If he refuses his food, it is most likely from your neglect of him after dismounting, and not the distance, unless you completely galloped him off his wind and brought him home hot: in either case, foment the legs in warm water, make a large soft bed, let him drink if inclined, and then muzzle for two hours, when he will probably have regained his appetite.*

As to the considerate Christian, who, after a hog-hunt, or long day's journey, will carelessly hand his

* There is one time that massallahs, or cordial drenches, prove excessively hurtful. When severe distress is occasioned by a long run, and you find that your horse can with difficulty go forward, give a drench on the spot, if you have it, loosen the girths, and lead gently home; but by the time he reaches his stall, he will be feverish, and cordials or stimulants then may do serious harm. A seer of steeped ground grain in a good bran mash, sweetened by a lump of ghoor, a drachm of emetic tartar, and two of nitre being put in it, will, in this case, with a large bed and loose cool stall, be the quickest and safest restorative, repeating the same on the following morning. A clyster of salt and water must also be given on arrival at his stall.

When a horse drops exhausted, bleeding three or four quarts is recommended, giving also a pint of wine or beer, to enable him to crawl home after an hour or two of rest. When home, treat as above, with bran mash, emetic tartar, &c.

faithful steed over to an ignorant syce, perhaps to be tied up and left in the not unfrequent position of a bullock at the slaughter, the hind-legs being tugged afoot too far backwards, and the head pulled down to the tent-peg, and he march off to fill his own greedy stomach—may nought but hot beer and tough meat welcome his palate for dinner, and the night-mare, in a buggy bed, be his only nocturnal repose!

"THE HORSE TO HIS MASTER.

Take care of me a mile out, and a mile in.
Up the hill spur me not,
Down the hill push me not,
On the plain spare me not,
In the stable forget me not."

Massallah Balls.

Caraway-seed	3 drs.	Mustard	½ dr.
Cardamom-seeds	2 „	Cinnamon	½ „
Ginger	1 dr.	Black pepper	1 „
Turmeric	2 drs.	Anise-seed	2 drs.

The whole to be finely pounded, and baked up with gram flour and ghoor, into a thick ap. A little ghee or fresh butter can be added, if the horse is fond of it. Some of the light pleasant stimulants will frequently be eaten with a relish, if the horse is accustomed to bread and they are given fresh.

Cordial Drinks.

Any one of the following, given in half a pint or a pint of warm beer :—

Brandy, Gin, or Rum ... 2 wine-glasses.
Hot water ..... 3 „
or
Port or sherry ..... 4 wine-glasses.
Hot water ..... 2 „
mulled with nutmeg, &c., just as you like it yourself.*

Beer Tonic.

Gentian	**2 drs.**	**Ginger**	**¾ dr.**
Anise-seed	**2 „**	**Black pepper**	**½ „**
Turmeric	**1½ dr.**		**in a pint of beer.**

One-half of each article, in only half a pint of beer, is the proper strength to begin with after illness, increasing to the full dose in a week.

For a horse whose dung always remains soft after the grain has been reduced, and there is no apparent reason to account for it;—but not to be given until he has first had a mild warm drench of physic, and been afterwards fed for a fortnight on partly boiled grain, with a little gruel always added to his water,—this will frequently harden the dung into balls of itself; and then the following ball or drench, given every other morning, when gradually brought on again to dry grain and dry grass, will help to keep it so.

Ball.

Gentian, quassia or chreeate	**1½ dr.**	**Ginger**	**½ dr.**
Catechu	**½ „**	**Bhang**	**$\frac{1}{15}$ „**
		Opium drench	**$\frac{1}{15}$ „**

* It is a bad practice accustoming a horse to cordial balls or cordial drinks. It is like a man accustoming himself to drachm-drinking after a time he cannot do without them. I would never give them but when a horse is in a state of exhaustion from overwork, or when requiring cordials and tonics when recovering from sickness.—ED.

this, mixed in half a pint of beer, or in one wine-glass of brandy, and three of hot water. These relaxed constitutioned horses should, every other evening, have their grain boiled, and not be fed wholly on gram.

A pill, the size of a pea, is often given by natives in cases of fatigue, or in gripes. The pill is thus concocted, but it requires a native to make it, who has been taught how to prepare it, and then it will keep for years:—Six drachms of solid arsenic, and six quarts of onion or moolee juice are put in a pot, and placed on the fire for eight hours, till all the juice is absorbed. About one drachm of the arsenic is lost in the residue, and the remainder is made into sixty pills; each pill, therefore, contains about five grains of arsenic.

The following is also given by them when a horse is fatigued, and necessitated to continue his journey on the following morning:—

Opium	½ dr.
Bhang	1 „
Alum	1 „
Googul	2 drs.
Turmeric	4 „
Ghoor	4 „

made into a ball with bajree flour.

Native Massallah Ball.

Ginger	½ seer.	
Mustard	⅛ „	
Black pepper	¼ „	
Asafœtida	6 drs.	
Garlic	½ seer.	
Kala jeera	½ „	cummin seed, or like it.
Kucha methee	½ „	fenugreek like it.
Umbee huldee	¼ „	a turmeric-coloured zedoary.
Gora waj	¼ „	also called butch; the orrisroot, or like it.
Kootke	¼ „	a febrifuge, a small round stick.
Underjow	⅛ „	a small long seed, like the oleander.
Bungrekan	¼ „	a white, mineral-looking substance.
Gulloo setwah	¼ „	also called iskung, the root of a tree; colour, yellow.
Lindee peepah	⅛ „	a long pepper.
Jerkatolah	⅛ „	a kind of flat bean.
Palas papra	⅛ „	a very thin flat bean, crimson outside, pale yellow in.
Ajowan	⅛ „	a very small seed, half the size of a caraway.

This is enough for twenty balls.* These ingredients should not be all made up at once, but kept pounded, a little being mixed up fresh with ghoor, when wanted; then, it will often be eaten, if the gorawalla keeps his fingers playing in the angles of the mouth: or, as there is nothing bitter or nauseous in it, a small quantity may be baked up in a large ap. If forced down, the balls must be soft, in the shape

* These massallah balls I would never give. If a horse requires any thing of the kind, the cordial ball or cordial drinks at pages 140 and 141 are much better.—Ed.

of an egg, but not more than two inches in length, nor more than three quarters of an inch in diameter. When a balling-iron is used, it should always be covered with leather, and the tongue very gently held.

TRAINING,

That is, for the turf, is the highest artificial state of existence the horse has to endure; and artificial means must be resorted to before any horse can possibly be brought on the course in a fit state to run a contested race with a chance of winning; but there is no necessity to follow in the wake of admitted errors, much less to practise absurdities, or to block up the animal's nostrils every night at nine o'clock with that confounded little muzzle: when the muzzle is required, let it always be of the shape and size before directed.

Near one-fourth of those that are put in training, are never brought to the post, and no wonder: the very stable management of many is radically bad; too much attention is paid to trifles, and the essentials, the very life of training, neglected. Some are over-physicked and over-fed, and then over-physicked again; the bowels, of course, get out of order; the stamina are weakened; and the legs, as a natural consequence, fail: others are over-clothed or over-sweated; and almost all, over-stabled. Shutting the stable-doors close in India can only have one advan-

tage, and that is, nobody can then be aware of the nonsense that goes forward within. How often does training commence on the slight, and perhaps delicately-constitutioned Arab, with two drastic doses of physic? Before two months are passed over, the bowels become greatly deranged, and the legs bunged: it is then declared, he (which ought to signify the trainer) "went all wrong;" and this wrong, or rather wronged horse, has to be physicked again, and laid up in bandages for a month, and perhaps blistered, before he can be got rid of. Sometimes a sturdy animal will stand a great deal of mismanagement, and yet appear at the post, and even win, if entered for a race that answers for him and running on a course suited to his make, &c., but this is not gaining by judicious training; much oftener, however, he fails to show his countenance, having been ruined by a regular adapted process, as the nasty dry coat, disordered inside, over-worked legs, and contracted feet most abundantly testify. A well-made horse, indifferently trained, may beat a bad-made horse well trained; and one of high caste, indifferently trained, may beat one of not so good caste well trained, &c., &c. It is through these, and various other distinctions, that bungling training is always hidden. Horses, no doubt, will continually go wrong when being fine-drawn in the last dangerous month; but, unless from accident, &c., it should not occur until

that time. It is useless your attempting to train, unless you punctually visit your stables at every feeding and watering hour, (have a good watch, for you should never be more than five minutes out in your time,) then carefully inspecting the grain, grass, water, &c., and always look in the last thing before you go to bed, and the first thing at daybreak in the morning; for, however good your gora-wallas or jockey may be, you should never lose sight of the proverb, that "He who works with his hands, has seldom too much in his head."

Training in India and training in England are certainly different,* but there is no difference in the

* The Arab, it has been justly said, cannot be trained in India as the English horse is in England: but the remark holds good reversed; the English racer could not be trained in England as the Arab is in India. The difference of country, climate, food, and the manner in which the two, from infancy are reared, require an alteration of system, and render much of the artificial means pursued in England unnecessary; but the English race-horse, notwithstanding his wonderful performances, will, perhaps, not stand more training, more actual galloping, before he comes to the post than a high-caste Arab; and yet an English thorough-bred will easily outstrip the best Arab, in either pace or distance, and beat him in daily journeys, or continuance of labour; though, for these two last, the Arab might prove victorious, if on his own native soil.

Arab blood is a little in disrepute just now, owing to those that have been exported from this country having been of a very mediocre caste; but I do not believe that a thorough-bred, genuine, unblemished Arab horse, and certainly not a mare, of the proper build and stamp, would be objected to—the one for a stallion, and the other for a brood-mare; or either to be crossed with suitable-sized English thorough-breds—by any of the great racing breeders at home: and if twenty genuine, high-caste Arab horses

principles; and those consist in condensing the greatest quantity of pure muscle into the smallest possible bulk, and so gradually to raise the muscular powers and wind to that degree of perfection that every atom of speed may be drawn out without any distress being exhibited, and without the slightest damage to the legs. In effecting this consists the whole arcanum of good training; and, by attending to the following, you may accomplish your end, and be able to cope with your neighbours; but if you enter on your sporting career, intending to train all horses in the same manner, without any regard to their peculiar constitutions and temperaments, their likings and dislikings, thinking the whole secret consists in making

and mares were now to be sent to our cold climate, and as judiciously crossed, and well taken care of, for the next fifty years, as the race now in existence have been from sire to son during the last half-century, they might equal in size, and surpass in speed, all those of the present day.

When last in England, my opinion was asked of a late importation from the Bomb Proof, price twelve hundred rupees, and if I thought he was a *real* Arab. I replied, to my confiding querist, I thought he was a real Arab, quite as much a real Arab as a cathammed horse he had purchased of his baker a few days before was a real English horse: and I recommended that he should embark that to his munificent donor in India in return, who would then have a real English horse, and a most equitable exchange, too. An Arab, to be worth acceptance at home, must either be very showy and handsome, answering for a lady's park horse, or else have proved his blood by the very best performance, displaying besides the cardinal points of a good stallion; for the very best performance of an Arab, in India, would be very third-rate at New-market. A few of the genuine caste are annually imported to Bombay, but three parts to seven-eights bred are what chiefly fall to our lot.

them thin, you will generally succeed, to your heart's content, in bringing down all the flesh and fat of the body, depositing it, unluckily, about the legs and ancles : when degraded to this state, they should be shut up in the dark stable ; for such horses, " there is no place like home."

Stable and Clothing.

If fresh air and a cool stall are necessary to put a horse into condition, how much more so must they now be, when his powers are about to be exerted to the highest pitch they are capable of attaining ; but, as the more inducement to rest, and to lie down, a horse in training has, the better, close up the sides with tatties that he may not be disturbed during either the day or night. One single blanket jhool, made as described at p. 113 with the single neck and head-piece, will generally, during all November, be quite sufficient clothing ; and these are only to be worn at night, and during the morning's walk. In December and January, if it should be cold at night, the thermometer below fifty-five degrees, another body blanket can be added at nine o'clock. It is sudden changes from warm clothing to no clothing, and from warm stabling to cold air, that hurt a horse, not a cool and uniform temperature. The thermometer, at noon, in the stall, during the cold months, varies from seventy to eighty degrees.

Stable management.

Fresh air, without a thorough draught; keeping the stall cool, not cold; and the horse warm, not hot; good grooming; proper clothing; daily airing of the jhools; cleanness of litter; strict attention to the food and water, and punctuality in giving them; constant regard to the appetite; careful observance of the dung and urine; never disturbing while at rest; stopping the feet every day and night; shoeing once in every twenty-five days; physicking only when actually necessary; sweating only when actually required; and enforcing the mildest of treatment,* but at the same time never teasing with kindness, constitute the essence of stable management.

What description of horse are you about to train? Is he a slight, or a narrow-chested, or a flat-sided, or a delicate, or a hot fiery horse; or the reverse of all this? A powerful-limbed, fine-chested, well circular-barrelled, strong, quiet-dispositioned horse: and is he young or old? for, according as he varies between all these, so must your training vary. Has he been only a short time out of the stables, and just advancing into something of that state we left him in at p. 131, under "EXERCISE," or was he half-trained last year, and been kept in tolerably strong exercise, with

* Gora-wallas are seldom known to wrong a horse; though covering themselves, by mistake, on a cold night with his jhool, and pilfering a little of the evening's feed, are no very uncommon occurrences.

proportionate hard food ever since? The latter has a decided advantage; and if he has been down the gulph two years, and half-trained each year, he has of course a greater advantage still. Racing colts in England, it is true, come to the post at three and four years old, in fine order, most of the great stakes being for young horses; but a couple of years' extra hard food, with as much exercise as can be given, without rounding the shank-bone or bringing a wind-gall, adds no little to strength and vigour.*

Time required to Train.

A horse in good hands is deemed capable of proving his utmost speed in five, or, if taken from grass and no exercise, in seven months: many for the Bombay turf, fresh from the Bomb Proof, are brought to the post in three months; but so short a period is not sufficient, unless they have been in strong and regular exercise for some time previous. Others, again, which have been laid up to fatten and get a decent external appearance, are in this state (only half way to condition) put into training exercise, in the hope of bringing out their full powers in as brief a space, by dint of sweating and galloping:† these, like the former, are almost sure to get thrown over

* In India a little more soiling is thought necessary than in England; and, if so, then a less quantity of physic is needed.

† See the note under "SWEATING."

in some way or other, for should their legs prove of that solid texture to stand this hasty training, it is more than their bodies will—they "fly to pieces." The sudden high feeding on grain is also apt to make them foul, or they grow stale, and, if there has been any taint in the pedigree, perhaps sulky into the bargain ;* besides, a three-part bred Arab, though of the most willing temper, will never stand the training of the genuine blood. The legs, likewise, sometimes swell during the day at the commencement of training : this arises often from debility, and more from the horse not being ready for a good canter at all, than from the exercise having been too severe for this stage. Call, therefore, the first two to six months, according to the time it may have taken you to putting into condition ; when you have accomplished that, three to four more will be ample to carry him up to the mark.

A day's Routine.

Every morning, precisely at daybreak, your trustworthy horsekeeper (if he does not merit that appel-

* A good caste Arab will sometimes sulk too ; the General, of 1839 and 40, for instance ; but then the cause, whatever it may be, seldom arises from strong training. The misfortune is, they generally take it into their heads to sulk the day of the race, by which one would suppose the concourse of spectators intimidated them, or perhaps reminded them that this is the day of whip and spur. Gelding well sometimes cure sulkiness, though there should be nothing wrong with the testicles ; and I should be inclined to risk that operation to backing an uncertain horse.

lation you had better change him, or sleep in the stable yourself,) is to look quietly in at the stall, to discover whether the horse is lying down; if so, let no noise be made; there he is to remain till he voluntarily rises: if standing up, put on the snaffle bridle immediately, for it is just at this time, after the night's rest, that he will commence nibbling at grass, one blade of which is hurtful, now that he is going out to exercise. First, with a perfectly dry towel, wipe the ointment, given at p. 192, out of the heels, or the dust and dirt may adhere, and the ointment become the cause, instead of preventing cracks. One gora-walla having been placed on each side, take off the night-clothes, and brush him down well for the space of five minutes; then replace one body jhool and head-piece, &c., or your English set, if you have them, putting the saddle above. The jockey or gora-walla is then to mount, ride him to the course, and walk him about, not at a snail's pace, nor at an amble, but in the manner described at p. 124, and at the utmost extent of his walk, for an hour and a half. The clothes are now to be taken off, the saddle placed well back, (the work properly of less than a minute,) and the canter given; after which he is to be walked till he is cool, then brought into the stall, and the head tied up *instanter*, one end of each snaffle rein being buckled forward to the wall, and the mouthingbit in his mouth; the heel shackles

(with heel ropes attached, if he is disposed to dance from side to side,) having been also fastened, the muzzle put on, and one gora-walla, as at daybreak, placed ready on each side, take off the saddle, and wisp him down briskly with dry grass till every hair is smoothed, and the skin is becoming warm again under the belly as well as the body; that finished, do the same down all the legs and fetlocks, till warmth succeeds to the rubbing. The hands are to be used next, to get off as much hair as possible; the syces must dip them in water, and heave strongly against his sides, for after a month all the loose coating ought to be off. One man will now do, if you cannot spare two, to go on with the cleaning. Commence first with the curry-comb, taking care it is not too sharp, and that it is used very gently if the skin is at all fine or tender, the curry-comb being only to raise the dust: the hand-rubber second; and that can scarcely be laid on too strongly for some horses; but if the skin and coat are fine, and the horse restless, that must be gentle too. The hair-brush well applied thrice to the horse and once to the curry-comb, not thrice to the curry-comb and once to the horse, is to follow thirdly; a dry cloth to wipe away all the loose hairs, fourthly; and then throw a light blanket jhool over the body, to be taken off about half-past ten, if the weather is warm, after he has finished his grain and grass. Now offer him

the small dhool of water; take the chill off if it strikes you as cold, or he hesitates to drink it without, which is often the case in December and January, in the middle of the day as well as in the morning, especially if drawn from a well; then proceed to the legs, commencing with the hand-rubber, and finishing with the brush. Lastly, pick the feet out, and wash them in warm (not too hot) soap and water; but above all things, be particularly attentive to drying the heels quickly, not scrubbing them too strongly, with a coarse towel: when dried, put on the ointment. The eyes and face may be washed in the warm water too, with a large sponge, this being as comfortable to your horse as to yourself. The grain is now to be given: half an hour afterwards a dhool of water, and then four or five poolies, nine or ten pounds, of sweet dry grass.* Half a bedding of litter is also to be thrown down, (but this may be omitted during the first month, unless an inclination to lie down in the day is observed,) always putting some of the old well trodden, that has been dried in the sun, uppermost; for this answers three most useful purposes: it prevents any long blades of the fresh bedding running into the eyes. The sight of two horses I have seen almost destroyed from this, the thick stem having nearly cut through the cornea: it is the softest, and

* This is by far too much. Not more than three or four pounds at the outside ought to be given at any one time.—ED.

therefore disposes him to lie down; and not being fresh, he will abstain from picking at it when he has finished his allowance of grass. Take away the remnant of the poolies at half-past ten, and leave him undisturbed in a loose state till half-past twelve. Water is then to be offered, the coat to be brushed, and the legs to be malished and shampooed. At one, the mid-day feed, with two or three poolies of grass, and he is to be left again to himself unapproached till half-past three or four. The stall at this hour is to be swept clean, water given, and the afternoon grooming to take place: the feet and heels are not to be washed again at this hour, but the morning's ointment is to be wiped out. About five o'clock put a light single set of clothes on, (during October and the early part of November he may go without any at some stations,) and have him led for an hour, though many, if strong or fleshy, will do much better ridden at this time also; then brought into the stall, well rubbed down with the hand-rubbers and hair brush, his feet picked, a little water offered, and the grain given. The bedding is now to be made, a little of the ointment put in the heels, the clothing changed for warmer, if it becomes chilly at this time, and as much grass placed in one corner as you intend he should eat during the night. At half-past eight give a small feed of grain, varying from half a mäp up to two mäps, adjust the clothing, and

leave him for the night, as he ought to have been during the day, in a loose state, unmuzzled, unhead-stalled, and unshackled. Thus you have one day's routine, which is to be adhered to with but little variation so long as he is training.

About one morning in every seven should be a holiday, and Sunday will be a fitting one. Let the walk take place at daybreak, as usual, the saddle being above the clothing; and whilst out give free scope to his play, and allow him to gape, yawn, move about, and halt at his pleasure. The neck is neither now, nor at any other time, to be kept constrained with a martingale, which impedes the free action of the forelegs, and causes tripping. This annoying rein is sometimes resorted to to keep the horse's head in its proper place, and steady; but oftener because the rider is unable to keep his own in its proper place, and steady. Unless to a bolter, or a ewe-necked determined stargazer, a martingale should never be used. English young colts, half-broken, may require different kinds of reins and martingales, but a decent rider can dispense with them all upon an Arab.

Training Weight, Physic, Exercise, etc.

You should always train with as near the weight as possible that he is to carry in the race; never more than an extra half a stone, and that is half a stone too much for a slight horse. A strong-legged, straight-

spined, high-caste animal may not prove the worse for it, but a slighter horse will have his stride shortened an inch in the course of three months, and an inch in every stride will make two or three lengths at the end of a race.* You ought also to have some idea as to how much flesh your horse can go to training with, so as not to have to sweat off a whole lot, to the detriment of his health, and at the expense of his legs. This is difficult to tell, perhaps, with a new purchase, but a strong-built, good constitutioned horse, will naturally throw up flesh quicker than a slight carcased or irritable one; and a young horse, (if not quite a young colt,) up to seven or so, will throw it up quicker

* There are very few Arabs in India whose owner's weigh above ten and a half stone, that have not their action in some degree hurt; and a light weight, who is conscious of the nimble active step of a perfectly fresh nag, would detect at once a kind of shortened step, or slight deficiency of spring. By a perfectly fresh nag, I mean a horse that has been taken care of so long as he had a colt's tooth in his mouth, and never had more than eight stone on his back up to five years old. I have mounted numerous "beautiful light going horses" for sale, the property of owners of about eleven stone; in lieu, however, of fine airy action, they hammered away, in comparison with fresh nags, like so many stone paviers. No man, of course, can be expected to acknowledge this until he has sold his "finest going horse he ever saw." Being a light weight, I would always willingly give three hundred rupees extra for a valuable horse, if he had never been mounted.

A horse frequently pulls up lame from some very trifling wrench or strain, or interfering knock. Halt on the very spot for half an hour, and apply cold vinegar, (a bottle of which should always be taken to the course,) then lead gently home: two days after, you will often find him able to take his canter again, if no subsequent swelling ensues. Further advanced in training, over a hard course, tenderness comes over the feet: poultice for two nights, and put a piece of narrow wax cloth, trebled, between the shoe and the crust: it will remain firm if the crust is rasped level for the shoe.

than after that age; but whatever description of horse you are about to run, he is to be put well in condition before the training commences; and if between the age of five and eight, with a large carcase, it is not desirable he should be so plump as one that is rather young or rather old, slight constitutioned, or flighty; for, these large carcased strong horses, especially just at that age, of five, six, or seven, throw up flesh so quickly that their legs would be in danger, from the great work and sweating required, to draw them out fine. You must endeavour, therefore, at the end of condition (if your object is the turf) to have more or less flesh on him, according to circumstances.

Presuming, then, that he has been enjoying the cool pendal, devoid of heavy jhools; that his feet are well open at the heels; that he is to escape the antiquated and prejudicial drastic purgative balls, now that he is in good condition, and that you have full three months before the races, commence with one mild drench of physic—the high feeding, and strong exercise, about to be pursued, render this indispensably requisite, as a safeguard against the numerous little ailings that are so apt to arise and cause an overthrow. In accordance, therefore, with the universally adopted precept, that "no horse can run without physic, give the aloes and Epsom salts, &c.;*

* See "PHYSICKING."

and if he belongs to any of the first enumerated class, under "DIFFERENCE OF FORM," he should not purge more than six or seven times, and he must have nothing but walking exercise for the first three days after the physic has set. That over, proceed to the daily canters; and though he has been in strong exercise before, commence with only a mile and a half: in a week increase it to two miles; in a fortnight to three, and then quicken the canter, with due regard to the state of his body, &c., never urging him beyond a hand-gallop during the first month; and during the whole time of training he is never to be pressed to his full speed, excepting at the trials; but the pace is always to go on so gradually increasing, without any irregularity in the distance, that his powers may be arriving at their highest pitch by the commencement of the last fortnight. If you are training for a long race of three miles, the morning's canter should be nearly four, in order to get "the length" well into him. If for heats of two miles, the canter should be full four, or there is no objection to your giving it of only the length of the heat, walking a quarter of an hour, and then repeating the same. If for a short race, or short heats, the gallop and distance may be proportionably quicker and shorter, but the exercise is never to be so severe as to produce the least distress.

You may put it down as a rule that every Arab,

during the first two months of training, should be above his work; if there is an exception, it is where there are fine, broad, flat limbs, added to a well-formed barrel, and the horse a little sluggish or vicious, then an extra gallop may be taken out of him with impunity, and often with advantage. But the weak points must always be the guiding mark: for instance, when there is a fine round barrel and well-ribbed home, yet slight or rather tied-in legs, the latter must be the guiding mark as to the quantity of exercise; but when the carcase is too small, or flat-sided, with a long hollow flank,—making, perhaps, a rather washy horse—yet the limbs strong, the former must be the guiding mark.

Trials.

Towards the end of the second month it will be advisable to take a few trials, that you may be able to judge if he is likely to be qualified for the race you wish. Try a mile and a half with eight stone four pounds, and three days after, the same distance with ten stone, if he is to run for a welter. After an interval of another three days, try two miles and a half with eight stone four pounds, and after three days more, the same distance with the eight stone four pounds again. From this you will learn how he carries weight, and whether he is capable of running a long or short distance best. In all these trials push

him well, closing the heels to his sides if of the lazy tribe; but never punish unless he is a regular sluggard. Continue henceforward taking trials weekly, or every ten days. At one time start at score from the post, and keep him to his best pace the whole way: at another time bound off at a good gallop, but keep in hand the first half of the distance, then push him strongly the remainder, always having every half mile carefully timed: thus you will gain a wrinkle through what means he is enabled to exert his pipes to the most advantage, and consequently when to make play during the race.* In all trials, unless the horse is of the most willing temper, and flies by a pull of the ribands and pressure of the calf, you should have a fresh nag in waiting at the last half-mile post, to accompany him home; but if you own a superior known good nag in your stable, which is in training, it is better to let him start with the one in trial; there are so few that will exert themselves to their utmost without a competitor, and it is easy to tell the rider to hold in the first part, if you wish it. Previous to taking a trial, or giving a sweat, always put the muzzle on at two in the morning, and the day succeed-

* The stride of a well-built running Arab of fourteen hands two inches, at the top of his pace, on a good turf level course, carrying from eight stone to eight stone seven pounds, will be found upwards of fifteen feet; and if, when at this speed, the hind feet overstep the spot, the fore ones have quitted, he has superb springy, and undoubted running action.

ing a trial or sweat avoid putting him out, for fear of his legs.

The Grain, Grass, and Water.

Bajree and mhut is sometimes used at the commencement of training, but it is not so good as gram. If brought into condition upon it, give it twice a-day for the first fortnight; then once, so as gradually to discontinue it; and it may be crushed a little in the grinding-stone, the same as gram. Gram* and oats, equal parts, would be good, but the oats are seldom procurable. Barley is the only other grain, and a very excellent one it is, but it must not be suddenly introduced. At each feed of gram, give half a mäp of barley that has been put in the grinding-stone with the gram, and daily increase it, so as that at the end of two months, one-third barley, and two-thirds gram shall be eaten at each feed. The grain that is relished most will generally be most easily digested; but gram and barley mixed is the most nutritious and best; and with many horses you may go on increasing the barley to one-half. No grain must ever be given new; it should be from eight to fifteen months old, and not more. The grass, of course, should be the very best, the gingwa, having a fragrant smell, and stacked after the rains, though a little fresh-dried Huryoli, that has been cut a few days, or

* When gram is mentioned, it is always the Chenna or Bengal gram that is meant.—Ed.

a little fresh lucern, may be given at night with advantage; putting it down at nine o'clock, when he has finished his last feed. Gram should be heavy, sweet, and fresh smelling, without holes in it, and a sufficient quantity of it laid in for the four months, that no change may take place in the quality. When the gram, or barley, are very hard, give them an extra turn in the grinding-stone, but do not moisten the grain. If too greedy a feeder, and any should be voided whole, chop one mäp of Huryoli grass very small into it at each feed, and also add one mäp of bran mash,* for the rule is now, in a measure, to be reversed; the grain must be proportioned to the exercise, and, therefore, everything must be tried to make mastication and digestion go on well. He should love both his food and his work, then he will thrive. Three drachms of rock or two of black salt, in every feed, is useful in training to all kinds of horses: begin with less. You must be careful to distinguish the difference between a poor feeder and a little eater: the horse that eats the little set before him with an appetite, increases in stamina, and keeps

* If he has a natural dislike to bran, even this handful must not be given, or it may have an opposite effect, and induce him to swallow down the grain still more greedily, in order to get rid of the taste of the bran. This would not be the case, when gram or barley is given while putting into condition all boiled and mixed with the bran; then he would be obliged, more or less, to masticate the grain. A handful of linseed-meal, if not disliked, may be tried in each feed for this purpose, in lieu of the bran, and it is preferable.

his flesh on, is no weak-constitutioned horse, and you should not give to such a one more than he actually requires. The dung should always fall in balls at the commencement of training, and you should endeavour to keep it of the consistence of cow-dung the whole time; but the high feeding and strong galloping, and the nature of gram itself, renders it very difficult, and, towards the last month, almost impossible to do so. Boiled coltee, one feed a day, is good for horses that are naturally rather lax, as well as others; or it can be given for the one o'clock feed; and again at half-past eight o'clock feed; this, with the cordial ball, will generally keep the dung of a proper consistence, if there is no over-feeding. The dung, towards the later periods of training, is occasionally found, not only a great deal too soft, but slate-coloured, and so slimy and agglutinated, that when lifted up with a cane it will hang on either side without breaking; hence the horse grows stale: this proceeds from bad stable management in various ways, irregularity, or over-feeding, particularly on gram, over-sweating, or over-work. The cure is bran mashes for two whole days, with four drachms of gum arabic in each mash; and a little green food in lieu of the dry grass: then the drench of physic, p. 96; and this, followed by a few carrots, and the beer tonic, p. 141, daily: boiled sago should also be frequently given afterwards with the *last half* of every feed, and a little linseed

meal mixed in the first half, with the gram and barley.

The quality of the water requires as much care as the grain. A change to softer water may not hurt, but from soft to hard infallibly will, and that often too evidently. When it cannot be avoided, three gallons of hot water should be poured on three mäps of bran, and strained off when cold; or a quarter of a mäp of linseed, and a small lump of ghoor, may be boiled in a gallon of water, and then more cold water added; either will, in some degree, correct the change; or a little chalk and clay may be added to the water, letting it stand for an hour in the sun. As to training on actual hard water, you might as well try to train on bricks; a horse will not keep in condition upon it. River water should generally be chosen in preference to well water, as being softer, and ten or twelve degrees warmer; but clean well water, if not harder or colder, is quite as wholesome. During the cold weather, water from a deep well, in the early part of the day, is as warm as river water. The water in the cold weather is frequently in the morning too cold for a horse in training to drink; but by stirring your hand round in the dhool for two or three minutes, you may raise it to a proper warmth, without the addition of hot water, the smell and taste of which, to many horses, are very disagreeable. Placing a tub of water in the stall all day, so that he

may drink *ad libitum*, is a mistake, arising out of the discovery, that, when water is always within a horse's reach, and before his eyes, he will drink less than when only watered twice a day; but the object is not only that he should never over-swell himself with water, but also that it should be drunk at that time that most facilitates digestion. This tub-system, like the English one of taking to a pond to water, exercising gently a quarter of an hour, and then bringing back to the pond again, may answer very well for some horses, when putting into condition, but not for training.

Quantity of Food, and hours of Feeding.

The grain is always to be given in a trough, never in a tobra; and if ever he appears indifferent about it take it away, and re-examine it: perhaps you are giving too much, or it may be you fatigued him in the morning. Never stand coaxing him, therefore, either to eat or drink; for there are some horses that, by this means, may be induced to swallow against their inclination. See that both are clean and good, but if refused, take them away till next feeding or watering time.

The more grass that is put before a horse at those times, when it is not left long before him, the better, because he will have the advantage of picking the best out of a lot, and I suppose you do not require to

be told that his taste in this respect should have precedence of yours. After morning's feed—as it is all taken away again at half-past ten, and after evening's feed, as you can take away as much as you like, when you look in at nine—it is of no consequence how large a quantity you put down. The grass and water, as a rule, should be very gradually reduced, a little every month; and the grain as gradually (at the rate of about a quarter of a pound a week) increased; but the appetite for all must be consulted to a certain extent. If the digestion keeps pace with the appetite, and there is no looseness, well and good; do not overstrain his stomach in order to see how much he "*really would take in:*" if the appetite is better than the digestion, there must be reduction. The following (supposing he was in good condition when the training commenced) may be considered a fair quantity for a full-sized Arab, of fourteen hands two inches high, actually to swallow in one day and night.

	Grass. lbs. of dry Gingwa and Huryoli.	Water. gallons, five bottles to a gallon.	Grain. lbs. of gram and Barley, or partly boiled coltee.
First month, in one day and night.....	9	8	10
Second month, in ditto and ditto.....	8	7½	11½
Third month, in ditto and ditto.....	7	7	12½
Last fortnight.....	6	6½	13

Remarks.—Gingwa grass, though very fine, is not English hay, neither is gram, oats; or a little less of the former, and more of the latter, might probably be considered a fair quantity.

Regarding water, the less a horse in training *is inclined* to drink the better, appears an acknowledged axiom: at the same time he is always to have as much as his appetite leads him to take, but not more than three gallons at each watering time.

Thirteen pounds of gram and barley is about seventeen seers, mäps, or powder-canisters' full, when ground.

The racer must be fed often, yet never till he is hungry, is a maxim in training; but as much as five mäps of ground grain, about three and three-quarter pounds, may safely be eaten at each feed; and whatever more is required, to the extent of two mäps (not beyond), it would be better to give it at half-past eight, to changing the hour to nine, twelve, three, and seven; the former interferes less with the rest during the day, and the small feed at half-past eight or nine may diminish the wish for grass during the night. Much, however, must necessarily depend on the horse eating largely, and swallowing too quickly. If a large feeder, and quick swallower, and no grass be chopped with the grain, five feeds may be requisite, at nine, twelve, three, six, and nine; but the first,

of nine, one, six, and half-past eight, (the last one always partly consisting of boiled barley or sago,) though not generally adopted, will answer best when properly fed. If you are a determined advocate for the hours of nine, twelve, three, and seven, you should enter upon that system when you commence putting into condition. Feeding at daybreak, even to the amount of a kutcha seer—the custom with some turfmen, who always muzzle at night—is not a good practice, unless for such a voracious feeder as requires to be fed six times a-day; and then, after being muzzled all night, it may be necessary. An hour before would, even then, be better than daybreak. If an English training groom feeds early in the morning, the exercise does not follow so soon after as it does in India; but there are as many little variations in different stables at home as there are in this country.

Muzzle, when required, Leather Dhool, etc.

If at any time during training, notwithstanding the old bedding being uppermost, you find he eats some of it, either during the day or night, after you have allowed a fair quantity of good grass, the muzzle, annoying as it is to some horses, must be used, but not put on till after nine, and then something must always be given at or before daybreak. In this case, however, you probably commenced training before

he was properly in condition; for the appetite of a healthy horse is mostly moderate both for dry grass and water, and he can scarcely contract the habit of feeding too greedily on either the one or the other, while under this regular regimen of training. The circular barrelled, large-carcased, strong horse, is more liable to overeat on grass than the light carcased; the last description frequently will not eat enough. Sometimes a horse will eat half his allowance of evening's grass by ten o'clock, and not touch it again till just before daybreak, and then fill himself: this should be discovered, and prevented by the muzzle being put on at two or three o'clock, when he may not be lying down. Always give the water four times a-day, allowing the small dhool once full at each time, with another half-dhool, if wished for, when he comes home in the evening, as advised under PUTTING INTO CONDITION. He will be inclined to drink less, on the whole, in this way, and thrive the better. You cannot be too particular with your leather dhool: gora-wallas are always putting curry-combs, brushes, and old shoes, in a wooden tub; besides, a tub is not half so nice as an old dhool, tinged with moss-like green from the continual use of water. A new tub should have a few shavings burnt in it before being used, and if once taken to wash the feet, a delicate horse will smell it for a week.

Cordial Balls, Horse-bread, etc.

The belly is sometimes a trifle larger than is altogether compatible with training; the horse may not be at all too fat, occasionally the contrary, but the belly droops a little. Feed more liberally, both on grain and grass, if the bowels will stand it, and give a cordial every other morning; but even if with this rather drooping belly he should carry too much flesh, neither the grain, grass, nor water are to be stinted; (over-feeding, which is worse, you have been before warned against). Give, in this case, six of the alterative balls, p. 97, one every other evening on return from his walk, and sweat gently once a week. The belly will go up of itself, if you are not in a hurry.

If the dung becomes too lax, or you find him getting heavy and out of spirits, and neither caused by over-feeding; or if by accident he has been a little over-fatigued, lessen the exercise a day or two, and give for a couple of successive evenings, at the last half-past eight o'clock feed,

1 map of Barley, boiled for two hours,
1 Do. Sago,
2 maps of Bran mash, into which put
4 drachms of Gum arabic, and
2 of Rock Salt.

Give also the following cordial-ball at daybreak, and a little bajree flour gruel just before his first watering time in the morning.

Cordial Ball.

Columbo root	1 dr.	Turmeric	½ dr.
Gentian	1 „	Sulpher	½ „
Anise-seed	1 „	Bhang	1/12 „
Ginger	¾ „	Opium	1/12 „

all finely pounded, and mixed up with the inside of a couple of figs or dates, or boil the figs or dates, skins and all, in a little sugar and butter, then mix in the ingredients.

Cordial balls, two or three a week, or half a one daily, are also indispensable to many delicate horses the last month in training; and the above is as good a one as at this time can be made up; yet it is very unpleasant to have to ball a horse continually, however expert the horsekeeper may be in doing it. A drench is preferable.* Half, or three-quarters of a pint of warmed beer is the best vehicle for these ingredients, or wash the ball down with beer.

Horse-bread, though seldom made in India, is beneficial on many occasions during training. When the muzzle is put on at nine o'clock, and nothing given till feeding-time on return from exercise in the morning, there is a fast of nearly twelve hours. This is just six hours too long. Half a loaf of bread, given half an hour before daybreak, two hours before

* To this I dissent. As before stated it is always much more annoying to a horse to give a drench than a ball.—Ed.

the gallop, would be very serviceable to many horses, even when muzzled at ten or twelve o'clock.

Horse-bread.

Gram or wheat-flour... ¾ lb. Anise-seed..... 2 drs.
Ginger ..... ¾ dr. Ghoor ..... 1 oz.
baked up together into a large thick ap, can be given instead.

The following recipe will make good bread, which the horse will in time grow fond of, if habituated to it by very slow degrees, daily giving a small bit when hungry:—

Wheat-flour and Gram-flour..... 3 lbs. each.
Anise-seed ..... 2 ozs. finely powdered.
Ginger ..... 1 oz. „
Ghoor ..... 3 ozs.

Add the whites of a dozen eggs, well beaten together, and as much beer, well up, as will knead it. Bake in an oven into three loaves, and commence giving when one day old.

Bandaging.

Training the legs, it appears evident, from the fatness so generally observable about the lower parts, is fully as difficult as training the body; but if the latter has been properly physicked, yet not over-physicked, the feet kept properly short, and stopped, and the strong galloping not too hastily introduced, the legs may be brought out as clean and wiry, and the fetlocks as smooth and undented, as during the first canter.

Flannel bandages to the legs are undoubtedly of great benefit, inasmuch as cold applications strengthen the sinews, and keep the legs fine, but the wetting of the bandages every hour annoys the horse and disturbs his rest, and lying down is of greater benefit to the legs than bandages. Do not take it for granted that the less a horse lies down in the day the more he will at night: it is the nature of some horses to lie down easily, while others, though fatigued, cannot always be induced to do so, even with a dark, quiet stall, and nice bed. Neat flannel bandages, kept well wet with cold water, or a little nitre added, I am a great friend to while putting into condition; but, on the whole, it will be quite as well to omit bandaging in training, unless the state of the legs seem to require it: which is too often the case, certainly. If bandages are used, do not put them on again after the walk in the evening.

Dry flannel bandages are used, by those who praise them, during the whole time of training. They may be of benefit to some legs; and they have this advantage, that, if so wished, they may be kept on during the night; and the rest is not disturbed by wetting them during the day.* If worn, they must be put

* I am an advocate for dry flannel bandages. They promote the insensible perspiration, and keep the legs of a horse in hard work much finer than wet ones. They have the advantage also of a horse not being continually annoyed by wetting the bandages, and they can be kept on during the night.—ED.

on a trifle looser than the wet bandages; but I recommend you to abstain from these, unless you have some reason for their adoption. Whatever bandage you choose, be careful to lay the folds even, commencing immediately under the knee, and bringing each turn as high in front as behind, so that the *bottom of each turn* of the bandage shall be fully as tight as the top of each turn. Each turn is to descend exactly one-half of the bandage, yet in such a manner that each looks straight in front. If any part of the bandage should be tighter than another, it is that close about the fetlocks, and not immediately under the knee; an extra piece, of inch-broad flannel, should be tied in front round there to prevent it coming undone, and the same also round the pastern. Dungaree bandages should never enter into a good stable, and a dry dungaree bandage is more likely to cause than to cure a bad state of the legs.

Warm flannel bandages, used for one hour after a severe morning's exercise, (put on after cleaning,) hot water being thrown on them every five minutes, will supple the joints, and prevent swelling; or, if you prefer shorter work, foment the legs with hot water, as hot as you can bear your hand in, for ten minutes after he is cleaned above; or let the feet, one by one, stand in the large tub from four to five minutes, with the water *well above the fetlocks*, not forgetting to thoroughly dry each quickly, as it is taken out.

Sweating.

Sweating would not have to be so much practised if the flesh were only brought to a firm and solid texture before being put into training. A strong dose of physic, and two severe sweats, will, in a fortnight, metamorphose a large carcase into a tucked up, gaunt belly; but Mr. Green would never attempt such palpable mismanagement as this on a horse that was intended to run to win; and the legs must be of good timber indeed not to thicken under it. Horses, whose legs have small back sinews, or which are tied in, or are crooked, will, notwithstanding every care bestowed on them, often show an evident inclination, after a good sweat, to counteract Nature's errors in some degree, by throwing out a little more substance about the hocks and ankles: heavenly creatures like these were never intended for such wicked work as this. All heavy sweating is dangerous to slight legs; still, sweating must be had recourse to, if the horse is determined to get fat, when at the same time he is not overfed: it is an evil, but not so great as having to physic again. The use of sweating is to take off the carcase, or neck, any little extra flesh, not deemed necessary, and which the exercise fails to do; and if done at the proper time, and not overdone, the benefit is often very great; for, two of the grand desiderata are, to strengthen the legs and diminish the weight of the body: the continual regular exercise

accomplishes the first, and this will effect the last. A gentle weekly sweat, towards the conclusion of training, is, therefore, necessary for every horse carrying too much flesh; it finishes him off, and draws him out fine; but with many of the light, highspirited, overwilling ones, it may often be wholly advantageously omitted. If the neck is a little too fleshy, he should, from the commencement, have a couple of head jhools on at the morning's gallop; the under one without ears; but they must be thick indeed to produce any visible effect on this part: the neck will fine a little of itself as the flesh becomes firm. If by nature a large-bellied horse, and a little too fat when commencing work, he should always have one set of body-jhools round him at that time. Some horses, however, will carry more flesh about the carcase than others; and if it is only muscle well condensed, you perhaps have a trump, and too much sweating, or extra physic, to draw such a belly over fine, will often be only running to certain ruin. Loaded with three complete sets of jhools, a three-mile gallop, with a couple of half-mile spurts in it, is as long, and strong, and heavy a sweat, as the most gluttonous Arab would ever require, if not too rapidly brought forward. Some turfmen always give the exercising gallop in one jhool to every kind of horse: that may do very well in England, but is not so well suited for India; it is unnecessary to many, and does harm to others.

After a sweat, he must be quickly led home,—not to have other jhools heaped upon him, but all the lather immediately scraped off with nice bamboo hoops, not too sharp. The wisping must then instantaneously succeed, or there is great fear of his taking cold. When cleaned, and the jhool put on, give the cordial, if you think it required, and then mix half a wine-glass of ghee, not oil, with half a wine-glass of brandy, and, immediately the legs are dried after the washing in the hot water, rub it well in, about the hocks and ankles particularly. If the shanks should become sore, rub in the lotion, for tender shanks, p. 193.

A lot of frothy, greasy, dirty, scurfy stuff, occasionally comes from horses with any superfluous flesh that have not been sufficiently long in training; and some also from others during their first sweat; so, if on the second or third sweat he sweats quickly and with much of this, it is proof he is in a foggy state, and it must be repeated again in a week or ten days, giving four or five of the alterative balls, p. 97 one every other evening, as mentioned before under "Cordial Balls," &c., if rather too fleshy; and if not too fleshy, one drachm of tartarized antimony, and a quarter of a drachm of ginger instead; but if he proves difficult to sweat, (which he generally will by the third sweating-time, if in condition before the training was commenced, and you are not

sweating too hastily,) and that sweat is like water, and he dries quickly after scraping, he is getting into prime trim, and will go on well without any more.*

* To train a horse suddenly up to the best of his mark the time will admit of, that is not in condition, but all soft flesh, needs a man that has seen a good deal, reflected a good deal, and has great confidence in his own knowledge; for, as mentioned under, "TIME REQUIRED TO TRAIN," p. 150, he is so very liable to grow stale, or go off his feed, or to injure his legs. After the mild dose of physic, sweating must here be commenced, as well as concluded with, if at all too fleshy, so as to get some of the useless fat out of both the inside and outside, before taking the stronger gallops. When brought home after a morning's sweat, extra clothes, contrary to what is stated above, are to be put over him for a quarter of an hour or more, until he sweats freely in the stall,—in his fleshy case it is necessary, for it acts as a protection against illness;—and when scraped down and dried, a fresh single set of blanket jhools is to be immediately put on, a small cordial given, and also half a gallon of luke-warm water offered: ten minutes after, he is to be taken out to have another quarter or half mile canter and then brought in for the regular grooming to proceed. The time for commencing these sweats, the quantity of clothes that he ought to sweat in, the pace that he is to go at, and the length of the sweats, are all difficult affairs to manage properly, for the flesh cannot be taken off too suddenly even in the stall, and the untrained legs will not stand it being taken off in the gallop, on the whole, I would not risk the chance of injury to a favorite horse by this hasty training however good the stakes might be. If only two months are to elapse before he starts, a gentle sweat every week for the first three weeks, and then a couple of stronger ones the last fortnight, adapting the length, pace, &c., to the size and grossness of the horse, will be near about the proper time, observing that the first sweat should not take place until he can prove his wind,—until he can "blow his nose." After the first two or three morning canters, if he stands panting at the sides, and fails to throw the mucus from his nose, by that peculiar quivering snorting shake of the nostrils, a few days more should elapse before the first sweat is given. Grooms who pay attention to this work, I believe, say the nostrils should always be breathed in three quarters of a minute from the time he is pulled up; and that some consequence is attached to these minutiae, I know, for I well remember a groom anxiously looking out for the blow of a fine strong horse, that had been sick for a

The same kind of remark is applicable here as under "SWEATING." If in good condition, and no soft flesh, when put into training, the one does at the commencement will generally be all that is required. The shŏk that some people have for physicking a month or so before the races, not because the horse is, but because they are afraid of his becoming too plethoric, is the cause of many overthrows and breaks-down. Whatever arguments you may hear to the contrary, this is not the time for preventive physic. The giving daily laxatives also at this stage, and that without a sufficiency of bran mashes, is not only equally censurable, but dangerous and bad, fatal gripes often coming on just when the physic might have been expected to operate.* A craving, large barrelled, deep-ribbed, strong horse, and that has to run a long race, may sometimes need a couple of doses of purgative physic, besides sweats, during a four months' training, in order to prevent him growing stale in his body, or round in his legs; but a slight, hot, irritable, or weakly one, is not to have every fire pulled to

month, when finishing his first morning's canter, and though it did not occur for a minute or more, it at last came with a fine healthy quaver—a good sign, the air passages are clear—and he remarked, "If the tripes only improve, there is no fear of his bellows." A slight horse, when of delicate constitution, will, of course, not have the loud shake in his nostrils of a strong healthy one: and if the horse was thin when commencing this hasty training, the only sweatings must be the last fortnight.

* It is incredible what extraordinary things some persons do in training which had I not witnessed, I should not have thought of mentioning them.

pieces by drastic physic, even if he does throw up flesh too quickly ; three of the alterative balls, p. 97, one every other day, with a couple of gentle sweats, a week between each, will suit much better ; and he will then come to the post in better order, and run under greater advantage than if more purgatives or laxatives had been administered ; besides, it is two to one against physic at this time having the exact effect that is wished with a nervous flighty horse ; it very commonly disagrees, or it operates too strongly ; in either case more or less debility ensues, and a month is not sufficient to restore the strength. Should heats have to be run, a pound too much flesh, recollect, is far better, for a light clear-winded horse, than a pound too little. The last description of horses are generally easily purged, and, in lieu of physic, oftener need gruel with four or five cordial balls, or cordial drenches, every week : they should have gentle training, and not be put to trial until within five weeks of running, and then never overtasked.

If a slight sprain, blow, or other injury occurs ; or the space between the shank-bone and suspensory ligament, and between that and the back sinews, should " fill," and there is no time to be lost, a drench of physic may be admissible, as the only means of cooling the leg and preventing loose flesh being thrown up ; and the horse afterwards very gradually

brought about by flannel, or *linen* bandages, *single*,* being kept on during the day, and wetted every half hour with the sal ammoniac lotion, p. 193. The greatest difficulty, however, always attends the bringing any "go-wrongs" up again, even in the best hands; and if the sprain has been at all severe, it is useless, as well as cruel, to attempt it, for you run the risk of ruining the leg for life.

As a general rule, then, no purgative physic should ever be administered within six weeks of running. When exceptions do occur, however short time you may have to spare, never be persuaded to ram it down the throat without some little preparation, for if gripes, in training, supervene from this cause, inflammation is generally close at hand. Bran mashes should at least be given for one day previous at all the feeds, and not more than two pounds of grass allowed until the muzzle is put on at nine o'clock; but unless a sprain or hurt has taken place, and the horse is about to be laid up, more than one day cannot well be spared for bran mashes without his going somewhat back in condition, for one full day must also be allowed after the physic has set, so as gradually to increase to the former diet, as any sudden introduction of dry grain into the bowels immediately

* In bandaging for a strain, let the bandage begin at the bottom of the pastern, close to the hoof, and go spirally upwards, instead of downwards.

the purging ceases is always hurtful, and frequently dangerous. At daybreak on the following morning, water first, the chill being off, and then give the drench, p. 96, to which you may had three drachms of Turkey rhubarb,* but as it is of the greatest consequence that a horse in training should purge, and not be merely sickened with the physic and thus so many days be lost, if a spoonful even is spilled in the giving, another should be added, and the clyster there recommended never omitted : wash the mouth cleanly afterwards with the salt and water, and if no strain has been the cause of the physicking, lead for an hour's walk. Read "PHYSICKING" in Part II, for the other precautionary treatment. If gripes takes place,

Warm beer	**1 qt.**	**Anise-seed**	**6 drs.**
Ginger	**3 drs.**	**Ghoor**	**3 ozs.**

may be first tried, not omitting the other directions, under "SICKNESS UNDER PHYSIC :" or the following ball, fresh made and soft, to be washed down with a little beer :—

Balsam of capivi.....	**½ oz.**	**Camphor**	**1 dr. dissolved in ½ a wine-glass of gin.**
Anise-seed.....	**4 drs.**	**Anise-seed oil** ...	**1 dr.**

the yolk of an egg ; mix.

* Rhubarb is found to possess so little purgative action in the horse, that it is never given by Professional men. There is nothing equal to aloes and nothing better than the aloes ball, when the aloes are good and properly prepared.—ED.

Should the physic operate a little stronger than desired, take

Chreeate ..... **2 ozs.** | **Anise-seed**..... **2 ozs.**

boil in three quarts of thin rice congee, strain off, and add to the three quarts of congee one ounce of finely-pounded gum arabic, and a bottle of port wine: give a pint, cool, every two hours, allowing some wheat flour gruel also, but no water.

Conclusion.

As horses in India are not plated on the morning of the race, shoe three or four days before; you will then see that the shoes fit well; and always take a couple of extra shoes, with nails and hammer, to the course with you.

The day before running, never give too strong a gallop: a walk will be sufficient, if the slightest disposition to flag was evinced on the previous morning's exercise; but it is most advisable your training should have been so managed that he should be in that state of trim and freshness to be able to take, and to require, a moderate canter; but more should not be given.

When stripped for the race, the flesh on the quarters should be as solid as a camel's; and it should be well on between the last false rib and the haunch-bone. The carcase should be straight, the belly looking nicely drawn up, and every muscle fully

developed. The coat should be glossy; the hocks lean; the legs and ankles smooth, and cool: and this is all compatible with the foregoing directions, which will perfect the wind, and carry him through, if he can win, as well as if you had attempted a more mysterious and intricate system.

If you have to start late in the morning, muzzle at nine, or twelve o'clock, according as you know your horse, the same as if you were in the first race; but, when muzzled early, or starting late, a third of a loaf of bread should be given a little before daybreak, about two hours and a half before the running; some give half a mäp, or a mäp of ground grain; the bread, or ap, is preferable. When the race is run in the afternoon, as at Bombay, still you should muzzle the night before, or during the night; giving the bread, and a walk in the morning; only return early, so as to feed at half past eight, instead of nine o'clock; and grant his usual allowance of grain, and also an hour's eating of grass; but not more than a gallon and a half of water at each watering-time: muzzle again at half-past nine, while he takes his rest till twelve. At this hour, if eager for water, allow four quarts; brush him well, and shampoo his legs, and at half-past twelve give a seer of ground grain, and also the third of a loaf of the bread; or nearly double the quantity, if he is not to run till after five; muzzling again till three o'clock, when the grooming is to take

place ; and let the jockey that is to ride, give him a hundred-yard canter a quarter of an hour before starting ; it will stretch his legs, and do him good.

Between heats, wine or spirits are given, with benefit. Four wine-glasses (thirteen wine-glasses to a bottle) of sherry or port ; or a wine-glass and a half of brandy, mixed with two wine-glasses and a half of water, or the same quantity of gin and water, are the proper allowance. Choose from these four according to your fancy ; but take care to give it fifteen minutes before starting. There is always half an hour between heats, so he will have a quarter of an hour's breathing-time to be rubbed down in, before swallowing it. Give it two or three minutes earlier in preference to two or three minutes later : this is of consequence, in order to have the desired effect ; but do not half stifle him with "blue ruin" the instant the heat is over, or he will be flying under, instead of over, the turf.

If, after a hard-contested race your horse's powers should have been over-strained and his strength exhausted, rub him quickly dry, put the jhool on, and lead him under a tree, or to a cool, shady spot ; then give four drachms of carbonate of ammonia, powdered, and made into a ball with water and linseed-meal ; after which, handrub the legs well, give a little water to drink, and a cold bran mash. Lastly, lead him

into a cool, open stable, and leave him in a loose stall, with a large bed under him. Look in an hour or so afterwards, and if the distress should still appear to be great, the breathing quick, the flank tucked up, and the eye red, take away from three to four quarts of blood, and leave him again to himself for twelve hours.

The racer cannot be kept in that excited state at the top of his condition for any length of time : " At the top of condition, on the brink of disease." And if he has to run at other races five or six weeks afterwards, the exercise and food should be moderately lowered for a fortnight, and a few carrots, a little green meat, and evening's boiled food given ; but whether he has been trained up to his full mark, or not, it does not invariably follow he is to be physicked again ; that must depend on the state of the body, and when he received the last dose.

If two mouths are to elapse, the drench may be given, varying in strength according to circumstances, and followed by refreshing for a few days with carrots, green meat, &c. ; but, if only half that time, it will be better avoided.

Coming out of training, never let your horse down all at once, but always give a gentle canter every other day, for the first fortnight, gradually reducing the grain ; and if after that time you deem physic

necessary, give the alterative drenches, p. 97, without any calomel at night.*

Riding; entering for a race; and requiting a querulous visitor.

Riding a race is as different from all other riding as a Scotch salmon is from all other fish; and as many aspirants for a silver cup, unable to afford racing establishments, both train and jockey their own steeds, let me advise you not to give up your own natural-formed seat for one you have never tried before. Do not stick your back up like an angry cat, striving to imitate William Buckle, or any other celebrated rider; but keep your middle person from the waist to the knees firm, yet flexible, grasping well with your thighs, and resting the weight mostly in the saddle, not in the stirrups; and if you have occasion to use the spur, apply it as far under the belly as the legs can reach. The management of the arms is even of greater consequence: you may have as

* On a visit to a racing-stable, at half-past ten at night, containing four horses in training I was introduced to the following stable-management. The first horse had heel-ropes on, and those so tight that they prevented him lying down. The second, a large and gross feeder, had the muzzle off if a little hard piece of leather with circumference scarce sufficient for a dog's mouth can be termed a muzzle—fortunate, therefore, it was off, for the animal would have been stifled by morning. The third was half-strangled, from the tightness of the neck strap of the head-stall. The fourth was smothering under two heavy jhools, and purging away from over-feeding; yet the old, careless griffin of a master had been successful the year before. Who shall despair to win after this?

strong a seat as a first-rate hog-hunter, and yet ride a horrible race, from not keeping a steady unjerking pull (not a dead pull) on the mouth. The man who has trained the horse is the only individual who can direct the most advantageous way in which he should be ridden; though the person that has daily acted as jockey, if not a goose, must, of course, be able to form some opinion likewise. If you are about to ride a hard-mouthed horse, always put a bit and curb on; and this should have been done in training: for if in taking his exercise-gallops he once runs away, he will be continually attempting the same trick: for having now got to know his speed, he will be on the look-out to break away with you in any part of the race, perhaps swerve, or bolt; and he will rarely struggle honestly when challenged for the rally in; but, while he should always be prevented from all chance of running off, you need not go into the other extreme, and throw him out of his stride, either at starting, or during the race, by an endeavour to restrain him too much when from over-eagerness you see he will not suffer it; it is better, in these cases, notwithstanding any instructions you may have received as to lying by, to allow the horse partly to make his own race: you can try to ease him a little as this over-eagerness subsides. In riding a match, a good caste horse against an indifferent caste one, always, after the first hundred yards, rate him well

the whole way so as to make him fairly shut up by the time he arrives at the distance post; for his object, if he is up to it, will be (thus over-matched) to make a waiting race of it.

You cannot expect to carry off all you start for the first season of your novitiate: added to muscle, bone, blood, make, a good constitution, good training, and good riding, you must use the greatest discrimination as to entering for a long or a short race, or long or short heats, and also to the weights. Some horses will fly with eight stone, yet prove very sorry with ten stone. Some will carry the ten stone well enough for a single race, yet fail in heats. Besides this, the course being hard or soft, light or heavy, dry or wet, up hill, down hill, or level, all and each make a wonderful difference in performance. There is not a horse in existence who, in some race or other, will not gain an advantage from one of these causes. Straight pasterns and tender feet, for instance, are not adapted to a hard course, nor slight legs and slanting pasterns to a heavy one; so, however confident you may be in having the best horse, never be too sanguine as to gaining: the best horse is rarely the best for all descriptions of races, weights, and courses.

Many beginners are much dissatisfied at strangers or acquaintances running down their horses. Nothing is so favourable. I never wish to hear a horse of mine

praised, unless on the day of sale. If a man tells you the neck is too thick, say yes, and the nostril is a little too closed: if he declares the quarter to be short, say yes, and rather wanting muscle; and, for every point he complains of, you name another. By good management, neither too acquiescing nor too differing, you may turn to good account a querulous visitor of this kind, and lighten him of a gold mohur for each of his mistaken notions. These first-sight, guess-work observations, half to three parts wrong on an average, are frequently changed by the utterers themselves in a week, and although your stud may not be perfection, yet, as all must be judged by comparison, it is not improbable that his and many others are much worse; consequently, instead of being discomfited, always give plenty of encouragement to whoever intrudes his critical remarks, strictly coinciding in their general correctness; and when the first pause occurs, ask, (if you know your horse,) what odds he will give against the ill-proportioned one? If he takes the bet kindly, do not be in a hurry; wait a day or two, then offer to double it: in this way you turn the tables on him genteelly. Every person has a right to look at your horse going his rounds, and also at rubbing down afterwards, when done on the public course. A by-stander can never dive into the real state of his condition by a few minutes' superficial glance at the external appear-

ance; it is more likely to puzzle and mislead,—so rather court than avoid inspection here. Your "dark" horse will not grow lighter by this little piece of complaisance; but if ever you admit either stranger, acquaintance, or friend inside his stable for half a second without being present yourself, it will be useless your denying your relationship to our "useful" friend at p. 261.

Stopping for the Feet.

Equal parts of tar and kidney mutton fat boiled together, and first laid over the frog only, then cow-dung over the whole. A piece of sponge, or thick numba, cut the shape of the sole, and kept over it by two transverse slips of bamboo fitting under the shoe, and wetted with hot water every three hours, should be used occasionally instead of the cow-dung, as it gives a nice soft pressure to the sole; a little of the tar stopping being always previously laid over the frog, if deficient, to encourage its growth and save it from rotting.

Ointment for the Heels, as a preservative against Cracks.

One pound of hog's or mutton-kidney fat, boiled down to the softness of ghee; two ounces of sugar of lead, well mixed in afterwards. The heels to be very slightly smeared over after washing at nine in the morning, and after being cleanly wiped out on return from walk in the evening. Whenever the slightest

crack appears, always poultice for one day with a little fat, linseed, bran, and a mashed turnip, carrot, or piece of melon, all boiled together, omitting the canter on the following morning; after which, equal parts of finely-powdered burnt alum and calamine powder, mixed, should be put on the cracks morning and evening, at the same hours as before, having first washed them with warm soap and water. The former ointment is not to be omitted.

Lotion for tender Shanks.

The leg, from the knee to the fetlock, often becomes very sore after sweating or galloping on hard ground, and he flinches when being hand-rubbed, which he never did before. Having fomented each leg for five minutes, as high as the knee, in a tub of rather hot soapy water, and *gently* dried them, warm and rub in *gently* with flannel, morning and evening, the following lotion :—

Bruised poppies..... 4 ozs. | Nim leaves ..... 4 ozs.

boil in four quarts of water; strain off two quarts, and add three ounces of Camphorated spirits.

Sal Ammoniac Lotion.

Crude Sal ammoniac (almost always procurable in the Bazaar) ..... 4 ozs.
Sugar of lead ..... 1 oz.
dissolved in three pints of vinegar and one of water.

PART IV.

TREATMENT OF A FEW DISEASES.

CLASS I. Sickness under Physic.
Overpurging.
CLASS II. Gripes.
CLASS III. Cold.
Sore Throat.
Influenza.
CLASS IV. Strangles.
Swelling of the Glands under the Jaw.
CLASS V. Fever.
CLASS VI. Inflammation of the Lungs.
,, of the Bowels.
,, of the Liver and Spleen.
,, of the Stomach.
CLASS VI. Inflammation of the Heart.
,, of the Kidneys, the Neck of the Bladder, and Bladder.
,, of the Feet.
,, undiscoverable.
CLASS VII. Red Urine.
Bursautee.

No one but a skilful professional man can possibly treat a quarter of the diseases of horses properly; and it often requires great experience, and the most able scrutiny, to be able to discern, or sometimes even to guess, what is the matter with a horse, or in what part he may be lame. This Part, therefore, while adapted to as full a practice as the most learned amateur can ever safely venture on, is written merely to prevent your outstripping the bounds of prudence: but these remedies must occasionally be resorted to, for, under inflammations and affections of the feet, &c., when assistance cannot be obtained, to do nothing, is to let the horse either die or be ruined. Remember, however, "prevention is better than cure;" and if you diligently read over Part III. once in every month, and follow in that tract, you will have little need for Part IV. But I must here again remind you, an Arab horse is not an English horse, nor is India England; and consequently, I advise you to adhere as strictly as possible to the directions I lay down, and not foolishly to alter the quantities of physic, or substitute other ingredients, or give them at different intervals than those prescribed, without first consulting some person who is capable of fully explaining to you the nature of the illness, as well as the virtues of the pharmaceutical compound.

In all diseases of the sudden acute kind, a horse's fate is decided in less than one-half the time a man's

would be; consequently, you cannot be too prompt in rendering relief; at the same time, never commence with either lancet or physic till you have formed some idea of the nature of the malady; the grand art consists in "giving the proper medicine, in the proper dose, and at the proper time;" so rather stand by the animal for half an hour with the book in your hand, watching the symptoms as they gradually present themselves; for there are few amateurs who would not very frequently save their horses, if they would only wait and discover the disease, before commencing with their ever-ready fleam and dirty aloes; after which it often becomes impossible to tell what the complaint really is; and hence the horse is lost.

CLASS I.

Sickness under Physic

is caused, firstly, by the ball breaking in the mouth, and half of it, perhaps, sticking there for a couple of hours; secondly, by the physic being too strong; thirdly, by the horse not having been sufficiently prepared, or by having been suffered to drink too much cold water, or to eat too much, or to other bad management.

Symptoms of the first:—The horse looks dull, hangs his head, and the saliva is black with the aloes.

Treatment:—Give whatever remains of the ball, adding another drachm or two to it; for, at least, that quantity is generally slobbered away before it is discovered; and wash the mouth out with salt and water.

Symptoms of the second:—Looks dull, and hangs his head, as before; generally coming on towards the evening of the day the physic is given. Treatment:—If you think you have given too strong a dose, keep him perfectly quiet, with a bed to lie down on. By not giving any exercise, and only very little grass and water when it commences to work, it will, perhaps, not operate more than is desired: if it does, see "OVER-PURGING."

Symptoms of the third:—Looks dull, hangs his head, lies down gently, and occasionally, though seldom, rolls, throwing himself on his back, and yet without appearing griped. Pulse, natural. Treatment:—Leave this state alone too: it will sometimes occur, even when the horse has been properly prepared, and the physic good; but if the uneasiness continues for an hour, or more, throw up a clyster of warm salt and water. At other times, all the symptoms of severe gripes come on, and you must be quickly on your guard. Back-rake immediately; then clyster with

Soap..... 2 ozs. | Common salt..... 2 ozs.
in a gallon of thin warm rice congee.

Then drench with

Oil of peppermint..... 1 dr.
or
Ginger..... 3 drs. finely-grated.
Epsom salts ..... 1 oz.
in a quart of thin warm rice congee

and throw three gallons of hot (hot enough to burn and make him flinch, but not scalding) water over his loins and belly. The whole of this, excepting the back-raking, to be repeated every hour; trotting *gently* in the interval.* If, after three hours, the symptoms should not be alleviated, and the pulse should rise beyond sixty, take away three quarts of blood, continuing the above treatment. Should the physic be operating, or immediately it begins to do so, there must be no exercise; the Epsom salts must be left out of the drench; and the clyster be composed of one quart of thicker congee, with a tea-spoonful of laudanum in it.

If a horse purges more than twenty times, he must be considered over-purged; and, if unattended to, inflammation may succeed.

Treatment:—Continue the gruel, as directed under "PHYSICKING," till you have given it six times; then change to the following:

Prepared or common chalk 4 drs.	**Catechu ..... 1½ dr.**
Gum-arabic..... 3 „	**Anise-seed ..... 1½ „**

* Trotting in such cases I disapprove of.—ED.

well mixed in a pint of arrowroot, or thick rice congee: give this every four hours. The legs to be, also, well hand-rubbed every four hours; and, in the interim, bandaged up as high as the knees in flannel or grass; and, if the weather is cold or damp, the jhool and head-piece to be put on. He should have a large soft bed, but be muzzled, if inclined to eat; and not a particle of grain, grass, or water given. After three drenches of the above, that is, after twelve hours, if the purging remains undiminished, add to it

Opium..... ½ dr. | **Alum*** ..... ½ dr.

and give the same, also, by clyster increasing the quantity of arrowroot in the clyster to a quart. There will be no danger of fatal inflammation, and very little of the bowels being injured, if this treatment is fairly adopted; but if you allow the over-purging to go on for twelve or eighteen hours, and then suddenly stop it with over-doses of opium or catechu, you will, most probably, as suddenly stop his breath at the same time. Should the purging, after another twelve hours, still continue unabated, the distress appear great, and the legs and ears cold, he will be in much danger: blister the belly, take away three quarts of blood, and give the medicine, both by mouth and clyster, every three hours.

* Half a drachm of the former and a drachm of the latter may be added with advantage.—Ed.

Blister for the belly.

Flour of mustard	**½ lb.**	
Spirits of turpentine	**1 oz.**	
Spanish flies	**2 drs.**	**finely-powdered.**
Linseed-oil	**½ pt.**	

to be made into a paste and spread on dungaree a foot square, with a jhool underneath it; or else spread on the inside of a sheepskin, and kept close to the belly for two or three hours, by bandages tied over the back.*

On recovering from violent over-purging, a seer of well-boiled ground grain, (gradually increasing to two,) one mäp of bran mash, and three drachms of gum-arabic, should be given at each feed for a week; wheat, or bajree flour gruel, sweetened, always at the first watering time in the morning; only two gallons of water at each drink, and that with the chill off; very little grass after each feed; a little dried lucerne, of the previous day's cutting, is best; and for the first three days, not moved from his loose stall.

But inflammation may also come on from the badness of the aloes, (if the stuff sometimes procured from the bazaar can come under that name,) or from improper ingredients being mixed up, or from the physic being given when the bowels were overloaded; as well as being caused by over-purging, and the then deadly sudden stopping of the purging with strong

* It ought, however, first to be well rubbed in for twenty minutes.—Ed.

astringents. The symptoms are the same. A quick small pulse, from fifty to eighty, and scarcely to be felt; heaving at the flanks; distressed countenance, and eyelids very red: the feet also become cold, and the hind leg, up at the stifle, sometimes trembles violently. Treatment:—Bleed four quarts, or more if the horse looks as if he could stand it: blister the whole of the belly, tying the jhool over it: put each foot in water, as hot as he can bear, for ten minutes, drying each quickly as it comes out; then rubbing till well warm, and bandaging half way up to the hocks and knees in flannel.* Clyster with thin, warm rice congee, if the bowels are not open; and drench, also, with a quart every hour. Put on warm clothing, if it is cold weather: but the irritation having been so great as to produce inflammation, he generally dies.

CLASS II.

Gripes

are caused by drinking cold water, especially when heated by exercise; or too much water immediately after feeding; or by exercise, immediately after feeding; or by over-feeding. By green food with the morning's dew on it; or too much green food sud-

* Instead of half way, it ought to be up to the hocks and knees in flannel.—ED.

denly given; or too much, even when accustomed to it. By change of grain. By the cold air. By want of exercise. Also, when the attacks are frequent, perhaps by some occult disease existing in the bowels.*

Symptoms.—The pain comes on quite suddenly; he paws the ground; looks round to his sides; rolls, and rolls over; tries to strike his belly with his hind feet; and breaks out into a perspiration: gets up again in a few minutes, shakes himself, and not unfrequently begins to eat. Almost all these exist, in a greater or less degree; and, after a short interval, they all return. The belly is sometimes tremendously swollen; I have seen it like a bullock's struck dead by lightning, and he groans heavily.†

Gripes.

Distinguishing Symptoms between it and Inflammation of the Bowels.

The ears, and legs, and feet are scarcely ever cold till after the perspiration breaks out; and, when rubbed dry and warm, they do not become cold again till after he breaks out into another sweat. In inflammation of the bowels they are always cold, and

* Gripes are more often caused by indigestion than any thing else.—Ed.

† This inflammation of the belly is from the presence of gas emitted by undigested and fermented food. The best remedy is, one ounce aromatic spirits of ammonia, and four fluid ounces of solution of aloes given in a pint of warm water, with copious clysters.—Ed.

though good friction will make them warm, they quickly become cold again.

The pain decreases after a trot; and when brought back to the stall, he often stands quite quiet for two or three minutes, as if perfectly recovered. In inflammation, the pain increases after a trot. There are, also, short intervals of ease during gripes, but none during inflammation.

In gripes, he often rolls quite over: in inflammation very seldom.

In gripes, he frequently commences picking at grass: in inflammation, I believe, never.

Rubbing the belly relieves the gripes; it increases the pain in inflammation.

The pulse is generally natural at the commencement of gripes, becoming fuller and quicker after a couple of hours or so, but not rising above sixty. In inflammation, it is never natural, but much accelerated at the commencement of the disease; not full, but small, and scarcely to be felt, and rising in four or five hours to eighty or ninety, or more.

Gripes, are not so often mistaken for inflammation of the bowels, as inflammation of the bowels for gripes.

Treatment.—Gripes from bajree and mhut. These are generally easily cured. Three drachms of finely-

powdered black pepper and a quart of hot, (do not burn his throat,) greasy mutton broth.*

Gripes from Curby.

Black pepper ..... 3 drs. finely-powdered
Mustard..... ¼ dr.
in half a pint of warm ghee.

Gripes from Green Meat.

Black pepper ..... 2 drs. finely-powdered.
Cayenne do. ..... ¼ dr.
Tincture of opium ..... 1 oz.
in a pint of warm congee.

Gripes from Cold Water.

Sweet spirits of nitre ..... 4 ozs.
Ginger ..... 1 dr. finely-grated.
in a pint of warm milk.

Gripes when severe, or the Belly is greatly swollen.

Linseed-oil ..... 6 ozs.
Spirits of turpentine ..... 3 „
Tincture of opium..... 1 oz.
Aloes..... 3 drs. finely-powdered.
Oil of peppermint ..... 1 dr. mixed.†

Gripes when the Horse is known to have Worms.

Linseed-oil ..... ¾ pt.
Spirits of turpentine ..... 3 ozs. mixed.

Gripes when the Horse has Sore Throat or Influenza.

Linseed-oil..... ¾ pt.
Sweet spirits of nitre..... 1½ oz.
Tincture of opium ..... 1½ „ mixed.

* The treatment recommended in the last note is preferable.—Ed.

† The aromatic spirits of ammonia and solution of aloes will be found preferable to this also. If there is not evident relief in two hours, the aromatic spirits of ammonia without the aloes ought to be repeated, and copious clysters of warm water and soap every half hour.—Ed.

Gripes when the Bowels are very Costive.

The same as that for Worms.

Gripes when the Bowels are well Open or Loose.

Tincture of opium ..... 1 oz.
Oil of peppermint..... 1 dr.
or
Ginger ..... 3 drs.
in half a pint of warm congee.

Gripes which come of themselves.

Onion juice.....1 oz.
in half a pint of warm ghee.
or
Aloes ..... 3 drs.
Tincture of opium ..... 1 oz.
in half a pint of warm water.
or
Brandy, gin or rum..... ½ or ¾ pt.
in a pint of warm water.

Should the first dose of any of these not relieve in twenty minutes, always back-rake; after that, clyster,* repeat the drench, and throw hot water over the loins and belly, and continue doing these last three every hour, till cure takes place, trotting gently in the interval; but always give the same dose twice, or, if you like, thrice, before changing to another. If it is a severe case, and no relief obtained after three doses, that is, after two hours and a half, take away three quarts of blood.†

* Clysters ought always to be administered from the first appearance of internal pain.—ED.

† See "BLEEDING," p. 81.

The first clyster, may be

Spirits of turpentine... 2 ozs. | Onion juice ..... 2 ozs.
in a gallon of thin warm congee.

When it is time for the second, for the bowels have not been opened, the clyster every hour should be

Aloes ..... 4 drs.
or
Epsom salts ..... 8 ozs.
Ghoor ..... 6 „
in a gallon of thin warm congee;

but if he has dunged freely, and the pain should still continue, each clyster should consist of two ounces of tincture of opium, in only a quart of warm congee.

Do not forget to change the clothing, if any has been on, that has become damp with the perspiration.

Doubtful symptoms.—If the symptoms are not very clear, and you are frightened lest it should be inflammation, give

Linseed-oil..... ½ pt. | Spirits of turpentine... 2 ozs.
mixed, which repeat, in half an hour, with
Tincture of opium* ..... 1 oz. added

and let the clyster be

Soap ..... 2 ozs. | Salt ..... 2 ozs.
in a gallon of thin warm rice congee.

For the first twenty-four, or, if the gripes have been severe, the first forty-eight hours after recovery,

* Half a drachm of unadulterated or purified opium, is reckoned equal to about one ounce of the tincture. If you have none of the tincture ready, macerate the half drachm in a small wine-glass of brandy.

give only one măp of boiled gram, and two of bran mash, at each feed ; the water with the chill off, and very little grass.

Gripes in India are very common, and the cordial quart of Hodgson, with

Ginger	**3 drs. finely-grated.**
Anise-seed	**6 „**
Ghoor	**3 ozs.**

is, after all, the most grateful cure : and one that is generally at hand. In giving it, you may omit the little point of mistaken politeness, you so generously offer to your friend at dinner, begging him "not to wait," but to swallow it down, fixed air and all, the instant it is poured out. You are now only treating the inside of a horse, and that under disease, which, having too much wind already there, will not be cured by another quart being thrust down his gullet. Pour the beer into a clean cooking pot ; then put in the ginger, anise-seed and ghoor, and stir it round whilst on the fire warming : in this way the stimulants become properly incorporated with the beer.

CLASS III.

Cold.

A thorough draught, or letting a horse stand still when heated by exercise, are almost as frequent causes of cold in this country as in England. Stripping off

the jhool in the cold weather, and then taking him out of the warm stall to be led for his morning's walk, or, at the gora-walla's option, to stand still, common sense must tell you will chill; and suddenly bringing a horse from the open mydan, in cold weather, into a close Bombay stable, has an eqūal tendency to produce cold or inflammation. During the months that are very cold, if you purpose riding at daybreak, and your horse is used to a comfortable stall, always keep as much clothing over the saddle as he had on at night, until brought out, and then, when you mount, give a trot for the first quarter of a mile. This may appear over-particular; but, on getting out of bed in January, at daybreak, walk out yourself without dressing.

Symptoms.—The same as in the human being.

Treatment.—Decrease, and boil the gram, and mix one măp of hot bran mash with it, or more, if he is fond of bran; decrease the grass also, giving lucern or green grass instead; take the chill off the water by boiling a handful of linseed, and a lump of ghoor in three quarts of water, then adding to it two gallons of cold: clothe him a little warmer than usual if it is cold, but let the stall be open, well open, and cool, not damp. If he coughs, put

Antimonial powder ..... **1 dr.**
Nitre ..... **3 drs. in 2 wine-glasses of hot water.**
Simple Oxymel ..... **3 ozs.**

mix and give it twice a day, before his morning's and evening's mash, which must now have less boiled gram in them; and do not exercise him beyond a walk. If the cold is very bad, he must not be moved from his loose stall, but clysters of warm soap and water used every other day. In some of these severe colds, when the pulse is much quickened, the glands below the ears perhaps swollen, the skin dry, and the running from the nostrils thick and plentiful, taking two or three quarts of blood is necessary, giving six ounces of Epsom salts in a pint of thin warm congee, every morning and evening, until it begins to operate.* Should the cold and cough not be of this severe acute kind, but remain hanging on him ten days or more, and the pulse be under forty-two, leave off the antimonial powder, &c., and resort to the mild beer tonic stimulant, p. 141, once a-day, in the morning; giving also half a drachm of assafœtida in half of one of the fried vegetable bringals, every afternoon. A little of the Bamboo-ke-puttah, is also useful in this case.

* The following treatment is better; half a drachm of digitalis, one drachm tartar emetic, one drachm aloes, and three drachms of nitre made into a ball, given morning and evening until the fæces become softened, after which the aloes ought to be omitted. The glands below the ears and the throat well rubbed in with the following liniment. Liquor ammonia two parts, spirits of turpentine one part, and olive oil three parts. The head well steamed twice a day and a hood put on. Clysters of warm water and soap every morning.—Ed.

Simple Oxymel.—Two pounds of honey and a pint of vinegar, simmered together over the fire.

If a horse takes cold, and, instead of becoming feverish, remains cold, perhaps slightly shivers, which sometimes happens, the hot instead of the cooling treatment may be begun with

Warm beer	**1 pt.**
Ginger	**2 drs.**
Camphor	**1 dr.**

dissolved in it may be given, which repeat in twelve hours, the gram being boiled, and mixed with a hot mash.

A horse is seldom the better for having a cold hanging on him a week or ten days, and nothing is more annoying than hearing a favourite one coughing; therefore these easy directions should not be neglected: besides, as colds of all kinds (both feverish colds, when the pulse is quickened, the skin dry, and the breath hot, &c.; and those when the pulse is not quickened at all, and the running from the nostril remains thin and watery) are so liable to lay the foundation of rheumatism, or leave the horse more or less debilitated, and pre-disposed to take on other disease, the beer tonic, p. 141, should always be given every morning for a week after the cold has gone off; or when given during a chill or lingering cold, be continued for a day or two longer.

For a horse in training, or that is sufficient in

vigour and spirits to work off a slight cold in his exercise,

Nitrous æther..... ¾ oz. | Warm beer..... ½ pt.
may be given, for three successive nights:
or

Nitre..... 4 drs.	**Opium..... ½ dr.**
Black pepper..... 1 dr.	**Honey or ghoor..... 1 oz.**
Anise-seed ..... 1 „	

to be put in the half pint of beer.

A hot bran mash should be added to the evening's gram, for these three nights, and the bowels also once opened by a clyster consisting of

Soap ..... 1½ oz. | Aloes..... 3 drs.
in a gallon of warm water.

Sore Throat.

Severe cold, or a cold neglected, may have sore throat connected with the cough, which is sometimes a serious affair.

Symptoms.—The food is not properly swallowed, but lumps of grass are often quidded out again. The water is not freely gulped; he sips and slavers in the dhool; it, in fact, hurts him to swallow it. The discharge from the nose is thick; mucus is coughed up, and the cough is so painful that he sometimes stamps with his foot. There is a wheezing noise when he breathes, and the glands below each of the ears are often swollen.

Treatment.—Take away *two* quarts of blood;* in

* See "BLEEDING," p. 81.

this case never more than three. Blister the throat. Give

Digitalis,* ..... ½ dr.
Antimonial powder ..... 1 „
Nitre..... 3 drs.

dissolved in two wine-glasses of hot water and then mix in three ounces of the simple oxymel, p. 210, three times a day. As much warm bran tea† as he will drink should also be given, and the dhool held up as high as the chest, that he may not be distressed by lowering his head. Not more than half a seer of boiled gram must be put into each bran mash, but lucern or green grass should be put in the trough as high as his chest. Dry grass irritates the throat; so, if no green meat is to be had, more boiled food, as turnips, carrots, &c., should be allowed instead. He must be left to his loose cool stall, with a bed, and well jhooled at night and the early part of the morning, if it is the cold weather. The digitalis is to be discontinued after three days, and the antimonial powder and nitre, after six. The bowels must be opened by clysters every other day, consisting of

Soap ..... **1 oz.** | **Epsom or common salt... 4 ozs.**
in a gallon of warm water.‡

* The treatment the same as recommended in page 194 with the addition of bleeding if the horse is in high condition, also blistering the throat sharply.—Ed.

† See "Bran Tea," p. 94.

‡ Clysters of warm water and soap ought to be used every day.—Ed.

This disease fortunately, is hardly known in India;* but severe colds and sore throats are not uncommon, both in the hot and cold weather; and if these were properly attended to and taken care of at their commencement, and the horse, on recovering, neither over-fed, nor allowed to drink too much gruel, which often brings back the complaint in all its force, requiring a second bleeding, added to the greatest care, to save him from danger, we perhaps should see nothing resembling it. The real epidemic may, however, some day be transported here: it is said to be very infectious, and that when it does visit a neighbourhood, many cases will occur about the same time.

Symptoms.—At the commencement of the disease, the coat is a little rough, the breathing somewhat hurried, the eyes inflamed, and the throat sore; hence, the swallowing is difficult, the food and water being sometimes returned through the nostrils: a discharge also takes place from them, as well as from the mouth, and the legs are swollen. Fever is present, and great weakness. The pulse varies, being in some patients only fifty; and in others, ninety; depending on the degree of fever.

Treatment.—*If* the pulse is full or wiry, and above fifty-five, and the membrane inside the nostril red,

* So far from this being the case on this side of India, it appears almost yearly in an epidemic form. I have repeatedly, year after year, had from two to three hundred on my sick list with this disease.—ED.

bleed three quarts;*† and *if* the dung is hard, give

¾ pint of linseed-oil with a drachm of ginger in it.
Clyster with a gallon of warm water and
4 ounces of Epsom or common salt.

Insert a seton in the chest, and also at the top of the neck, and blister the throat. Foment the legs with hot water, and keep the stall cool. The following morning, if the pulse still continues full and quick, and the membrane inside the nostril is redder than usual, bleed again three-quarts; but no more purgative physic is to be given. If a second bleeding is not required,—or, if it is, then, after twenty-four hours, give—

Carbonate of ammonia... ½ oz. | Nitrous æther..... ¾ oz.
twice a-day, at eight and four:

also,

Nitre	**2 drs.**	**Colombo-root**.....	**1½ drs.**
Gentian.....	**1½ „**	**Ginger**	**1½ „**

twice a-day, when the stomach is a little empty, at

* See "BLEEDING," p. 81.

† Bleeding ought to be resorted to with great caution, unless the horse is in high condition, and the inflammatory symptoms run high, it is likely to do more harm than good. Prostration of strength and debility is so great and rapid, that every drop of blood is wanted. The following treatment I have generally found successful. The bowels gently stimulated with from two to three drachms of aloes, and daily clysters of warm water and soap. The throat sharply blistered and the head steamed twice a day with boiling water poured on hay, and the following ball given morning and evening until the mouth is slightly touched. Calomel one drachm, linseed meal four drachms made into a ball with Venice turpentine. After all inflammatory and febrile symptoms have subsided, mild vegetable tonics to be given.—ED.

twelve and six:* these last balls may be washed down with a half pint of beer, or a quarter of a pint (a wine-glass and a half) of port wine, mixed with a wine-glass of water. Clysters, if the bowels are not open, and hot fomentations to the legs, must be used every day; and if the legs continue much swollen, they are to be scarified. The first twenty hours, while the physic is in his inside, feed on sweet fresh bran mashes and a *little* dried green food: after that, as it is the real epidemic, liberally on gram-flour gruel —three quarts, three times a day—boiled gram with bran mash, boiled carrots, fresh lucern, &c., and also thin gruel for drink—not water. Continue this for a week or ten days, when improvement or death will most probably have taken place; if the former, lessen the medicine daily, and be careful to keep the bowels open with clysters.

CLASS IV.

Strangles.

Strangles occur between the age of one and five, oftenest about three. There are three kinds of it. Strangle fever, without any abscess; true strangles, with the abscess under the jaw; and bastard strangles

* Instead of the nitre I would put a drachm of camphor in each of these balls.—Ed.

when the abscess bursts inwardly; but they are not so common in India as in England.*

Symptoms.—A slight fever, dulness, and disinclination to eat or drink occasionally comes over colts at two or three years old, either with, or without any cold, which keeps them weak and sickly for some weeks; and no abscess forming in the channel to mark the complaint, we are at a loss to account for the ailing: it may possibly be the strangle-fever. When an abscess forms in the channel under the jaws, then he has the true strangles; and it is most desirable that it should form, ripen, and be discharged; for the constitution is then said to be renovated by it. There is always a nasty discharge from both nostrils, with a choking kind of cough. When the abscess forms and bursts inwardly, it is called bastard strangles—though other forms of the complaint, and when it hangs long upon the horse, also come under that name. Sometimes during the strangle-fever and abscess, or, perhaps two, will form, not under the channel, but in some part of the body, such as the thigh, groin, &c.: this, fortunately, rarely occurs, as it renders the complaint very puzzling. True strangles, from mismanagement, may run into bastard strangles, and that into glanders; but bastard strangles has never the good luck to run into true strangles.

* All colts have strangles in India as well as in England.—Ed.

Treatment.—The strangle-fever, either with or without a cold, is to be treated the same as under "COLD," feeding on warm bran mashes, green food, sliced carrots, &c.; and if there is sore-throat, the same as under "SORE-THROAT." The true strangles is to be treated the same; but as the bleeding delays the abscess forming, he is not to be bled, unless the ears and legs remain cold, which shows that the lungs are becoming slightly affected. A tobra (not the one he eats out of) should be kept constantly half full of bran mash, and hot water poured on it every hour: the head being held over; this will promote the discharge from the nostrils, and be of great benefit; but if the breathing is difficult, this cannot be borne. Immediately the abscess begins to form the liquid blister is to be rubbed over it, and when soft and pointing, opened with a lancet; but the matter is only to be very gently pressed, never forcibly squeezed out. A poultice, kept continually warm, is then to be applied over the whole channel for two or three days; after which, a little Friar's balsam, or tincture of aloes, is to be daily squirted into the sore. Should the abscess burst of itself, enlarge the opening. When the abscess bursts inwardly, still promote the discharge from the nostrils with the tobra of hot mash, feeding on green food and mashes. If the abscess forms in any other part, instead of under the channel, treat the same. After the strangles are

over, a mild dose of physic is always requisite, followed by the "BEER TONIC," p. 141.*

Swelling of the Glands under the Jaw.

After five years old, a tumour sometimes forms in the centre of the channel under the jaw, without being accompanied by fever. A warm poultice should be applied till it becomes ripe, when it is to be opened, as before mentioned, with a lancet, and healed with he Friar's balsam.

A large hard swelling occasionally remains in the middle of the channel, the effect of strangles, cold, &c. Blister it, and if not lessened after a month, repeat the blister, or rub in the discutient lotion. If the glands below the ears remain hardened, treat them the same.

CLASS V.

Fever

is as common among horses as ourselves, and the after effects of it equally debilitating. More horses, I am inclined to believe, die from the effects of fever than from attacks of any of the acute inflammations;

* The treatment here recommended is judicious, but the best plan of steaming the head is putting some hay into a bucket, pouring boiling water over it, placing the bucket under the horse's nose, and covering up the horse's head with one end of the blanket or cumbly, and the other end hanging down and enclosing the bucket.—ED.

and for this reason, the danger is so often overlooked for the first three or four days, until inflammation of the lungs or bowels is approaching; then death is too near for bleeding or physic to be of any avail.

Symptoms.—The three principal and distinguishing symptoms of fever are, the great disinclination to exercise, the almost total loss of appetite, and the coat feeling dry and hot, and generally rough also, about the ribs. There is a peculiar soostiness all over the animal, but no expression of pain. A direct cold fit sometimes first occurs, as with us, and this comes on quite suddenly. The mouth is dry, the breath foul, and the tongue pale. The pulse may be quickened, or it may be weaker, or almost natural; but, during the fit, the legs, like the body, are more or less cold. When the cold fit is over, a warm one frequently succeeds, and a slight perspiration. In this state he remains, the fever often returning on the following day, and near the same hour. He is also generally flatulent, and the bowels are costive, but occasionally they scarcely at all alter, and the difference of warmth in the feet, some hot and others cold, often not at all remarkable until the fever has existed some days: but if at all thin before, the falling off in flesh and great prostration of strength, will be clearly manifest in forty-eight hours.

Treatment.—During the first cold fit, (not when

the warm one has commenced,) if a quart of warm beer, with three drachms of ginger, or a pint of port, and a pint of hot water, with a little spice were given, a jhool put on, and the horse gently trotted in hand for a quarter of an hour, (or let the trotting go on while the wine or beer is preparing, only be quick about warming it,) the fever might be nipped in the bud, and no more seen of it. Three horses I have cured in this way; feeding on bran mashes and green food the following two days. If unfortunately not observed at the onset, (which it rarely is, except by a man who lives half the day in his stables,) bleed* five or six quarts if the horse is fat and the pulse much quickened; but if the pulse is not much quickened, or not above fifty-five, never bleed, for the horse will frequently sink under it. Always back-rake, then clyster. Give

Calomel ..... ½ dr.
Emetic tartar ..... 1 „
Aloes ..... 1½ drs.

made into a ball, and ten hours after, another ball of the same.† After ten hours more, give

Linseed-oil ..... ½ pt.
or
Epsom salts ..... 6 ozs.
in thin congee;

and if the dung is not softened, repeat it after another

* See "BLEEDING," p. 81.

† Instead of giving linseed-oil or Epsom salts, I would continue the ball until the fæces become softened.—ED.

ten hours: nothing more, however, than three or four extra evacuations are allowable; purging is strictly prohibited. If the pulse was high and bleeding has been resorted to,

Digitalis	½ dr.
Emetic tartar	1 „
Nitre	3 drs.

are now to be given, morning, noon, and evening, in about three ounces of warm water* but leave out the digitalis after two days, or three, at most. Two drachms of cream of tartar may be given once in the day, as a cooling drink, when the digitalis is left off. The food must be warm bran mashes, with a little green grass or lucern; a little thin gruel occasionally; and the water have the chill off. Clothe warmly, if the weather is cold, making a large bed. The emetic tartar and nitre are to be left off when the appetite returns, and he becomes a little lively; but if the attack has been at all severe, you must not think of mounting him for a month at least, not even for a walk. Beer tonic, p. 141, should be commenced a week after recovery.

CLASS VI.

Inflammation of the Lungs.

Before commencing with inflammatory diseases, I must caution you against taking away a little blood,

* This ought to be given in a ball, and the digitalis continued until the pulse becomes intermittent.—Ed.

as a preventive when inflammation is coming on: this will render the disease very confused to you, whatever may be the case with a professional man. Wait an hour or so, until the disease fairly shows itself; the symptoms will then be fully developed, and you will know what part is attacked; and one full bleeding at the proper time, with the other treatment steadily pursued, will generally make a safe and speedy cure.

After an attack of inflammation of any vital part, it will take a fortnight, and often a month, before the horse can be again mounted. Boiled food; a little sweetened gram flour gruel in the morning before the first watering-time, and green grass, lucern, or carrots, are always needed to recruit the strength; but nothing is more dangerous than surfeiting with any kind of food after these attacks, in order to raise the condition quickly. Starvation, that is, as far as keeping the horse hungry, is the only safe system for ten days after recovery: the muzzle must be used at night if the appetite increases too quickly; but if, on the contrary, the appetite should flag, and the spirits not revive in that time, the beer tonic, p. 141, should be immediately resorted to; and even if the recovery is perfect, still, after ten days or so, a few mild beer drenches should be given. There is no disease in India, whether cold, fever, inflammation, or any

other, for which blood has been abstracted, and whether the appetite is regained or not, that tonic stimulants, of some kind, are not beneficial afterwards; and the beer-tonic, in proper quantities, is as good as any; the danger consists in commencing it too early.

Inflammation of the lungs is caused by sudden changes from cold to heat, rarely, it is said, by the contrary; by over-riding; by drinking cold water when hot, &c. It occasionally comes on quite suddenly: at other times the horse may have ailed a day or more previous to the attack.

Symptoms.—Veterinary writers have fortunately given us two symptoms, which mark this disease so clearly, that by common attention we can generally discover it: always, I may say, if not complicated with any other disorder. "The legs and ears are cold; of a deathy coldness; and the horse persists in standing, or, if he lies down, it is only for two or three minutes." The pulse is oppressed, rising to seventy or a hundred, and often imperceptible. The inside of the corner of the nostrils becomes of a dark crimson colour, the nostril itself being expanded, and the breathing greatly distressed. He appears stiff all over, and gently but anxiously turns his head round to his sides. Sometimes there is a cough; sometimes none.

Treatment.—Bleed six to eight quarts,* three to four (according as the horse is large or small, fat or thin) from each vein, *at the same time:* and it is particularly recommended in this inflammation to bleed from a large orifice with a broad-shouldered lancet, that the blood may flow freely and quickly. The bleeding over, back-rake; then clyster with

Soap..... 1 oz. | Epsom salts..... 4 ozs.
in two gallons of water,

and give

Aloes ..... 3 drs.
in half a pint of thin congee.†

Soak the feet and legs in hot water, then rub them

* See "BLEEDING," p. 81. Bleeding a particular quantity may appear to veterinarians equally as objectionable as giving a particular quantity of any medicine. Bleed, they say, till the pulse rises, &c., &c. I once saw a man bleed, and take a tea-cupful away, when he said the pulse rose. Bleed till the pulse rises, is about as useful a piece of information to many people as what I once received from a Surgeon I had written to regarding a horse that was dying from an inflamed vein; first, said he, "dissect it out." Bleeding a given quantity may, in some cases, be bad; but a quart too much or too little is better than a gallon too much or too little, and a certain quantity must be laid down. I have done the same with the dangerous medicine digitalis, &c.; for an acquaintance of mine once gave hellebore to a horse (strictly according to some professional book, he said) till the head drooped. It was done, however, too effectually; the head drooped for ever.

† This is dangerous treatment. If purging ensues from the aloes it is certain death. I would rather give very small doses of aloes, combined with digitalis, tartar emetic and nitre. The aloes to be discontinued immediately the fæces becomes slightly softened. Not a particle of aloes, however ought to be given until the inflammation has been somewhat subdued by the bleeding. The sides and briskets ought to be sharply blistered, and copious clysters of warm water and soap administered twice a day.—ED.

well till dry, and bandage them up to the knees and hocks in thick flannel bandages, or cover them round thickly with dry grass. If it is cold weather, put on a warm jhool; but the horse must remain in as cool a place as possible; if in a stable, every door and window should be open. In six hours after the bleeding, should the breathing still continue laborious, the ears and legs cadaverously cold, and he still stand in that peculiar stiff position, with the forelegs rather wide apart, open both veins again, and take from each another two or three quarts, and give

Digitalis	½ dr.
Emetic tartar	1 ,,
Nitre	3 drs.

made into a ball with linseed meal and liquorice water, or put them into a drench. Two ounces of the simple oxymel,* are also to be given once every eight hours; and clyster every day, for three or four days, if the bowels are not open, with

Soap ..... 1 oz. | **Epsom salts** ..... 4 ozs.
in two gallons of warm water.

Nothing must be given to eat for the first forty-eight hours but *cold* bran mash, (boiling water poured on bran, and allowed to stand till it is cold,) a handful of green meat occasionally, and the water with the chill taken off. The digitalis must be left out after two days; *i.e.*, after six doses, and the emetic tartar

* See p. 210, for the "OXYMEL."

and nitre after three days.* Beer tonic (p. 141,) to be commenced a week after recovery. If you blister the sides, the proper time for it is an hour after the first bleeding; but never do this in the hot weather, nor during the rains, unless it is cool.

Inflammation of the Bowels

is common enough everywhere. The causes are, cold suddenly applied; cold wind blowing on a horse's belly, picketed outside, that has been accustomed to a stable; or a draught of wind when under physic; drinking cold water, when hot; costiveness, unattended to; which last is often produced by gora-wallas, when *en route* by themselves, not searching for good water.

Symptoms.—The horse lies down, apparently without much pain; after a few minutes, gets up, and then lies down again. The breathing, is a little quickened; the pulse, at the commencement, increased to sixty or seventy; and the ears and legs rather cold. In the course of an hour or two all these symptoms

* A horse I had was attacked with violent inflammation of the lungs at six o'clock in the afternoon: he stood stiff as a post until two the following morning, when he fell, to all appearance, dead, and cold as marble all over. I was dining out, and did not return till that hour, but a veterinary surgeon was at hand. He was bled in both veins, back-racked, and clystered. Five hours afterwards, both veins were opened again: the horse recovered, notwithstanding the severity of the attack and the neglect of the first eight hours.

rapidly increase; the pain of the belly becomes very great, which is fully evinced on pressure; the pulse rises to eighty or ninety, and the ears and legs get much colder. His haggard countenance is often anxiously turned towards his flanks, as he groans and rolls; but he seldom tries to strike his belly with his hind feet, as in gripes. The bowels are always costive;* the belly is sometimes swollen, and the urine, if passed, is in small quantities, and with pain. The mouth is hot and dry; and the eyelids red, looking gorged with blood.

Treatment.—Bleed† from six to eight quarts; three to four quarts (according as the horse is large or small, and fat or thin) from each vein at the same time. The bleeding over, back-rake, and clyster with

Epsom salts ..... 6 ozs.
dissolved in two gallons of thin rice congee.

Foment the belly with hot water for half an hour, while a blister is preparing to be applied, about a foot square.‡ Give

Linseed-oil ..... ½ pt.

which repeat every seven hours till an evacuation takes place; but no purging is allowed here any more than in the previous disease; merely the removal of the costiveness, so as to cause a softened

* This is not always the case. Inflammation of the intestines is sometimes accompanied by severe diarrhœa.—ED.

† See "BLEEDING," p. 81.

‡ The blister ought to be well rubbed in, to the belly and sides.—ED.

state of the dung.* The legs must be well hand-rubbed to restore the circulation, and then bandaged with flannel or tied round with grass. A jhool, if it is cold or damp, should be put over the body, but the horse kept in a cool place. Warm bran mashes, with two drachms of finely-powdered gum-arabic in each mash; linseed or bran tea; thin gruel, and a little green grass, is all the food allowed for the first three days. If the bleeding has been properly performed, and none of the other treatment omitted, the force of the disease generally yields in six or eight hours; but if after that time the inflammation should not be subdued, and the symptoms of uneasiness and pain continue as at first, bleed again two or three quarts from each vein, and draw the firing-iron over the belly two or three strokes, a little behind the blister. The costiveness must be fully removed before the inflammation will subside; but immediately the dung is softened, no more physic is necessary, nothing but a soothing congee clyster.†

Inflammation of the Liver, of Spleen.

The first of these is much more common at many stations than the preceding disease. The causes are,

* If there is constipation, I would at once give a dose of Aloes, guarded with Opium.—Ed.

† A handsome Arab, that had gone through a morning's parade without the slightest sluggishness, was attacked with inflammation of the bowels when led to his stall. The bleeding, &c., having been delayed for twelve hours, he died on the following morning.

the climate, added to not giving physic, or giving that which is of no use, when, from a foul habit, there has long been evident necessity for it.

Symptoms.—Are somewhat between inflammation of the lungs and bowels. The pulse increases as the disease advances, and rises to seventy or a hundred. The horse will sometimes stand, but not with his forelegs so fixed and apart as when the lungs are affected. The eyes, mouth, and nostrils are always more or less yellow; but the legs are not so particularly cold. The bowels are always costive, unless the inflammation is very trivial, and then there may be a slight looseness. The head is, generally, being continually turned to the side that is affected, and which, if pressed on, near the middle of the false ribs, will be painful.

Treatment.—Bleed* from six to eight quarts. Blister the side you think affected, about the middle of the false ribs for about a foot square, six inches above and six inches below the finish of the ribs; back-rake, and clyster with

Aloes.....6 drs.
dissolved in a gallon of warm water;

and give—

Calomel.....1 dr.
Emetic tartar.....1 „
Aloes.....1½ „

* See "BLEEDING," p. 81.

made into a soft ball, every six hours, until the dung is well softened; but if the bowels were relaxed from the commencement, which is the case sometimes, though very rarely, bleed not more than from three to four quarts, and give—

Calomel ..... 1 dr.
Opium ..... ½ „
Chereeta ..... 1 „
every twelve hours, for three times;

after which,

Linseed-oil ..... 1 pt. | **Congee** ..... 1 pt. **mixed.**

Diet to consist of warm bran mash; water with the chill off; and, after twenty-four hours, a little dried, green grass. On recovering, you must rigidly attend to the leading paragraph, CLASS VI, p. 221, for a relapse is very likely to occur; to prevent which, a dose of physic should be given about three weeks after, consisting of

Calomel ..... 1½ dr. **over-night.**
Aloes ..... 4 **drs. in the morning.**

Inflammation of the spleen seldom takes place. Symptoms and treatment the same as "LIVER."

Inflammation of the Stomach

may also be said to be very rare, and not to be distinguished from inflammation of the bowels. Symptoms and treatment are the same as in that disease.

Inflammation of the Heart

is another very rare disease, and more resembling

inflammation of the lungs. If a horse dies of the latter, the former may be found inflamed; the same as when a horse dies of inflammation of the bowels, the stomach on dissection may also be found inflamed: but, as primary diseases, they are both allowed to be very rare.

Inflammation of the Kidneys; the Neck of the Bladder; and Bladder,

are caused by cold; continued rain dropping on the loins; over-riding, or strain of the loins; not allowing the horse to stale, if he requires it, whilst riding; long feeding on musty gram, &c.

Symptoms of inflammation of the Kidneys, the Neck of the Bladder, and Bladder, are the same.—The mouth is hot and dry; the pulse greatly increased; and the head anxiously turned towards the flanks, as under other inflammations: but here there is often observed a stedfast gaze towards the affected part; the head being turned over the body, and the muzzle almost put on the loins, instead of slightly directed towards the belly. The most distinguishing symptoms, however, are the hind legs. He stands wide apart; straddles broad when walked; and shows great pain, by shrinking when the loins are pressed on. The testicles are occasionally drawn close to the body, and then let down again. He strains to stale, and the little that is passed is high-coloured, or

bloody; but he is not so continually lying down and getting up again as when inflammation of the bowels is just taking place.

Treatment.—Bleed,* from six to eight quarts, three to four from each vein, at the same time; back-rake; clyster with

Linseed-oil..... 4 ozs. | Aloes..... 4 drs.
mixed in four quarts of thin warm rice congee.

Give a drench of physic, consisting of

Linseed-oil..... 4 ozs. | Aloes..... 4 drs.
in a pint of thin congee.

Put a mustard poultice† over the loins, laying the same on the inside of a sheep-skin, if at hand. A blister is not allowed; so if the mustard poultice cannot be had, foment the loins every half hour with nearly boiling water, and keep them well warm during the intervals. Cover up with a warm jhool, if in cold weather: hand-rub the legs, and bandage them up to the knees in flannel, or grass; but keep the horse in a cool place. When the physic has operated,

Digitalis ..... ½ dr.
Emetic tartar ..... ½ „
Gum-arabic ..... 2 drs. finely-powdered
made into a ball with linseed meal or ghoor,

should be given once in every eight hours: the digitalis

* See "BLEEDING," p. 81.

† Mustard Poultice:—Mustard-flour and linseed-meal, equal parts: mix them together with an equal quantity of hot vinegar.

to be left out after six doses, and the emetic tartar and gum after ten. Warm bran mashes; water with the chill off; and, after twenty-four hours, a little dried green grass may be allowed for the diet. Clysters of plain warm congee should be used once in every eight hours during the day of the attack, in order to foment and soothe the inflamed parts. On recovery, treat as under leading paragraph, CLASS VI., p. 221. Should the hindlegs remain stiff for more than a fortnight after the disease is subdued, put a charge over the loins.*

Inflammation of the Feet

is caused by hunting, or hard riding, or driving over stony ground; by severe training, particularly if the course is hard; by washing the feet in cold water whilst hot, or, *vice versâ*; when in a high latitude, by putting the horse into a hot stable, and bedding the feet up in deep hot grass the instant of return from the cold air; or, by inflammation being transferred from the lungs, or any other organ, to the feet. The two fore-feet only are generally attacked, but sometimes all four.

Symptoms.—The pulse becomes greatly increased;

* Charge:—Pitch, one pound; tar, one pound; bees-wax, quarter of a pound; resin, two ounces; heat them together, and, when a little cool, spread thickly over the loins, strewing over flocks of short tow immediately it is on. This will stick fast for six weeks, or two months; during which time the horse must be well walked; but not mounted.

the inside of the nostrils red; and the countenance distressed, as in other inflammation: but the horse is here evidently very uneasy on his legs, and when he lies down the muzzle is often rested directly on the affected feet, which are always found intensely hot; and the arteries on the sides of the pastern throbbing distinctly. These last symptoms are what we are told to rely on, as the distinguishing marks of this complaint. He occasionally breaks out into a sweat, but never attempts to paw with his feet, or to kick at his belly; and when down, though he may groan quite as much as when any of the previously mentioned vital organs are the seat of attack, yet he never rolls. Under any other inflammations also that had existed only an hour or two, the horse would be able to walk without exhibiting any pain in the extremities; but in this, the fore-feet are lifted up and put down again, similar to a man suffering from broken chilblains, and having a boot on; notwithstanding, you will seldom discover the disease till all the symptoms become fully marked.

Treatment.—Take the shoes off very gently, filing away the clenches, and slowly withdrawing each nail separately, for the pain is intense. Thin the sole all over as well as you can, cutting away small chips from the crust also, with the drawing-knife, and then filing even. You must do this when he is down, if he is unable to stand to allow it. Bleed three quarts

from each affected foot;* if you cannot get that quantity, take as much as you can, and then bleed six quarts from the neck vein. The bleeding over, put the feet into warm poultices of bran and linseed meal. Back-rake, clyster, with warm soap and water, and give the mild aloetic salt drench, p. 96; also—

Emetic tartar ..... **1½ dr.**
Niter ..... **2 drs.**
Cream of tartar ..... **2 „**

morning and evening, for three days, dissolved in half a pint of warm water.

Make a large soft bed, that he may lie down as much as possible; but, for the first three days, the diet must be only bran mash; water with the chill off, and occasionally a handful of green grass. On the following day after the attack, if the inflammation has not subsided, bleed again, three quarts from each foot; and if the horse is fat, change the warm poultices to swabs of cloth, which wrap round the hoofs, keeping them well wet with

Nitre .... **2 ozs.**
dissolved in a quart of the coldest water.

On the third day, if the inflammation still remains, the pastern, round about the coronet, is to be blistered; but unless the blistered part is well covered up with a cloth, and also something put round the neck to prevent him getting at it with his mouth, he

* See "BLEEDING," p. 81.

will tear it off, and the pastern will then for ever remain denuded of hair.

Inflammation Undiscoverable

would not be so common, if people would only take the trouble to read over quietly the different inflammations in the stable, in presence of their horse when reported sick; yet some, who do take that trouble, are often in such a fright, if the horse is valuable, or in such a hurry, that they fail in catching the symptoms indicative of the part affected.

If the lungs are the seat—the purple nostril; the stiff fixed standing, with the forelegs rather apart; and the deathly clay-like coldness of the ears and legs, are the principal distinguishing signs.

The bowels—the redness of the eyelids, when turned down; the lying down and getting up again; and coldness of the ears and legs.

If the liver—there will be yellowness about the eyes, added to the other symptoms of inflammation of the bowels: and if you here make a mistake, the treatment for this will possibly cure the other, although, of course, not so well, nor so safely, as purging is dangerous when the bowels are the seat.*

If the kidneys—the stradling of the hind legs, and shrinking on pressing the loins.

* Not when your Aloes are good, and guarded with Opium.—Ed.

If the feet—they are hot, instead of cold; the muzzle often resting on them when lying down, and the great disinclination to stand.

The note at the bottom of the page, however, ought to serve as a guide and warning, to convince yourself that inflammation really does exist in some part before you commence to cure.*

CLASS VII.

Red Urine

in India, must be considered as arising more from some little derangement in the digestive functions,

* A horse that had been for some months having only very gentle exercise, was taken out and galloped after a hog. On return to his stall, he immediately lay down. The alarmed master, without feeling the pulse or asking anybody's opinion, took five quarts of blood. The poor animal not being much refreshed by this, a friend recommended he should be clystered and physicked. The horse, after this, getting more "gureeb" still, they both allowed that it was a most extraordinary case, but that, as he appeared so near dead, bleeding again could not do harm, if it did not do good. At this stage, as I lived close by, my advice was solicited. I had seen the horse on his first return: his case was plain enough—a little over-galloped when not in wind, and brought home hot. Out of pity for the poor brute, I undertook to do all I could, saying, I had a bottle of Elixir lately sent me from London, just adapted for these obscure cases.

I then mixed some sawdust, red-ink, and blue-paint, in a quart of sour claret for the master to smell at, and putting the sufferer, half-dead from treatment, into a loose open stall, with a large soft bed, washed his mouth out with the Elixir of warm water, and left him with a little green grass till the following morning. The horse survived; and the owner gave me a gold mohur for a pint of the mixture; and his hunter, from henceforth, went by the name of, Impostor.

than as a primary disease of the kidneys ; and what would alarm in England, is here allowed to pass almost unnoticed. The native remedy is

Ghoor and ginger..... 6 drs. each
mixed, and given daily, for three days ;

and if the pulse is not increased beyond its natural standard, half of this may be tried. If there is too much excitement, a pint of the dhye (sour milk) sweetened with a small lump of ghoor, and given every morning, would be preferable. Some boiled food, with a little bran mash, and dried green grass, or lucern, should never be forgotten.*

Red urine, as a disease of the kidneys, or from inflammation, must be treated of separately.

Bursautee.

Various are the remedies that have been employed for this disease, and it is rather unconsolatory to find, that no veterinary surgeon, at either of the Presidencies, has yet condescended to favor the public with a paper on the subject. If, then, we are to be doomed to grope out a cure ourselves, I cannot too soon add my ideas to those of other amateurs, who

* Bloody urine where there is no disease of the kidneys, is generally caused by indigestion. I have seldom found the following fail in effecting a cure. Aloes one drachm, Nitre four drachms, made into a ball with honey or treacle. One to be given morning and evening until the fæces become softened.—Ed.

have already written for our benefit in the "Sporting Magazine;" such recommending the madar, blue vitriol, &c.

In accordance with its name, the months of June, July, August, and September, generally the two middle ones, are the periods at which this disease breaks out. The worst cases are always in low, damp, and "feverish," situations. It rarely occurs in a bad form in the Deccan; a dry "liverish" climate, like this, is one of the most effectual ingredients in the cure. When a horse is predisposed to break out with bursautee, and happens to be at an unfavourable station, and he is at the same time neglected, the usefulness of the animal is then destroyed for seven or eight months, and he will too often retain the scars, and loss of hair, for ever. It being my opinion, that the cause lies in a constitutional predisposition, which cannot be discovered till the complaint appears, and the only likely safeguard, against its breaking out severely, being a dry climate, I shall proceed to recommend a trial of that which in most cases will be found to expedite the cure. On the first appearance of the disease give a mild, warm drench of physic,* consisting of Aloes and Epsom salts, with one drachm of ginner, in rice congee.† Three days after the physic

* See "PHYSICKING," p. 87.

† I would rather give a ball, consisting of aloes and ginger.—ED.

has set, give half of a common masallah ball daily, for three days, then omit one day, and commence with

Cantharides	**2 grs.**	**Gentian or Chreeate**	**1 dr.**
Ginger	**1 dr.**	**Anise-seed**	**1 „**

in a ball;

this to be given every evening, after the last seven o'clock feed. After six days increase the cantharides to four grains, and after twelve days, to six grains. After eighteen days, if the appetite improve, increase the cantharides to eight grains. After twenty-four days, discontinue the ball altogether for three days, and then commence again with the first quantity of only two grains, increasing to the second, as before, and so on through the whole monsoon.* The food during this time, whether the horse is fat or thin, is to be boiled sago, boiled barley, boiled oorud and sheeps' heads,—try everything to induce him to eat the sheeps' heads, or in default, any strong meat broth,—and the more black salt he will *willingly* eat with his gram and bran mash the better.

If the horse is gross, and unable to take much exercise, from the largeness of the sores or swelling of the limbs, still liberal (not over) feeding on this kind of diet cannot be dispensed with: you must muzzle

* Instead of the foregoing, I would give the following ball morning and evening. Cantharides five grains, Dinioclide of copper half a drachm, Gentian two drachms, Pimento one drachm. The Cantharides to be discontinued for a few days, when the secretion of urine becomes increased. —Ed.

occasionally at night. Lucern and green grass, cut the day before, I am also friendly to in small quantities. Keeping the horse in a dry loose stall, well littered at night-time, is of course to be remembered, and as much walking, or gentle trotting exercise should be given, morning and evening, as possible, for which you may as well take off the shoes. The external application for the sores (and to apply which, you should endeavour to obtain the assistance of a clever native farrier) is the native poultice :—

Leem ke putta, as much as the size of an egg.
Chitrawal ke putta, ditto.
Vikmar, as much as two peas, or two gram.
Fulkeree, a quarter of a rupee weight.*

When the sores are small,

Koorkum-ketail.. ..... 4 tea-spoonsful,
Blue-stone ..... ½ tea-spoonful finely-powdered.
Alum..... ½ a one, all mixed,

will frequently dry them up. As a preventative for the ensuing monsoon, I should recommend the same treatment to be commenced with on the 1st of May, and carried on to the 1st of July, but not increasing to the


* I have seldom found the following applications fail in effecting a cure, when the Cantharides and Dinioclide of copper ball is given at the same time. The sores to be touched every second morning with the following escharotic solution, with a feather, until they look healthy. White wine vinegar two ounces, three and a half drachms, Bisulphate of copper, two and a half drachms, Sulphuric Acid, three drachms—and dressed morning and evening with the following ointment. Carbonate of zinc, one ounce, Venice Turpentine, one ounce, Suet, six ounces. The feeding throughout to be liberal.—Ed.

eight grains of cantharides, unless some symptoms of breaking out again show themselves, in which case you may gradually go up to ten grains.* The cure, of course, consists in eradicating all tendency to the disease from the constitution, as the sores would generally heal of themselves by October or November. Many persons are advocates for the application of the hot iron, or the caustic madar, to the sores, and giving large doses, internally, of blue vitriol in solution. It is not for me to say anything against this treatment in severe cases to those who understand how to use these remedies, but there is no more analogy between bursautee and farcy than there is between a common cold and glanders.

LAMENESS.—Ed.

[As "Blunt Spurs" does not describe lameness beyond "inflammation of the feet," and as I consider it desirable that more notice should be taken of so important a subject, I shall point out as briefly as possible the parts generally affected, the causes that give rise to lameness, as well as the treatment required to be adopted; but before proceeding to do so, being convinced that in some seven cases out of ten, the seat of lameness is in the foot, I will here copy from the "Illustrated Horse Doctor" Mayhew's foot bath, as tending materially to alleviate suffering and restore Horses lame in the feet, to a state of usefulness, in a much shorter period than any other means at present adopted.

* I prefer the ball given in the last note at page 238.—Ed.



"Of all inventions intended to mitigate the sufferings of the horse, none perhaps is so generally useful as the foot bath; certainly, not one is so decidedly beneficial in its operation. It consists merely of a wooden or iron trough, one foot deep, the shoes of the animal should, if possible, be taken off before the hoof is allowed to tread within the bath, or, if such a measure be not possible, then the burden of the horse's body, should be counterpoised by means of weights. This precaution is always prudent, for should the shod horse occasion fracture or breakage, an alarm might be excited which probably would ever after prevent the employment of the foot bath for the same quadruped.

"The water should always be mixed, without the building, it is never well to excite an animal's fears by allowing it to witness unnecessary preparation. The author is fully aware that most people assert the horse has a very limited comprehension, so it may have; but it has an active terror which is apt to misconstrue the simplest of motives; whoever has seen the busy eye of the quadruped watching all that takes place around it, and noting every triviality whenever any unusual movement gives intimation

to the animal that something is about to be attempted, will readily allow the need there is for excessive caution. The horse may comprehend nothing, but it is not, therefore, the less to be propitiated. Its terror has to be soothed and its confidence has to be gained: the last is soonest won by avoiding anything which possibly might excite the first.

"Always have the heat of the water ascertained by a thermometer. Sensation is only a relative test with regard to the presence or absence of warmth; were it not so, the coarse hand of a groom, nevertheless, might easily endure that degree of temperature which should pain the foot and leg of a horse. Let the fluid, in the first instance, stand at 70°, after the animal has entered the bath, gradually and without noise, increase the temperature up to 90°.

"At that standard the water ought to be maintained: the hoof should remain soaking from four to six hours at each operation; the groom, doubtless, will complain of having frequently to fetch warm water, and when not so employed, of being obliged to watch a thermometer; but the present book is not written to please the likings of any individual. To contribute to the welfare of the horse is the object of the writer; that he has not unnecessarily imposed an irksome duty upon any human being, the purpose for which the bath is introduced into the stable should be sufficient evidence."

"The horse's hoof is of considerable thickness, it is far from unusual with stable men to saturate the healthy hoof with various greasy preparations, therefore it will require some time before the heat and water can soften that which is, as it were, prepared to resist their action. The hoof should be rendered perceptibly soft when the object is to relieve a painful lameness: the warmth and moisture should not only saturate the covering to the foot, but should also soothe the internal structures. The pressure of the horn may thus be mitigated, and the deep-seated inflammation likewise be ameliorated.

"When the bath is removed, the foot should not be left exposed to the air, as the horn then quickly dries; it soon becomes harsh and brittle. In this condition it is likely to do more injury to the sensitive parts than good which was anticipated as the consequence of its immersion. The hoof when taken from the water, should be encased in warm and air-proof bandages: the intention being to retain the heat, while evaporation is prevented. The bandages likewise answer another purpose; they protect the foot, which being without a shoe and covered by horn, that has been deprived of its resistant property, is, therefore much exposed to accidents.

"To obtain the full benefit of the bath, the animal should enter it night and morning, the foot should be subjected to its operation for at least four hours each time, and the ingenuity ought to be exerted to prevent the hoof from becoming dry in the interim. Perhaps nothing is better for this purpose than the leather case, which is lined with sponge, and which can be procured of most tradesmen who deal in Veterinary Instruments. It is made to fit the foot, also to envelop the pastern. The bottom portion is formed of the stoutest leather, and will afford all desirable protection, while the sponge will retain the moisture which this material permits to be renewed should circumstances, such as the heat of the hoof, or the warmth of the weather, cause the fluid to evaporate. However, such additions must always be made with warm water. Cold water being unsuited for the purpose.

"These particulars have been thus fully detailed, because lameness constitutes no inconsiderable portion of equine misery, and because such ailments are more frequently encountered, than special forms of disease. To judge quickly and surely of such affections proves, in no small degree, Veterinary proficiency. In every shade of lameness, the gentlemen, unless more than usually practised in such ailments, had better be guided by an educated opinion. Where it is possible to mistake another's

misery, it displays no boldness to risk chances upon our own judgment."

Quittor

arises from injuries inflicted on the coronets, as well as on other parts of the foot. Ascertain with a probe the depth and direction of the sinuses, and inject into them a solution of the sulphate of zinc (20 grains to an ounce of water): should suppuration have taken place to any extent, it will be necessary to allow the pus to escape by opening out the sole, and removing such portions of the horn as have become separated from the sensitive part of the foot. It is a serious matter when the disease assumes this form, and the sooner the patient is placed under professional care the better.

Thrush

is caused from neglect and want of cleanliness. The frog becomes diseased and emits a very offensive discharge. Pare out the foot and poultice it for a few days, after which have the horse shod. Keep his feet dry, and with the assistance of a piece of bamboo in the shape of a spatula, insert into the cleft of the frog as deep as you can, equal parts of Calomel and Alum.

Corns

are of common occurrence and often produced from bad shoeing. They generally appear in the inner quarters, and are caused by the heels of the shoe pressing on the sole. Inflammation is set up in the sensitive part of the foot, the minute blood vessels are ruptured, and hence the red patches called corns.

Pare out and poultice foot, until lameness is removed, and then, with great care put on a bar shoe.

If pus has collected in the foot, act as directed for Quittor.

Sand Crack

generally appears in the inner quarter of brittle fore feet, extending for an inch or so from the coronet. It should be opened

out, the foot, poulticed, and when sufficient sound horn has grown from the coronet, pass the firing iron over and under the crack, and when the horse goes sound, shoe as described in the annexed cut.



Pumice Foot

results from inflammation of the foot when the toe of the coffin bone presses on the sole, and causes it to present a concave, instead of a convex surface. A horse so affected is of little or no value, and all that you can do is to put on a bar shoe, and protect the sole with leather.

Bruises of the Foot

are caused by shoes pressing on the sole, or this portion of the foot being pared away to such an extent, as not to afford sufficient protection to the sensitive part, which becomes injured whenever the sole comes in contact with a loose stone. Treat as recommended for corns.

Canker

is a very obstinate and often incurable disease, even under the most skilful treatment. It is frequently the consequence of thrush. Poultice the foot, remove every particle of diseased horn, and dress with Sulphuric Acid, and Tar, one ounce of the former, to one pound of the latter.

Navicular Lameness

may be attributed to a variety of causes. The symptoms are pointing and contraction of the affected foot. Blisters, setons and rest will afford relief, but the Horse will go lame again when put to work. Neurotomy is the only means by which he can be made serviceable.

Injuries from bad Shoeing

arise from nails penetrating, or pressing on the sensitive parts

of the foot. If the former is the case, the horse will show lameness at once. The shoe should be removed, the hole made by the nail opened out, and the foot put into a poultice. If the latter, lameness will not appear for some days after shoeing. On examining the injured part, it will very likely be found in a state of suppuration, open it out well, and after searching it with a probe, remove as much of the horny sole as is separated from the sensible one, and dress with tar and large pledgets of tow, so that there may be pressure on the parts pared out, keep down proud flesh by applying Sulphate of Copper (blue vitriol) or Caustic to it, and when the sole is sufficiently strong, protect it with leather, and give the horse moderate work until you can dispense with it.

When you suspect that a horse is lame from shoeing, get a hammer and strike on the clenches all round, when the hammer comes in contact with that part of the foot that is injured, the horse will evince pain and endeavour to extricate his foot.

Punctured Wounds

are caused by horses stepping on nails, treat as for injuries arising from defective shoeing, but it sometime happens that the nail penetrates the navicular bone, or the tendon inserted into it, in either case, the lameness is incurable, and the sooner the horse is destroyed, and relieved from his sufferings the better.

Having been described at page 10, I have here merely to point out the treatment. Reduce the inflammation in the parts, with cold application, and then blister repeatedly if necessary. Should lameness still continue, fire, and give the animal rest before you put him to work again.

Bog Spavin, Thoroughpin, and Wind Gall

are bursæ distended to an unnatural size with Synovia, and are generally produced from violent exercise. They are gradually developed and scarcely ever cause lameness, but are very

unsightly. Should they however interfere with the usefulness of a horse, treat as for bone spavin.

Curb


is frequently caused by horses being brought suddenly upon their haunches, and arises from sprain of a ligament and tendon that pass behind the back part of the hock, and bears the appearance described in the following cut No. 1 from Mayhew's Horse doctor. Treatment as for sprain.

No. 1



* Curb.

No. 2



† An Indian Rubber bandage for keeping wet cloth on a curb.

Strains of the Tendons or Back Sinews

are produced from violent exercise, and cause lameness, heat and swelling. If the injury be very severe, put on a shoe with raised heels, reduce the inflammation with cold applications, (ice in preference to anything else); after this is done, and should the tendon remain enlarged and painful, blister, and fire, if necessary.

Shoulder Lameness

is of rare occurrence and may arise from a sudden turn at exercise, or from entangling the leg with the head rope. The lameness is very great, and the horse threatens to fall at every he takes. Foment frequently with hot water; when

inflammation has subsided, put on a mild blister, and let the animal have rest.

Hip Joint Lameness

arises more frequently from falls than anything else, when in motion, one hip appears higher than the other. Treat as for shoulder lameness.

Dislocation of the Patella

is caused when a horse is suddenly turned round, whether standing still, or in motion, the leg is quite stiff and stretched out behind, and when the horse moves it is jerked forward: walk him, and whilst he is advancing the injured limb, press on the out side of the stifle, until you hear the patella slip into its proper place; lead the animal into his stall, tie up his head, blister the stifle, and after a few weeks' rest, he will be fit for work again.

Lameness from cutting

arises from defective shoeing, weakness, or awkwardness; the inner portion of the fetlock joint is the part that suffers from the foot of the other leg coming in contact with it. Sometimes an injury is inflicted between the knee and fetlock, and is called "speedy cut." If there is much heat and pain, use a bran poultice, and when this is no longer necessary, apply a little tincture of Iodine or Myrrh to the part abraded.

If you think the injury has been caused by defective shoeing, remedy it by raising or lowering the quarters, but never think of rasping away the inner portion of the foot. If from weakness, give the animal a liberal supply of food, and rest till he regains his strength. If from awkwardness, use a boot.

Rheumatism

arises from exposure to cold: sometimes all the legs are affected. The patient is feverish, pulse and respiration are accelerated, and the bowels constipated. At other times, the attack is not

so severe, lameness exists but in one leg, this no sooner recovers, than another goes wrong, and so on till all four have had their turn.

If a small horse, give

Aloes ..... **4 drs.**
Calomel ..... **½ dr.**
Gentian with ginger, &c ..... **2 drs. to form a ball.**
Aconite ..... **20 or 30 drops**
in half a pint of congee,

every third hour, until the pulse is natural; then give every morning a ball composed of

Nitre ..... **4 drs.**
Camphor ..... **1 dr.**
Gentian ..... **2 drs.**

Rub the following liniment on the legs morning and evening, and cover them with flannel bandages.

Spirits of Wine ..... **1 qt.**
Camphor ..... **2 ozs.**
Sweet oil ..... **2 „**

In all cases of lameness do not omit to take off the shoes, and give a dose of physic. Prepare the horse for it by putting him on bran mashes for two or three days; but if it is necessary to act on the bowels without delay, give him a ball at once.

If you are treating horses just imported from England or the Colonies, do not attempt to give a full dose of physic, but administer drachm doses of aloes daily till their bowels are moved; from the low diet they are placed upon during their voyage out, they land in poor condition, and I have known superpurgation produced on large horses from a four drachm dose of aloes.—ED.]

WARRANTING.

Caveat Emptor, at p. 253, says, "It is known that horses have secret maladies, which cannot be discovered by the usual trials and inspections; therefore the buyer requires a Warranty of Soundness, to guard against such latent defects." I have taken the liberty of extracting a great deal of the language, that follows, down to the line at p. 255, finishing, with the words "knowledge of the seller," from the above author, and converting the same to my own purpose; for which plagiarism I offer every apology. The arguments and recommendations, however, in various parts of "Caveat Emptor," in support of Warranty, are not exactly applicable to India, nor can I agree with them even for England, being opposed to warranty in every shape: so, with all due deference to the ability and the pleasantry displayed in the writings of that author, I am about to advise you to swamp all warranty, for these reasons :—

1st. If you sell a horse to-day that, either has a slight cold on him, or catches one during the time of sale, and that cold, from improper management or neglect, degenerates into a chronic cough, the purchaser may, perhaps, declare he had a chornic cough on him at the time of sale, and bring evidence to prove the horse coughed the minute he came out of your hands, and has coughed every day since: if

warranted, in law you *might* be liable, and have to take him back.

2nd. If you sell a horse, that should die two months afterwards of chronic diseased lungs, and a veterinary surgeon was to declare, from appearances on dissection, that the horse must have been diseased for a period of three months, and, consequently, must have been so at the time of selling: if warranted, in law you *might* be liable, and have to refund the money.

3rd. If you sell a horse that, four or five months previously, had put out his hip, strained his shoulder or back sinews, or had been lame from navicular disease; and a fortnight or so after purchase, he again puts out his hip, strains his shoulder, or back sinews, or becomes lame again from navicular disease: if warranted, in law you *might* be liable, and have to receive him back.

Caveat Emptor, p. 304, quoting from Lord Ellenborough says: "I have always held, and now hold, that a warranty of soundness is broken, if the animal at the time of sale had any infirmity upon him which rendered him less fit for present service."

1st. Again, if you sell a horse perfectly fresh and unblemished, and that horse, a week afterwards, throws out a spavin: if warranted, in law you *might* be liable, and have to receive him back.

2nd. If you sell a horse perfectly fresh and unblemished, and that horse, a month afterwards, becomes blind, from ophthalmia; and the purchaser proves that the sire and dam of that horse were blind from that cause, it is an hereditary disease: If warranted, in law you *might* be liable, and have to receive him back.

Caveat Emptor, p. 313. "Where, however, the proof of pedigree and hereditary disease are both accessible, it seems clear that a constitutional taint is unsoundness."

From the foregoing two sets of examples, with the quotations from law at the bottom of each, and which have been brought to bear on exactly similar questions, you may judge of the difficulties you might occasionally be placed in, as a seller, by warranting. I have used the word *might* throughout them all, nothing regarding horseflesh in law being positively certain, for so much depends on particular circumstances. "Unsoundness itself is sometimes sufficient "to break a warranty; at other times there must "have been knowledge of the unsoundness. Most "cases are questions for the jury, rather than of law. "No legal contract can be founded on fraud, and "wilful deception amounts in law to fraud." This is plain enough; yet, if you take your case to the law, the chances are always nearly equal, whether it will

be decided for or against you; "not from any defect in the law, but because both buyer and seller have always proofs of the shameful transaction."

Suppose you gain your cause: if you have been a seller, your horse may be returned to you half-ruined; and if you have been a purchaser, you are always bound to return a horse in as good a state as he was when taken from the seller's hands. Here is a second affair that may upset your first, and cost you another large sum. Avoid law, if possible, and never enter into any discussion: "your character, if you have any, will not be enhanced by embroiling yourself in a quarrel with a cheat;" but in order to prevent disputes, as well as litigation, never warrant, nor ask for a warranty. Do not commit yourself either, by saying, "He is sound as far as I know:" this is a qualified warranty, and the purchaser may maintain *assumpsit* upon it, "if he can show the horse was unsound to the knowledge of the seller." Such might be fair in some cases, but very unfair in others, and it might lead to great disputes; for every man who really knows a horse, must be fully aware there are not ten in every hundred that can strictly, professionally, and legally, be called sound. This, therefore, should be your only warranty:— "There's my horse, his price is——rupees, ready coin; you take him with all faults and diseases; I allow you a quarter of an hour's inspection, and I

will send him over when you send the money." There are even objections to allowing a man to try your horse. A friend of mine had a chesnut Arab for sale. A purchaser called to inspect him; he appeared to suit; was sound, wind and limb; fresh, unscarred, and four years old; price 1,200 rupees. "I like him much," said the purchaser; "might I throw my leg over him?" "Yes," said my friend, "ride up and down here as long as you wish, in my presence." He mounted; walked, trotted, and cantered; the action was good in every respect. You imagine, perhaps, the horse was sold. No, he now discovers two objections; he did not want a chesnut, he wanted a gray; and he did not like to go to so high a price as 1,200 rupees. The chap ought to have been forced to take him. I wonder how an imposition of this kind would be decided at law? If fifty heavy men were to play this trick on a slight blood Arab, his action, of course, would be none the better for it.

Having recommended you, as a seller, never to warrant, and consequently, in equal fairness, as a buyer, never to ask for a warranty, there is the greater reason in the latter case for you to proceed with caution, and if distrustful of your own judgment, to have a friend with you. When, therefore, a horse is brought out for inspection, if the appearance, figure limbs, &c.,do not satisfy, make your *congee*, but do not

abuse another man's property when at sale. If you are pleased, and fully certain that it is a horse you want; that the colour and price will suit, and that you have got the money ready to pay, take ten minutes' examination or allow your friend to do it for you, then solicit five minutes for a walk, trot, and canter; in a quarter of an hour let your decision be final; *if undisturbed* by the owner, this is ample. In England the case is different; there it is always advisable for an inexperienced person to have a new purchase submitted to a veterinary surgeon besides, for a couple of hours, a day, or two days, as he may think necessary. Half a guinea is all you have to pay, and this, with your own, or your friend's eyes, to boot, is abundance of warranty. Some persons, however, expect too much from a veterinary surgeon. A professional man can only tell you of any disease or remains of disease, or fault in the build, which is likely to produce disease or strain. He cannot tell you, merely by looking, if a horse is subject to gripes, rheumatism, or inflammation, unless some outward sign or symptom remain. He cannot tell you if a horse has ever been sprained, unless there is enlargement, mark as of blister, or something externally to denote it. He cannot say either if one horse is more liable to become blind, throw out a curb, spavin, or splent, than another, unless there is some visible sign or malformation, or he knows the sire and dam,

or grand-sire and dam, had these defects; and then he may say, "These diseases being often hereditary, or this build being faulty, they are more liable to occur in your horse." Beyond this, no uninspired veterinary surgeon can caution you. When a recruit presents himself before the surgeon of the regiment, on enlisting, do you suppose the surgeon could tell if he had fever last year, or sprained his leg last year, unless some evident weakness or enlargement remained? How can he tell if he is subject to gripes?

Buying a horse blind in both eyes, it is said you cannot return him as unsound. Caveat Emptor, p. 274. "But it has been held that a warranty against visible defects is bad in law, the purchaser being expected not only to possess ordinary skill, but to exhibit ordinary precaution." But a large splent extending on to the back sinews, a large spavin, large curb, or contracted, foundered foot, anchylosed pastern-joints, are all as visible defects as blind eyes: they are palpable defects, yet they constitute unsoundness: but "law is law."*

Much has been said against dealers in England, and dealers in India too. I have seen some black tricks in both countries; notwithstanding, I think, in the long run, dealers are as much sinned against

* The whole of this Chapter on warrantry is most judicious and correct.—ED.

as sinning. In either country, every man is bound to be wide awake, or, as the judge says, expected, not only to possess ordinary skill, but to exhibit ordinary precaution; and experience tells me to trust a dealer quite as soon as a gentleman.*

* Addison's definition of the word gentleman is "a term of complaisance, sometimes ironical." And gentlemen, and passers for gentlemen, are as often mystically mixed up together in one house, as thorough-breds and passers for thorough-breds are in the same staple.

At a dinner-party of eight, some few years ago, the conversation turning upon horse-flesh, I happened to let fall my ideas of the little general honesty existing in any part of the civilized world, in selling a horse. My *vis-á-vis* exclaimed, "Impossible! no gentleman would ever attempt to pass off an unsound horse." Five more of the party chimed in to this most creditable speech, so well calculated to delude the unwary, leaving only one of the same opinion, or rather, who acknowledged the same opinion, as myself. Clearly seeing that I had got into company with either knaves or fools, most probably a little of both, I thought more nourishment was to be gained at this house for the body by drinking, than for the mind by talking, so I allowed the subject to drop. Three months had not elasped when my *vis-á-vis*, the "impossible" gentleman, asked me to look at a horse of his that had been sprained, and blistered, but was still lame.

The horse had a ringbone on one pastern, besides something else internally wrong in the foot of the other, arising, most probably, from concussion; so I pronounced him incurable under eight months,—very probably, not then. "What shall I do?" said he, "Write on half a sheet of paper," I replied, "that the horse has been lame for two months—is thought to be incurable—and that he is in the market for sale." The man burst out laughing. He afterwards tried to get rid of him, but failed; for I took especial care that none of my acquaintance should be deceived by such an apostate as this. He then sent him to a dealer, who refused to sell him as sound. At last, he handed him over to a friend to dispose of, from whom he received nearly the full amount of his original cost. I did not discover the unlucky weight that was imposed upon; but, fancy this hypocrite crying out, "Impossible!" Here lies the honest distinction between a man wanting to buy, and one wanting to sell.

Every novice in horse-flesh is satisfied with his new hobby for a week. A horse, however badly bred, or faultily built, if only in good external condition, will always catch his eye before a thin one; and bog-spavins, thorough-pins, capped hocks, and windgalls, as well as the round shank bones and dents, are all less likely to be taken notice of when the nag is in plump order; many having a bone-spavin, a contracted long foot, or founder, have I seen pass through three or four hands, each new possessor alike unconscious of anything wrong: these treasures, whether latent or patent, not being always discovered till the horse is hunted, or suddenly becomes lame. Kind Griffins, then, for whom this volume is chiefly written, I most fully exonerate, and acquit you of the charge of intentionally deceiving; yet, for reflecting on your judgment, and for daring to assert, that if you want knowledge you must begin with "BLUNT SPURS," I know you would like to see me "regularly bitten." Friends, impose upon me with every fault and infirmity the horse is subject to; from the day this book issues from the press, I know I must be considered fair game; but if you have the slightest compassion for a man who has endeavoured to save your limbs, as well as your money, I implore you, never ask me to buy



A CHEAP HORSE.

SHARP SPURS HAVE FINISHED HIM.

Always on hand, for sale, at
MR. GREEN'S REPOSITORY.

	RUPEES.
A good horse...	400
A very good horse	500
A superior horse	600
A very superior horse...	700
A first-rate hunter	800
A very handsome horse..	900
A superb horse, carrying a magnificent head and tail...	1,000
A perfectly fresh officer's charger	1,100
A high-caste, five-year old horse, without any blemish	1,200
A racer	1,300
A handsome racer	1,400
A three-mile welterer	1,500

It was my original intention to have sent this Treatise forth with a brief exposition of every complaint and accident the horse is subject to, which I had divided into thirty-five classes, under a hundred and fifty heads, but at present I am prevented doing so :

at some future period these may possibly be added. On the request of an old acquaintance, not to fail to subjoin a few lines on red urine, and bursautee, I have dwelt a little on these two diseases, though out of their place.

I have now to acknowledge my thanks to Professors Spooner and Morton, of the Royal Veterinary College, for their very great kindness in perusing the foregoing before publication; Mr. Morton has, also, most obligingly undertaken the trouble of seeing the whole through the press for me; and if there are any little points not in strict accordance with the views of these scientific gentlemen, they are, perhaps, of no great importance. I can, however, assure you my manuscript received due praise, especially for the forcible expounding and illustration of the Foot and Heel; and I therefore am entitled to expect the work will soon realize me half a fortune; in order to succeed in which, I am about to change half my name. In conclusion then, critical gentlemen, I respectfully caution you to "Ware Name,"* and if you do not liberally patronize these pages, drawn out by the head of BLUNT SPURS, you may yet get punished with a heel, and be brought to the ground by

SHARP SPURS.

* "Ware horse," I suppose, you know, is the caution given in England to the by-standers, when a horse is brought out for sale.

BURMAH PONIES.

By "Vagrant."

"The Burman horses are small but spirited and "strong. There was one in 1842 in the menagerie "belonging to the Zoological Society of London. It "did not stand more than twelve hands high, but "he was a beautiful little fellow and a picture of "strength."—*Youatt on the Horse.*

"The horses of Burmah and the Shan country "which are imported into the Provinces, are small "ponies resembling the little Spanish horses that "run wild in Missouri, and the other western parts "of America."—*Mason's Burmah.*

The admirable ponies above described by Youatt and Mason were until the first Burmese war, almost unknown in India, but of late years they have won for themselves such high credit as safe and powerful hacks for mountain work, and so hardy withal that they are almost indispensable to all sojourners on the Neilgherries or in the Malabar Provinces, and for all the general uses that ponies can be put to, they are even greater favorites than the Mahratta galloways and tattoos once so common.


The sudden introduction of "Arab" and Indian blood into Burmah since the second war has however caused the "Burmah," or to use a more correct term, the "Shan" pony to differ now very widely from the animal described by Youatt and Mason.

Although perhaps the animal is now hardly so handsome or sturdy as formerly, the change is in most material points for the better, and it is difficult to conceive a more perfect hack for the purposes so small a creature can be turned to than the "Shan" pony of the present day.

The Shan Pony has doubtless sprung from the same origin as the mountain ponies of Upper India or the Tartar States, viz., the wild horse of Northern Asia. Blyth, the famed naturalist pointed out to the present writer that many of them still bear the Zebra-like, or, as he termed it, Hippotigris stripes. The breed has evidently been carefully looked to by the Shans, who resemble in industry and intelligence their cousins the Chinese, more than do any other of the Mongolian races, and who have hitherto been so jealously afraid of losing it that no stallion or mare, unless she was supposed to be barren, was to be found in a lot brought down for sale in the British provinces.

The "Shan" (or, as he used to be called the "Pegu") pony of a few years ago was seldom above 12-2, but of immense power in loins and shoulders, the latter,

and the neck, perhaps a little too heavy and the pasterns too short to please every eye, but as the animal was hardly ever required to go out of an amble under a heavy weight, these could not be considered such draw-backs as they would have been in hacks for lighter work, and although his neck was sometimes thick, his head was generally beautifully set-on, his muzzle and eye showed almost as much breeding as do those points in an "Arab," and his feet were naturally so perfect, that they could stand almost all the ill-usage that even an Indian "Nalbund" could inflict upon them. His pace was almost always an amble, out of which it was difficult to make him break.



Men formerly had quaint ideas regarding these ponies which were generally supposed to be proof against any unfair treatment, hardly ever getting sick, but when they did, to be incurable, and they were also believed to be invariably hard pullers and to require bits of curious forms to keep them within bounds.

In the narrative of the mission to the Court of Ava in 1855, by Captain Henry Yule, Bengal Engineers, this belief is thus referred to.

"* * * * As we had no opportunity of seeing a "Burman review, I borrow the description of one "from Col. Burney's Journal (January 2nd, 1831.)" "* * * * * * The business of the day commenced "by a party consisting of some of the Woongyis "and Atwenwoons and first Officers of Government, "dressed in their military helmets and uniform "appearing at one end of the open space, mounted "on small Burman ponies. They ambled one after "another past the king towards a target elevated "on a pole about twenty-five feet high, at which "they threw a small light lance, five feet long, as "they turned their horses round to return to the "spot whence they started and to take another lance "from one of their attendants, and amble again "towards the target as before."


"They each threw about a dozen lances, and the

" target was not struck more than ten or eleven times.
" .....When their stock of lances was exhausted,
" they dismounted, and the most successful of them
" came and kneeled down before the king, whilst the
" queen sent to them a piece of red muslin and some
" flowers. * * * * * The next display consisted of the
" Cavalry marching past the king, and afterwards
" trying, two at a time, to pick up an orange with
" their lances whilst in full gallop. Only two oranges
" were so picked up, one by a mahomedan who com-
" manded a portion of His Majesty's Cavalry, and
" who is said formerly to have served in Skinner's
" horse."

" The Cavalry consisted of three bodies, of about
" eighty or ninety men each. One party had curious
" high head-dresses, said to be the old uniform of
" the Cassay horse. Most of them appeared to be
" Muninpoorians or their descendants. They were
" armed only with lances, having neither sword nor
" carbines, nor pistols. I should have taken them
" to be very bad riders, but the manner in which
" they struck down upon the oranges whilst in full
" gallop, and recovered their lances, proved them
" to be good horsemen. The Burmese always ride
" without shoes, and can insert only two or three of
" their toes in their stirrup iron; the large gilt flaps
" of the saddle prevent the use of spurs, but the noise

"of these flaps, with which they urge their horses, "can never, of course, prove so effective as a good "pair of spurs. In riding, they bend their bodies "far back, and seem to hold on by the bridle, which "must be the cause of all Burman ponies having "such hard mouths. The legs of the rider also, with "the knees much raised, are in constant motion, and "do not appear to belong to him. Yet, from the "Burmese saddle having a high peak, both behind "and before, it is difficult to be thrown out of it."

The "Shan" pony has of late years increased an inch, if not two in height, is a lighter and more active, although hardly a less powerful animal very often displaying most unmistakable signs of the Indian "Arab" or "Persian" blood that has from time to time been left in Burmah.



Fortunately "Arabs" and "Persians," for the simple reason that more sires of those breeds have been left behind (as their masters obeyed the long hoped-for order of "relief") have done more than "Country breds," towards altering the type of pony; but traces of the wiry little "Mahratta" the varmint half English looking "Country-bred" from Calcutta, and we write it with sorrow the small eye and shut nostril of some of our worst Indian breeds are now often to be seen. It is interesting to note these signs: where traces of the two last-mentioned are found, the cross has generally come from the dam, and, as in most such cases, the produce is larger than it would have been with a full-sized sire, and a pony dam. Many of these ponies however have with their mother's merits inherited some of the irritability or timidity so often found in "Country-breds," and also occasionally a large, and, it may be a weak foot instead of the small hollow goat-like one of the real "Shan." Exactly the reverse is the case where the "Arab," "Persian" or "Mahratta" blood can be traced. There the cross has, by being from the small dam, apparently not so much increased in height as in lightness, activity, beauty, and docility, (this last virtue however the "Mahratta" has done little towards adding to) the animal will still work under as heavy weights, but he will now jump and gallop as well as any "Deccanee," and, instead of

pulling, the chief difficulty with ponies fresh out of a "Shan" lot is to get a bit light enough for them. Their feet are so perfect, that unless they are required for harness work over the stony roads of Rangoon, they need never be shod, and it is almost impossible to bring them down, unless they are very young and so over-ridden as to throw out splints, which, while forming, will always cause a horse to stumble.

It is as well to mention however that while, unless for harness a Shan pony with good feet need never be shod in Burmah, nor need he be in most up-country Indian stations, he should always be ridden with a little extra care during the first fortnight or so of the rains, as his feet have been worn down by six months' work over dry roads, and while the damp perhaps makes the hoof more tender, the stones have had their usual coating of dust washed off, and are more likely than ever to do harm.

This extra care need not however last more than a month, and the little additional thought required to make a man ride on the soft side of a road during that time is well paid for.

Perhaps the greatest drawback the Burmah pony has had to contend against in his struggle for favor is the notion, that although he seldom gets ill, when he does go wrong, he is more difficult to cure than any other horse. Hearsay has begotten, and indo-

lence has fostered this strange notion; indeed, this very chapter had its origin in the request of a Connoisseur in these ponies who implored the writer to say why they would not recover when once they got ill?

The reason is, not that a Burmah pony, whether right or wrong, requires treatment different from his stable companions, but, the master believes the animal to be thoroughly hardened against disease or fatigue, and takes it out of him or neglects him accordingly.

Over exertion, when not quite fit to work, puts the little animal "off his feed," or makes him listless or perhaps worse for a day or two.

This would be at once discovered and its evil effects guarded against were a *horse* out of sorts; but the groom, like the master, believes that a pony should never be ill and so neglects the first warning. The simple loss of appetite, or what the master would in his own case term "dyspepsia," grows into colic, and even this is not, because the victim is a pony, thought seriously of until it has turned into inflammation. The master even now does not like to believe that his tough little hack can have much the matter with him, and stoutly asserts that, "he will be all right "to-morrow." By to-morrow, however, the most severe treatment would be of little avail, and ere

many hours, there is ready for circulation another wonderful example of how impossible it is to cure a sick Pegu pony.

After long experience of Burmah and its ponies, the writer can very confidently affirm that the rules for their treatment, whether sick or well, are those laid down for other horses in the Griffin's Aid-de-Camp.

As these ponies are very hardly dealt with and vilely mis-managed when they are first purchased, the following remarks may perhaps be interesting, if not of service, to residents in Burmah.

The writer who has been long in that country, is well aware of the difficulties every lover of horse-flesh has to encounter there: but he is not the less able to affirm, that, with a little care and system, condition may be just as well kept up, and that whether with horses or ponies the coats and eyes may be as blooming and bright in Rangoon as they should be in a good stable anywhere in India.

The ponies are, at the end of the monsoon, after having been entirely grass-fed, and many of them hardly ever mounted, brought down in hundreds from the Shan States, N. E. of British Burmah, by perhaps the most original and most amusing horse-dealers in the world.

Although perhaps more industrious with his hands than any other Eastern (Chinese and Japanese of

course excepted) the Shan is as lazy in all other respects as is the Burman.

An intending purchaser looking over a lot must do every thing for himself and may pull the animals about as he pleases; but although his proceedings are regarded with indolent curiosity by the owners, it is almost impossible to get one of them or even their servants to show an animal off or to take an interest in his sale. A likely looking pony may strike the buyer's eye, but not one of the stout good-tempered-looking lads lounging in the wooden caravansary, round which the "lots" are tied as close as they can stand, will be at the trouble of moving the animal out, much less of mounting him, and most startling to a nervous purchaser would be the reasons given. Rather than exert himself, the Shan will without hesitation cry down his pony, "he is afraid "to amount him." "The pony is very violent and "will throw the rider off—will run away—will fall "down," &c., &c.

The only way is for the buyer to pull the animal out and to try it as he pleases. These ponies are, with hardly any exception, very gentle, and although they will at first very naturally wince and shrink from the saddle and bridle of civilisation, they may, as has been scores of times proved by the writer, be mounted bare-backed and perfectly managed with the halter by any

ordinary rider; they must be sound, for they have neither been shod nor used; hereditary disease must be almost unknown amongst them, and they are far too active and clever to meet with many accidents during their colt-hood. A purchaser, therefore, although he may pass over a really good pony, can hardly hit on a bad one. Let him, however, unless he has made up his mind to buy at once and take his new purchase away with him, beware, how he shows any preference, for if he be supposed to be a judge, eagerly watching his movements are a lot of most ill-favourd Natives of India, owners of hack carriages, or purchasers for the Madras market, who knowing nothing of horses themselves are prepared to go a few rupees in advance of his offer so that if he commence a bargain in the morning, he is very apt on returning to finish it in the afternoon, to find his animal gone.

These difficulties got over however, we will suppose the purchase concluded, and, that, for from one hundred and fifty to two hundred rupees, a stout gentle and handsome little hack, worth at least twice those sums in India is carried off by the exulting buyer. Now the poor little animal's difficulties and mismanagement commence. His temper and digestion are at once violently assailed, and when after a few months of constant seige one or both give way, the master blindly ignoring cause and effect, wonders

what ails the animal, and is apt to crave sympathy for his bad luck.

Up to the time of purchase the pony has been fed on a short allowance of green grass. A *short* allowance because his former Shan owner was too lazy to cut a full feed; he is at once allowed to eat as much as he can gorge of the miserable grass brought into Rangoon during the dry season, and besides this, is generally over-fed on chenna and paddy. The consequence is that sooner or later according to the stoutness of his digestion and in these matters horses are as delicate and variable as their masters, the pony goes wrong, gets "hide bound," "lampas" and uncertain in appetite, all the more so that he is very probably like many grass-fed animals afflicted with worms. The monsoon commences, rich and rank grass springs up everywhere—the pony suddenly gets as much of it as he can eat, his gram-feeding is still kept up, and the consequence is that if he is not carried off by an attack of inflammation, his skin is affected and that mange or still more often that abominable disease "Prurigo" and occasionally Ringworm, so common amongst the Burmans themselves, and now and then warts, swollen legs, and other symptoms of thorough want of condition show themselves. Occasionally even in gentlemen's stables although it is generally confined to those of the hack

carriage proprietors a disease much too like "farcy" to be bearable shows itself. So much for mismanagement in feeding, now for the attacks on the temper of the pony, after he was purchased.

Up to this date he has never been shod and his model feet are at once cut up, and cut out of shape by one of the wretched Nalbunds of Rangoon. The shoes are for a *harness* pony on the Rangoon roads, necessary evils, but the vile system of shoeing there cannot but annoy the pony and perhaps cause him to blemish himself for life by cutting or brushing. Until purchased, he has never had any bit in his mouth, except a rope or a common Burmese Snaffle, his master, at once, furnishes him with a severely bitted double bridle, tightly curbed,—the pony naturally resists this and the odds are that the dread of having his mouth tortured by a violent wrench at starting makes him fidgetty to mount and fretful when moving.

The owner and his friends are not surprized at this, for "Pegu ponies almost always pull"—and the evil is met by a still more powerful bit which ere many weeks, if hauled at with average strength and clumsiness effectually confirms their opinion and so deadens the once tender mouth that the pony can afford by hanging on the rider's hand and giving him the whole of its head to carry to furnish in the

most orthodox fashion another proof of how sure a Burmah pony is to pull.

The writer hopes that the following hints may be of use to some of his old friends in Burmah.

In the first place he ventures to advise every one fond of horses in that country to stack, as soon as possible after the rains cease, as much hay and straw as will keep their animals for at least eight months. There cannot be much risk in doing this, for, if not required for home consumption, the fodder will always sell and grass-cutters can be dispensed with.

The Burmah grass makes excellent hay, which, as horse owners are now beginning to appreciate will each season be more easy to procure. A Burman never cuts his straw but leaves it after the ears have been taken off, to be trampled down by his buffaloes, and eventually to be set fire to as a lazy and therefore a thoroughly Burmese way of manuring for next year's crop.

If this straw be cut as soon as possible after the ears have been removed and while the rich grass which springs up with it is still in flower, it makes when mixed with ordinary hay, fodder which all horses will greedily eat and thrive on well. Any Burman farmer, if fairly spoken to, will allow his straw to be removed, but, as the villagers themselves are too indolent to work, there may be a little diffi-

culty in getting Indian coolies to go out into the country to cut it. It should be stacked upon a platform raised as are the Burmese houses, and while both hay and straw are being pressed down, common black salt or salt and water should be freely sprinkled over the forage. The stack should of course be securely thatched over, and on the side the monsoon beats. So much for preparation to receive a horse or pony.

When first purchased which would probably be in January, a fresh pony should be sparingly fed—half of his allowance of grass may, if the owner pleases, or doubts whether he has stacked forage sufficient to last through the monsoon, be that daily sold in the Rangoon market, but it is better to avoid this if possible. Chenna and paddy bruised and mixed in proportions of one of the former to two of the latter should be given, but sparingly at first and increased as the animal's digestion and appetite may guide. Indeed it is a safe rule with fresh horses of all breeds, and, the same rules apply with equal force to Burmah ponies as to Australians, Capes, or Arabs, always to be underfed at first and increased by slow and small degrees. When condition is obtained however, keep it up on as high or as liberal feeding as the animal's constitution and the exercise he gets will admit of.

The above precaution alone should suffice to bring

an animal through his first introduction to work, into sufficiently good order to enable him to stand almost any hard usage he may be called upon to encounter afterwards.

As the rains set in, the hay should be slightly increased or, if there is not enough of it to feed entirely on, a very little of the rank grass now springing up may be given, but always dried as much as possible, and the gram rations may perhaps be slightly added to. Grooming must now, more than at any other season, be insisted upon for nowhere in India is it so difficult to keep up the action of the skin as it is in Burmah at the commencement of the monsoon. Should the animals coat get wrong a change of diet for a few days from bruised to boiled gram will often, at once prevent further mischief. "Cooltie" is not often to be had in Rangoon but "Oorrud" can always be obtained and makes an excellent change as does boiled instead of bruised paddy. The writer is convinced that with horses as with men, half the cases of irritability of skin if not indeed of temper, a Doctor has to deal with spring from damaged digestion: should this change however not be sufficient and symptoms of mange or "Prurigo" appear, although the feeding on boiled gram should still be kept up, a few of the seeds of the "Butea Frondosa" or "Palas Papara" *vide* the list at the end of this work will do wonders internally, while a little Kero-

sine oil rubbed over the irritable places, will, although it may for the time being, act almost as a slight blister, allay and remove all propensity to rub. Few men know the value of the "Palas Papara" seed, whether as a means of putting a horse into condition or as a vermifuge. It should however always be given when the stomach is nearly empty, say before morning grooming; six seeds or so, pounded down and given daily either with a very little gram or "Goor" *i.e.*, coarse country sugar, for a week or ten days—then discontinued for about the same period, and if necessary recommenced with, again and again, will generally brighten the eye and coat better and more safely than more remedies. It is complicated known to every horse dealer in India, can be had for a few farthings in every bazaar, and every servant knows it by its native name.

The seeds are thin, flat and kidney-shaped.

In his book on "The useful plants of India" Major Heber Drury mentions this tree under the following name:—

BUTEA FRONDOSA. *(Roxb.)* Nat. Ord. Leguminosæ.

Bastard Teak, Eng.	Palasie, Malayalum.
Porasum, Tamil.	Palas, Hindoostanee.
Moduga, Teloogoo.	Palas, Dhak, Beng.

and says that when in flower it has a very striking appearance from the gaudy appearance of the bright

scarlet blossoms—that "the name of Plassy so cele-
"brated in Indian History is nothing more than
"Palas or Palasie, the Hindoostanee name for this
"beautiful tree." Also that the gum is used in medicine as a powerful astringent administered in the form of tincture and powder, and also that two or three seeds deprived of their outer covering are frequently given in cases of tape-worm.

The next disease fortunately is little known except amongst the ponies of hack carriage proprietors but, as it sometimes occurs in stables which should be better managed, it may as well be mentioned here, although in doing so the writer frankly owns that he knows neither the cause, the name, nor how it should be dealt with. It is caused, however, he presumes by injudicious feeding and bad grooming in a climate under which it is very difficult indeed to keep the action of the skin up to the mark. If it is not "farcy" its symptoms resemble that disease so strongly that, were a pony of the writer's attacked with it the brute should immediately be shot to keep him from contaminating others. For the symptoms, *vide* those of "farcy" in any book on horses. It is treated and apparently with success by the hack carriage men, by the barbarous firing, natives delight in, over and all about the sores. Of the three cases in a gentleman's stable the writer has met with, the first died within a few days. Another on which his owner

tried a constant change of remedies without following any particular one for a week together, and would not destroy, became an object too loathsome to keep and probably died. The third was under the writer's charge at first—and was thus dealt with—the prescriptions having been got from a Veterinary Surgeon.

Arsenic, three grains
Cantharides, two grains

given in morning and evening feed, the sores well bathed three times a day with warm water and a solution of sulphate of copper afterwards applied. Liberal and varied diet was recommended but as that was difficult to obtain in Rangoon, the pony had in addition to his usual feed of grass, strong beef broth mixed with boiled rice every morning, as many carrots as could be procured and boiled Oorrud in place of the usual Burmese feed of chenna and paddy. This was continued for ten days, at the end of which the animal's master returned and startled at the liberal ration of beef and carrots his pony was receiving, took the advice of a native and had the animal fired after their fashion. The pony although blemished for life certainly did recover and is still working in Rangoon. Whether the cure was caused by the medicine and high feeding or the firing iron, or both combined, the present writer does not venture to say.

He now winds up this chapter by begging those who may take the trouble of reading it to believe, that Burmah ponies should be treated just as are other horses, that they do not, until their mouths have been spoiled by the absurd notion that they will pull, require bits unlike those of other hackneys, and lastly, that when a man is fortunate enough to have a good Burmah pony the little hack is well worth taking care of.

The names of some ponies well known at Rangoon in 1865 and 1866, may serve as illustrations of each stamp of animal mentioned at page 269.

Colonel Phayre's, "Garibaldi," "Dowager," "Lottery," "Chance Shot," "Earth Stopper," "Exchange," "Chowsah," "Sunshine," "Punch," and "Shandy Gaff."

"Dowager" and "Lottery," two of the tallest, ponies, (13 hands and 1 inch at least,) known up to that date in Rangoon, showed unmistakable signs of Indian "Country-bred" relationship, so did "Chance Shot," but in a less marked degree:—while the blue blood of the Desert must have contributed to give large eyes, open nostrils, broad brows, lean heads, hollow jowls and thin skins to the others. "Shandy Gaff" excepted, *he* was wonderfully like the old Mahratta tattoo once so well known too and loved by every true sportsman in the Deccan. Even to his

temper each of these ponies carried out the characteristics of the race from which he must have sprung.

The "Deccanee tattoo,"!!! how many a liver-trying ride under a roasting sun and in the teeth of wind which felt almost as scorching as a blast from a furnace: but still how many days of glorious sport and how many a cheery comrade do those words recall? No man, save the writer's old friend, Henry Shakespear, in the "Wild sports of India," has attempted to do justice to the game little slave to whose blood and gallant endurance as a hack, so many a day's sport, whether with gun or spear, has been due. We can vividly remember after many a broiling gallop, how welcome to the sight of weather-beaten rider and panting hackney used to be the varmint little "relief" as he stood under the scant shade of some "bauble" thorn. How alike in all essential points were the two ponies, how wide-awake and vicious-looking was the fresh one, and how often the little brute used to fight, until like Don Juan's hack, he

"Knew that he had a rider on his back."

When he would give up further mutiny, settle down to a stretching gallop and appear to enjoy it too; How we oft times used to reproach ourselves as we looked at the last pony, and how varmint and game were his looks as with big blood-like, but gentle eyes, the only *soft* things about a Mahratta, wide thin nostrils and lean head all brought out by

violent exertion; we pulled him up with quivering legs, heaving flanks and shaking tail, to bear witness to having been rattled along at twelve miles an hour over a bad road and under a midday Deccan sun. Never, their tempers excepted, were there better ponies, and sin and shame will it be if the breed, used up as it was for baggage animals during the mutiny by Rose's, Whitlock's and other Central Indian columns, not to mention the enemy, be allowed to die out. It is said however that Sir Richard Temple will prevent this, and if he does, the thanks of many a light weight with a lighter purse, should be freely bestowed on him.

Few, even among sportsmen, understand or appreciate the pleasure of a long ride of, say sixty, or even half as many more, miles at a stretch, with a change of horses at every stage. First, there are always the pleasures of change, then difficulties, and sometimes excitement or danger to encounter, and, where did man show greater knowledge of his fellows than did Scott when he wrote, as causes why they should stray.

* * * "I sought to drive away
"The lazy hours of peaceful day;
"Slight cause will then suffice to guide
"A Knight's free foot-steps far and wide.
"A falcon flown, a Grey hound strayed,
"The merry glance of mountain maid:
"Or if a path be dangerous known,
"The dangers' self is lure alone."

ADDENDUM.

HINTS ON THE EDUCATION OF A HORSE, COMPILED FROM THE WORKS OF VARIOUS AUTHORS.

(By the late Major Jenkins, Madras Army.)

Introduction.

A Horse in India when purchased from a Dealer's "Lot," has in almost every instance been backed, and probably has been ridden for years. Many persons are satisfied, provided the horse will go quietly along a road, and think that putting him through a course of instruction in a Riding School, is not only unnecessary, but actually injurious. There are others, however, who can appreciate a horse that has been taught to be "generally obedient;—light in hand;—to carry himself well;—to walk, trot, steadily and quietly, and always in hand;—to rein back freely, and close steadily to either hand;—to canter to both hands, and change leg;—to go about on the forehand, and haunches," &c.—but who may not know, how to teach a horse the accomplishments they desire, and it is for their use, that I have thrown together these *hints*, extracted from various authors; but principally from "Adams on Horsemanship," "Horsemanship by Captain Richardson," and "Training of Cavalry Remount Horses" by Captain Nolan.

By Monsieur Baucher's *Méthode*, recommended by Captain Nolan,* "The Horse is gently used, the progress is gradual, but certain. For a few days he is ridden on a snaffle, with a loose rein at a walk and a trot. He is then bitted, and a few simple lessons teaches him, to yield to the feeling of the rein, and the pressure of the leg,—then he is collected and got in hand, not by pulling and sawing at the horse's mouth, but by gradual pressing him with the leg, till he raises himself off the bit, and gathers himself up at a walk, then he can be controlled, and put together, to any extent required, by the judicious use of the spur. As all this is done at a halt, or at a walk, the horse undergoes no fatigue. Reining back, then perfects the horse in the use of his limbs, and in unqualified obedience to the rider's hand and leg. This once attained, a few lessons will teach the animal to canter, change leg, passage, and pirouette; the horse becomes a perfect charger in a very short time, without having, in any way suffered from his breaking, indeed! without having been over-tired, or over-worked, during the whole of his education, and from his mouth having been gently dealt with, it remains fresh and good, instead of being hard and callous."

The instructor should possess, activity and firmness, the greatest possible coolness, good temper, and

patience; he must gain the confidence of his horse; unless he does this, his work is one of brute force, and opposed to all the instincts of an intelligent animal. Horses are by degrees made obedient, through the hope of recompense, as well as from the fear of punishment; mere force and the want of skill and judgment, tend to confirm vice and restiveness, punishment therefore should never be inflicted on a young horse, except for decided restiveness and downright vice; even in that case, so soon as the horse moves forward, he should be treated kindly.

Care should be taken not to over-fatigue the horse, nor to wish to obtain too much at once.

The Longe.

Many persons mistake the meaning, or intention of longeing a horse, and suppose it is only to tire, fatigue, and worry a horse into subjection, the design is to make him familiar, and accustom him to be handled.

It is not recommended to longe a young horse, particularly without a rider on his back, if it can be avoided, as he is apt to rush round the ring, and strain himself; the longe is however useful, and often necessary with refractory horses, but the use of it, should be made the *exception*, and not the rule.

The young horse, having a plain snaffle in his mouth, with side reins, fastened to a roller, to which

there is a crupper attached, and a cavesson on his head, the noseband of which should be three inches above the nostrils, and not fastened too tightly, is led into the school, (which it is desirable should be an enclosed space ;—as the colt will work kindlier, when enclosed by walls, than in an open space, where he would be striving to enlarge his ground, or disengage himself ;)—care should be taken to see that the crupper is properly fitted, and that the hair of the tail be not doubled up, or entangled with it, and that the reins are of a proper length : the colt's head should not be confined too much, nor drawn down too tightly ;—the inward reins in longeing, should be a few inches shorter than the outward reins ;—so as to allow the rider of the horse, if mounted, to see his inward nostril.

A skilful person, will, single-handed, longe a horse in many figures ; and by heading with a whip, change him without stopping, and longe him in a figure of 8 ;—a horse should only be longed at a walk, till he circles without force. At first you will require an assistant, who should lead the horse round the ring, while you keep in line with his forehand ; if you get before it, it will tend to check the pace, and if you get behind the shoulder, it will excite the horse to increase the pace, and cause him unnecessary alarm ; when the horse has become accustomed to the place,

the Assistant had better be dispensed with, as the fewer there are about a horse, the better.

In longeing to the left, the whip should be carried in the right hand; and in longeing to the right, it should be carried in the left hand—the arm extended, the lash trailing on the ground; so that the whip may be readily thrown towards the hind quarter of the horse to urge him forward; but, the proper place to strike a horse is on the belly: study to keep the colt at a strictly regular pace, first at a walk, and then at a trot. When the colt has been longed to the one hand, for a sufficient time, draw him quietly to the centre, speaking to him and caress him; have in your hand a little sugar, or gram, to give him, but keep the whip out of sight, or let it fall at once upon the ground. Alter the length of the side reins, and put the horse round to the other hand. Do not let a lesson exceed half an hour.

When the horse has become tractable, you can teach him to change from the one hand, to the other, without a pause; the side reins in this case, being of course of equal lengths. In longeing to the left; when you wish to turn the horse to the right, pass the whip over into the left hand, retaining the cavesson as before; seize the tape at the length of the arm, with the right hand, and, giving the word "change," draw the horse round to the right, at the

same time throwing the whip in front of the horse, to check his forward movement; and cause him to turn, and pass round to the right; retain the whip in the left hand, and the tape in the right. To change from right to left; grasp the tape as before directed, at the length of the arm, with the left hand, and draw the horse round; at the same time, throw the right arm out to the right, with the whip extended; to check the horse, and make him pass round to your left.

Mounting.

Place a saddle on the horse, and with a plain snaffle in his mouth, and a cavesson on, lead him into the school, and walk him round a few times; then* "stand in front of the horse, raise the line with the right hand and play with it, speaking to the horse at the same time to engage his attention, whilst an assistant quietly mounts; no one else should be allowed near, as the more people there are round a horse, the more alarmed he is, and the more difficult to manage; as soon as the man has mounted, turn your back to the horse, and move on, leading him round, do not pull at the bridle nor look the horse in the face, but turn your back to the horse, and he will follow."

"In mounting a young horse, place the left hand rather high up on the mane, and with the right take

* Training of Cavalry Remount Horses by Captain Nolan.

hold of the pommel, *not* the cantle of the saddle, you can thus always swing yourself on to the horse's back; whereas, if your right hand is on the cantle, and the horse springs forward, or turns round, in trying to pass the leg over the horse, you must let go your hold with the right hand, and thus lose your balance and are thrown off. By the other way, you hold on with both hands, the saddle is open to receive you, and you can swing yourself into it in spite of anything the animal may do to prevent you."

The rider should have a whip in his hand but no spurs on,* "and his legs must not press the sides of the colt, if the colt begins to kick and plunge, you must keep a firm hold of the cavesson, and the rider must preserve his seat by balance alone;—there must not be the least violence, nor noise."

When the alarm of the colt has subsided, proceed to longe the colt with a rider on his back.

"In urging the colt to walk, the rider should bear slightly on the mouth with both hands, at the same time that the pressure is made with both legs. Avoid upon every occasion to allow him to amble, or break from a walk into a trot. If he starts, or shies at objects, caress him and take instant measures gently to quiet his alarm." † "The first point to gain is to

* Horsemanship by Captain Richardson.

† Training of Cavalry Remount Horses by Captain Nolan.

get the young horse to go forward and to go willingly, after which you can proceed to bend the horse's head first to the one hand, and then to the other. The rider should not play with the snaffle rein, but merely draw it gently to the side he wishes to bend the horse's head to, (of course always having a feeling with the outward rein;) and if the horse follows the indications of the rein only a few inches, his head should be brought straight again with the outward rein; this should be done first at a halt, and then at a walk.

To teach the horse to follow the indication of the rein, you should;—when given on the inward rein, draw the horse gently towards you with the cavesson, and ease off the line, when the indication is given on the outward rein; thus the rider can be assisted by a judicious use of the cavesson, in making the horse 'change,' and 'pirouette,' &c., so soon as the horse is teachable, the assistant and the cavesson, should be dispensed with.* "Begin to *collect* the horse both at a walk and at a trot, raise the horse's forehand, by keeping the snaffle in constant play, and do not allow the horse to lean upon your hand; practice the horse in changing from hand to hand, and in circles first large, and then in smaller ones; feel the horse up, with both reins at every step, and, if the horse is sluggish, make use of the whip upon the

* Training of Cavalry Remount Horses by Captain Nolan.

horse's shoulder, and work him well up to hand; but be particular, not to have a dead pull on the horse's mouth."

Mouthing and Bitting.

* "It must always be borne in mind," Captain Richardson remarks, "that the fineness of a horse's mouth, is not produced by lacerating the gums of the horse. The delicate, and beautiful skin which covers them, is never so tender and sensitive, after abrasion, as before. The mouth of the colt has to be formed to the usage of the hand; as the hand is the medium of the will of the rider, it is by very gentle and correct indications of the hands and of the legs, that the education of the horse is to be carried on, and perfected, not by ponderous bits and rough treatment; great care should therefore be taken in selecting a bit.† "The best for all purposes is a light one, with the cheeks of an average length, the mouth-piece merely sufficiently arched, to admit of the horse's tongue passing freely underneath it; the width should vary according to the breadth of the horse's mouth. A bit of this sort is quite sufficient to bring any horse under control, for it is a mistake to fancy, that the opposition a horse offers to the rider's hand, is caused by the peculiar shape of the mouth, or that one horse's mouth, is by nature, much

* Horsemanship by Captain Richardson.
† Training Cavalry Remount Horses.

more sensitive than another. The jaw-bone of every horse, is covered in the same way ; whether a horse be light or heavy in hand, cannot therefore depend upon the quantity of *flesh* between the bit and the jaw-bone, though many suppose this to be the case ; but the fact is, it is not the horse's mouth that is hard, but the rider's hand that is in fault." "If the horse has never been taught its yield to the pressure of the bit, and to bring his head in, he will, should he have a severe bit in his mouth, set his jaw against it, to alleviate the pain he suffers; and thus, he adopts a way of his own,which he will ever after recur to, in similar circumstances." The bit should be placed in the horse's mouth, so that the mouth-piece be one inch above the lower tusk, the curb chain laid flat, and fastened to that tightness, as to admit of one finger being passed, between it, and the horse's jaw.

Bending Lesson.

*"The balance of the horse's body, and his lightness in hand, depend on the proper carriage of the *head* and *neck*, and to these two points, first direct your attention. They should always precede, and prepare the horse by their attitude, for every movement about to be executed; and the rider has no power over the animal, until he has rendered both these points, susceptible of every impulse communi-

* Training of Cavalry Remount Horses by Captain Nolan.

cated by him. It stands to reason, that if they do not lead in all turns, and changes of hand, &c., if in any circling, they are not bent to the circle, if in reining back, the hand is not brought home, if their carriage is not always in keeping with the different paces, the horse may execute the movements required of him, or *not*, as he pleases; for his resources are still at his own disposal."

"A young horse generally attempts to resist the bit, either by bending his neck to one side, or the other, setting his jaw against it, carrying his nose high up, or low down, he must therefore, by bending him to the right and left, be taught to arch his neck, and to bring down his head on the reins being felt, and thus be rendered manageable. As a general rule, when the horse champs the bit, it is a sign that he no longer resists the action of the hand. It is of the utmost importance that the horse never be allowed to take the initiative." Always oppose the raising of the horse's head,—always lower your hands and bring it down. The horse being brought into the school, see that the bit is properly placed in the horse's mouth; and the curb chain so, that you can pass a finger under it; place yourself on the near side, in front of the horse's shoulder, facing inwards, the feet a little apart, to give you more power."

"Take the off bit rein in the full of the right hand,

close up with the ring of the bit, between the fore-finger and thumb, the near rein in the same way with the left hand. Thumb nails towards each other, and little fingers outwards, bring the right hand towards the body, extending the left one, from you at the same time, so as to turn the bit in the horse's mouth. The strength employed must be gradual, and proportioned to the resistance met with, taking care at first, not to bring the horse's nose too much in, or too close to the chest, which would make the bend very difficult; if the horse reins back, continue the pressure, until he finding it impossible to escape, from the restraint imposed upon him by the bit held thus cross-ways in his mouth, stands still, and yields to it."

"When the bend is completed, the horse will hold his head there without restraint, and champ the *bit*, then make much of him, and allow him to resume gently his natural position, but not to throw his head round hurriedly."

"Practice this in the same manner to the left. To prepare the horse to 'Rein in,' take the near snaffle-rein in the right hand, and the off rein in the left, at about six inches from the rings, cross them behind the horse's jaw, and draw them in, till the horse gives way to the pressure, and 'reins in' then make much of him."

You should now mount the horse, and bend the horse's head to the right and left.

"*To the right*, by passing the second finger of the right hand through the bit and snaffle-reins low down, so as to have the reins short on the off side; then draw them gently towards you, till you get the horse's head completely round to the right, in the same position as in the bend dismounted;—when the horse champs his bits, allow him to resume his natural position."

"When bending the horse's head *to the left*, pass the right hand over the left one, and placing the fore-finger through the near reins, proceed as before directed."

"Next teach the horse on the reins being felt, to arch his neck; and there remain steady, till he be allowed to get his head away again."

"To 'rein in' turn the little finger of the bridle hand towards the horse's head, lowering the hand as much as you can and keep it there; with the right, (nails down) take hold of the bit reins, close within the grasp of the left hand, and shorten them by degrees, drawing them through the left. When the horse resists much, and holds his nose up, keep the reins steady; do not shorten nor lengthen them; the legs closed, to prevent the horse from running back: he will remain perhaps a minute, or more, with his

nose up, and his jaw set against the bit, but will then yield, bring his nose in, and champ the bit; make much of him with the right hand, loosen the reins, and after a second or two, 'rein him in' again. This practice gives the horse confidence, for most young horses are afraid of the bit, and if frightened at first by a sudden jerk of the reins will never after go kindly up to hand."

"How to teach a Horse to obey the pressure of the Leg."

* "*To circle the horse to the right, on the forehand.* Apply the left leg well behind the girth, very quietly, and without touching the horse's side with the spur, press against him, till he takes a step to the right, with his hind legs; take the pressure of the leg, from him, and make much of him; then repeat the same, and get another step from him, and so on till he has turned about; the horse should not rein back; but his fore-legs remain steady, and his hind quarters, circle round the fore. At first you may feel the *left* rein, and use the whip very gently on the horse's side, near to where the leg is applied, but care must be taken never to hurry a horse; if done by degrees he will soon understand what he has to do."

"The leg opposite to the one which presses the 'hind quarters' to circle round the 'fore,' must be kept close to the horse, to assist in keeping him in

* Training of Cavalry Remount Horses by Captain Nolan.

his place, by communicating a forward impulse, whilst the other leg communicates the impulse, which makes a horse step from right to left, or left to right; and in order that the pressure of the one, shall not counteract the effect of the other, the leg applied to make the horse step to either hand, should be further behind the girth, than the leg used to keep him up to hand. Both legs should be close to the horse's sides at all times, the pressure on either side should be increased, as occasion requires. In circling to the other hand, *vice versâ*."

* "By circling the horse on the 'fore hand,' you have taught him, on applying the leg, to move his haunches to either hand, and as he has thus learnt to obey the leg, you can, by making use of it, prevent him from moving his hind legs, to the right or left, and can now teach him to circle on them."

To "circle on the haunches right about;"—bend the horse's head a little to the right with the bit; pass the right hand over to the near snaffle-rein, apply the left leg as far behind to girth as possible, to keep the haunches steady; make the horse step to the right, by feeling the outward snaffle-rein, and passing at the same time both hands a little to the right, keep the horse up to hand, with the inward leg.

At first the horse must be halted, and made much

* Training of Cavalry Remount Horses by Captain Nolan.

of, three or four times during each turn, and if his haunches are thrown out, they must be brought back again by applying the outward leg;—thus gradually led on, the horse will learn to go about to both hands, on his haunches; when the horse is broken, it will be sufficient to carry the bridle hand to the side, you wish to turn to."

The use of the Spur.

* "The *spur*, is not only used to inflict punishment, when a horse refuses to obey the pressure of the leg;—but, it is also, combined with a good hand, a most powerful agent in bringing into subjection the most intractable, and in infusing spirit in the most sluggish animals; it requires great prudence, and a thorough knowledge of the horse, to use the spur so as to obtain the proper results.

"The object is to unite the horse's powers at the centre of gravity, that is, between the *forehand*, and *haunches*; and it is by the combined use of the hand and leg, that we attain this.

"Suppose your horse at a walk bearing the weight of five lbs. on your bridle hand; when you close your legs to him, you will feel the effect of the impulse communicated in the additional weight thrown on your hand, and this weight augments, in proportion to the impulse given.

* Training of Cavalry Remount Horses by Captain Nolan.

"On feeling this additional weight on the bridle hand, do not give way to it, but keep the bit hand low and steady, and play with the right snaffle-rein; the horse finding the bit an insurmountable obstacle, will by degrees learn, instead of throwing his weight forward when the impulse is given by the hand, to throw it back, and bring his haunches under him; but should you, instead of closing the legs, or using the spur gently, put both spurs into his sides, the horse would throw so much weight forward, from the great impulse received, that he would probably pull the reins out of your hand; your object would thus be defeated in the beginning; and the horse having burst from control on the first application of the spur, by throwing his weight forward, would ever after do the same. It is of great importance therefore, that the spur be used with judgment, and a knowledge of the horse's temper, so that the impulse communicated to the horse with the spur, should not be stronger, than what you can easily control with the hand.

"Increase by degrees the use of the spur, until the horse will stand its application without throwing any weight on the hand, without increasing his pace, or without moving, if applied when standing still."

Reining Back.

* "Reining back should not be commenced until

* Training of Cavalry Remount Horses by Captain Nolan.

the horse is well bent, and obeys the pressure of the leg; during the reining back, the horse must be well in hand and well balanced; as he can then make an equal use of all his fore legs, and raise them equally from the ground.

"Having a steady feeling of both reins, apply both legs to the horse, to make him lift one of his hind legs; at the moment of his doing so, double the feeling of both reins, which will oblige the horse to recover his balance, by stepping backwards, and thus, the first motion in reining back, is produced; give the horse his head, and make much of him; at first a few steps backwards, is all that should be required from the horse, increasing by degrees; if he bring his hind legs too much under him, ease the hand, and apply both legs to make him regain his balance forward; for this reason always apply the leg first, and then an increased pressure on the reins, because if you feel the reins first, the horse throws his weight back, and it stands to reason," as Captain Nolan observes, "that the more weight he throws on his hind legs, the less able he is to lift them."

"The horse must never be allowed to hurry, or run back out of hand, nor diverge from the straight line; should he do so, place him straight, by bringing his haunches to the right or left as may be required."

Shoulders in.

The horse having learnt to follow the indications of the hand and obey the pressure of the leg, can now be taught the 'Shoulder in,' at first a few steps only being required of him.

Separate your reins, take one in each hand, and hold them rather short;—bring the inward hand down, letting the wrist rest just above the hip, draw the horse's head round, at the same time advance the outer hand, to admit of the horse freely bending, press him with the inward leg and he will move diagonally, the outward hand by light touches, animates the horse, and guides him, in the direction you wish him to go.

Passaging.

After a few lessons at the shoulders in, proceed to the half passage, and passage; that is, to teach the horse to move sideways. The motion of the horse's legs in passaging is the same as that in shoulders in, but the head is turned differently. In the passage, the horse looks the way he is going, and moves directly to either side.

The horse being well balanced, the feeling of the reins corresponding with the pressure of the legs,—supposing you are going to the right, lead the shoulders off first, by a double feeling on the right rein, at the same time, apply an increased pressure with the left

leg,—the horse will cross his legs, and move sideways to his right. After a few paces stop, and make much of him; should the horse not readily obey the leg, apply the whip gently on the horse's shoulder and flank.

Paces.

The Walk.—The walk is the least raised, the slowest, and the most gentle of all the paces, but it should be an animated quick step.

* "Before moving forward, the horse should be light in hand, the head brought home (and not with the nose stuck out) the neck arched, and he should stand evenly on all his legs."

"In urging the colt to a walk, close the legs and communicate a sufficient impulse to carry him forward, but do not ease the hand, because if you do, the head and neck may relapse into a position which may defy the control of the hand.

"You should always have a light feeling of both reins, and when the horse bores on the bit, keep the hand steady, use both legs, which, by bringing his haunches under him, will oblige the horse to take his weight off your hand."

† "Avoid upon every occasion to allow the horse

* Captain Nolan.

† Horsemanship by Captain Richardson.

to amble, or break from a walk into a trot; if he starts and shies at objects, caress him, and take instant measures gently to quiet his alarm."

* *Trot.*—A horse trots when he raises the "off fore and near hind leg," or "near fore and off hind," from the ground at the same time. Begin the trot in a very careful and quiet manner, and do not proceed to extend his pace, until he has become well collected in the slow trot.

"The hand must be constantly at work to retain the head and neck in their proper position, without counteracting the forward impulse communicated by the leg; thus the horse will acquire regularity of pace, increased speed, and that safety, which is natural to the horse that is well balanced and light in hand.

* *The Canter.*—"Is a repetition of bounds, during which the forehand raises first, and higher than the hind quarters.

"The horse being properly placed, light in hand and well balanced, throw the weight more on his haunches, by increasing the pressure of the legs, and restraining him with the reins, and according to the hand you wish to strike off to, throw the weight of the horse to the opposite side. He must not be

* Training of Cavalry Remount Horses by Captain Nolan.

allowed to canter with either fore-leg leading at his own will, but must be made to do so at the will of the rider.

To canter with the right or off fore-leg leading, the extra bearing must be made upon the right rein; and at the same time the pressure with the left leg must be increased. If the horse refuses to strike into the canter by the pressure of the leg alone, then use the spur instantly, or strike the horse with the whip on the left shoulder. If the horse still hesitates, pass the whip behind the back and strike the near hind quarter. To canter with the left or near fore-leg leading, make the extra bearing upon the left rein, with the pressure of the right leg or spur. Teach the horse to strike off on the circle first, than on the straight line after that, try him at changing leg.

"A horse may canter false, disunited with the fore, or disunited with the hind legs. If the horse in cantering to the right leads with the near fore, followed by the near hind, he is cantering false. If leading with the near fore-leg, the off hind leg remains further back, than the near one, he is disunited; in such cases, close both legs to the horse, to bring the haunches under him; feel both reins, but the hand you are leading to the stronger, to bend the horse's head slightly in that direction; throw the weight off the leg he is required to lead

with;—which will enable him to throw it forward *and* to canter as required.

To stop a Horse.

* "The horse, being accustomed at the pressure of the rider's legs, to bring his haunches under him, when you wish to stop, close both legs, and throw your body back, keep your hand low, and increase the bearing upon both reins, by drawing the arm back. If you do not use the legs, but merely pull at the bridle, you may by strength of arm, stop the horse; but it will be entirely on his forehand, and the horse would be under no control. As soon as the horse has obeyed the check, and remains still, ease the reins."

Leading.

In leading you should never be further forward than your horse's shoulder, with your right hand hold his head in front of you, by the bridle, close to the mouth, and if necessary touch him with the whip with the left hand.

Mode of constructing an open Circus or School.

Enclose a circle of a diameter of from 42 to 50 feet with thick planking, or a mud wall, 3½ feet high, with folding doors to be made to open inwards; and outside this, about three or four feet from the planking, have a mud wall built six feet high; so that the horse cannot see over it.

* Training of Cavalry Remount Horses by Captain Nolan.

Prepare the ground as follows :—

Dig up the whole of the earth within the circle, a foot in depth; hollow the centre about six inches, into the shape of a saucer, replace the earth with a few inches of good mould, free from stones, cover the whole with about six inches of straw, or old bedding, which has been well saturated, over the straw, place three inches of good mould, and have the whole well trodden down, so as to form a good foundation; the sides sloping upwards to the wall :—over all, put some cart loads of saw dust, or tan.

Longeing Whip.

The longeing whip should be 13½ feet long, the handle being 5 feet, and the thong 8½ feet made of hide, or leather split at the end.

Age and Work.

(From Mayhew's Horse-Doctor.)

"One horse could wear out two pair of legs," is an old jockey's phrase. Most men when purchasing a dumb slave, pay great attention to the lower extremities. If an animal be used up, or has performed hard-work, the indications are sure to be found on those parts; but what comment does the language the act referred to pass under the conduct of those masters, the history of whose treatment, or rather

whose abuse of a living creature is thus sought for and often found upon a breathing frame?

Before the strength has departed, or the legitimate number of years are exhausted, cruelty deprives a most obedient drudge of its power to serve. The history of almost every horse is a struggle to exist against human endeavours to deprive it of utility. Nature, when she made the animal, formed a creature hardly second to her master-piece in anatomical perfection; the legs are strong, but in his impatience and blind obedience to the dictates of fashion, man *will* put them to their fullest use before their structure is confirmed. Racers go into training, when one year old. Carriage-horses, machiners, cart-horses, nags, roadsters, may-birds and park-hacks, generally come into work about the third year. The animal, however, does not cut all its teeth, till the completion of its 5th birth-day. It is required to look upon eight seasons before its adult period is entered upon, and yet, at the 3rd year or before that period, it is put to such work as only a horse can or does perform.

When the horse was designed to be only matured, the frivolity of mankind pronounces the creature to be *aged*. The life is indeed, generally worthless before the 8th year is entered upon. The young flesh, bones and sinews, long before that time arrives are made the seats of poignant diseases. Work, not

in the first instance laborious, but sudden and energetic beyond what the frame of the *young* horse can endure, casts it out of the gentleman's stable. Once removed from that place, its descent is rapid. From the carriage to the hack is a leap often cleared in Equine history; but every change adds misery to its lot! It fares worse, lodges worse, and works harder with every new proprietor, till at length, as its years and wretchedness accumulates, Nature interposes and takes the sufferer to Herself.

Spavin, splint and ring-bone are no more the legitimate consequence of Equine existence, than noads and anchilosis, are the natural inheritances of human beings; yet, what would the world look like, if men had their motions impeded, and their joints firmly locked by bony deposits in anything like the proportion which such misfortunes are witnessed in the inferior life? The most useful, the most trusting, and the most joyous animal is one towards which man acts, as though his study was to abuse the authority entrusted to him. Its utility lies in its legs; its play also is a canter; but before its body is set, its limbs are disabled."

* * *

LIST OF MEDICINES, &c., GENERALLY USED IN INDIA, FOR VETERINARY PURPOSE.

English.	Hindoostanee.	Tamil.	Uses.
Aloes	Musambur	Kariapolum	Purgative 3 to 5 drachms. Alterative 1 drachm with 2 drachms of Gentian every morning, until the bowels are acted upon—1 drachm of Ginger Powder to each bolus.
Alum	Puttacarei	Paddi-karum	Equal parts of Alum and Calomel, is used in Cancer of the feet—3 drachms each of Alum and Catechue, is used for soft doing, when not arising from diseased liver—4 drachms of Alum in a Bottle of Water is used for sore mouth.
Ammoniac Salt... ...	Noasauger	Navacharum	With Nitre is highly recommended for removing inflammation in Sprains—the Bandage being continually kept wet.
Anise Seed	Sonf	Somboo..	In cordial preparations.
Antimony Sulphurated...	Surmah..	Unjunuckle	One part of Antimony, and two each of Nitre and Sulphur is used in Cutaneous disease—4 drachms of the powder daily in gram.

Arrow Root	Kooaka neshasteh ...	Kova mavoo	
Asafœtida..	Hing	Pérungyum	In drachm doses with 2 of Gentian or 1 of ginger daily, is very beneficial for Cold—in Flatulency, 1 drachm to a pound of water and a drachm of ginger, drench.
Bastard teak seed or Butea frondosa. ...	Palas papara	Porasum verray ...	Two drachms in powder to be given in gram daily, for Horses with worms.
Black Pepper	Kálee Meerchie ...	Molooghoo	2 drachms of Pepper with 4 of salt *(Common)* mixed in the Evening gram, is useful for Horses exposed to the open air in rain or cold season.
Bishop Weed, (Seed of)..	Ajiowan.	Womúm.	In cordial preparations and fumigation in Catarrh.
Bran (Wheat)	Guavm Ka boosa ...	Godumbay thowdoo ...	Instead of gram, a day or two preparatory for physic.
Bitumen Pretolum ...	Muttie ka tail... ...	Muntylum	In Mange dressing.
Bole Armenic	Ghilarmenie	Simie Kavakulloo ...	
Calomel	Paraka Caffoor. ...	Rassakapoorum ...	In Salvating and in Liver affections, also see Alum.
Camphor...	Kufoor...	Carpoorum	In Fever and for destroying worms in cancer.
Candle	Tsharbee kee buttee...	Moolookoo vuttey ...	

LIST OF MEDICINES, &c.—*(continued.)*

English.	Hindoostanee.	Tamil.	Uses.
Cardamom.	Elachie...	Yalersie..	In cordial preparations.
Carrot	Gajur	Carrot Kalung. ...	
Cassia lignea	Darchinie	Lawunga puttay ...	In cordial preparations.
Catechue...	Kutt	Kauskettee	See Alum.
Chalk	Veláitie Chunna ...	Semi Chúnámboo ...	With Tinct. of Opii for over-purging.
Chatties	Mattee ka bassan ...	Sattee	For keeping water, gram, &c.
Chilly, dry	Sookah Lalmaresh ...	Konjey Molacoy ...	Should be made into paste and applied on Mange parts and washed 4 hours afterwards.—See Black Pepper.
Cinnamon..	Kulmie Darchanie ...	Karaway puttay ...	In cordial preparations.
Clay, Potters	Chickni muttie. ...	Kali munnu	Equal parts of cowdung and clay, very useful in stuffing hoofs at night.
Cloth	Kupra	Pedavay.	For bandages.
Clove	Laong	Craumboo	In cordial preparations.
Copper	Tamba...	Samboo...	

Copper (Sulphate of) ...	Neelatoto	Tooreshoo	Externally as Caustic. Internally Tonic.
Coriander..	Cotumbaroo	Cottamullee	The young plants generally sold in vegetable markets are very useful when Horses are over-purged—from physic, give 5 or 6 handful.
Cotton	Ruhie	Panghee.	
Cow's dung	Gayeka Gobur.. ...	Pasavoo Shanee ...	In stopping feet.
Creyat	Crëatt	Kiriat or Nela-vembu.	A substitute for Gentian.
Croton	Jumal Gota	Nérvalum Cottay ...	Should be used by a Professional.
Cubebs	Dumke murchie ...	Val méllághoo.. ...	
Cummin Seed	Zera	Seragum	
Charcoal	Colssa	Addipoo Kurrey ...	
Egg	Anda	Moottay.	
Fennel Seed	Sonf	Perum Seerugum ...	In cordial preparation.
Fenugreek	Methie...	Vendium	In cordial preparation.
Flour of Wheat	Guavm-ka-ata.. ...	Godumbaymao ...	
Frankincense	Coondoor Oodh ...	Sambranie	

LIST OF MEDICINES, &c.—*(continued.)*

English.	Hindoostanee.	Tamil.	Uses.
Galls	Mahpull.	Mashakai	For Cancer and Thrush, take finely powdered galls, rub over the part with a spatula, dropping lime juice while rubbing.
Garlic	Lassoone	Vellipoondoo	When made into paste with two-thirds of its weight of Wheat Flour, forms a very powerful Blister which should be removed fifteen minutes after application, the part to be blistered, should be shaved.
Ghee	Ghee	Nayee	A quart bottle full milk-warm is given in Colic when nothing else can be procured.
Gingelly Oil	Meetah tale	Null-unnay	For preparing Blister.
Ginger, Green	Udruck...	Injee	In cordial preparation.
Ginger, dry	Sount	Chookoo	In Flatulency 2 drachms in powder thrown up in injection has proved beneficial.
Gram, Horse	Koolty...	Koelloo	When boiled is the common diet for Horses in the Northern, Southern and Ceded Districts, also in the Presidency.

Grass, Green	Gans	Pilloo	
Grass, Dry or (Hay) ...	Sookhah Gaus.. ...	Kaunjey Pilloo ...	
Great Millet..	Jarie	Cholum	Diet for Horses.
Green Gram	Moong	Pucha-payaroo.. ...	Do.
Gum Arabic	Kavet Ka Ghonde ...	Vullám Pissini... ...	4 to 6 ounces in a bucket of water is given to Horses suffering from inflammation of kidneys and bladder.
Hemp or Tow	Sanka djar	Shanapa-naar... ...	In pledgets for pressure to Cancer in the foot, and secured by bandage.
Hogs Lard	Suever Ka-tcherbee ...	Punnee Colopoo ...	For preparing Ointments.
Honey	Shahut...	Thayne	
Indigo	Neel	Neelum	Solution of Indigo is used for abrasions.
Iron, (Sulphate of) ...	Heera Cassish.. ...	Unna Baydie	As tonic.
Jaggery	Goore	Vellum	The syrup is used to keep boluses or masses moist, when a large quantity is made.
Leech	Jouk	Uttay	
Leather	Chumra	Thole	
Lemon or Lime... ...	Neemboo	Elimminchum pullem.	With Galls for Cancers.
*Liquorice Root... ...	Meeta lukree	Ademoodrum	In cordial preparations.
Lime, Quick	Choonah	Soonamboo	In solution it is used to correct fœtid discharge in Cancers.

LIST OF MEDICINES, &c.—*(continued.)*

English.	Hindoostanee.	Tamil.	Uses.
Linseed Oil	Ulsi Ka tale	Alliverray-unnay ...	A Bottle of Oil is given in inflammation of bowels.
Liquor, spirituous ...	Arrack...	Shoyroyum	In preparing Tincture, &c., when Alcohol cannot be procured.
Long Zedoary	Umbee huldee.. ...	Pulang kalangoo ...	
Mace	Jawatrie	Jadiputree	
Marking nut	Beelamboo	Sharang cottay ...	
Mercury	Parah	Rassum	
Milk	Dhoodh	Paal	
Musk	Meshk	Castorie	A scruple with two drachms of Gentian given daily to Horses suffering from Rheumatism or cold.
Mustard	Rai	Kadaghoo	Counter irritant for Rheumatism.
Myrrh Gum	Bol	Valitipollum pissini ...	Tincture is made of this Gum and applied to abrasions.
Neem Tree	Neem ka jaud... ...	Vaypum marum ...	
Nitre	Shorah....	Pottel oopoo	As Febrifuge and Diuretic.
Nutmeg	Djapul....	Jadecoy	In cordial preparations.

Naptha or Rock Oil ...	Muttie Ka-thale ...	Mun tylum	Externally with other Oils for mange.
Onion	Piaz	Vungoyum	A good cooling diet with Tyre and boiled Rice.
Opium	Afeeom	Abinee	
Pepper, black	Golemirsh	Mullaghoo	
Do. Cayene... ...	Lalmirsh	Mollagoy	Take an handful, make into paste with an ounce of Black Pepper, and apply to mange parts, wash after 4 hours and repeat after a week.
Radish	Moollee	Molinghe Kalungoo ...	
Rice	Schavul	Arisee	A substitute for gram when Horses are suffering from heat spots.—See Onions.
Rosin	Googole	Coongalingum ...	3 drachms as Diuretic with Nitre, same quantity.
Sarsaparilla, Country ...	Muckey	Nunnaru-vayr ...	
Senna	Soona Mukkee ...	Nillaverie	
Soap, Country	Sauboon	Soucarum	
Soap-nut...	Reetah	Poonga-cottay ...	A substitute for Soap.
Soda impure	Khar	Kharum	
Suet	Buckray-Ka-tcherbee.	Aatoo Kolupoo ...	For making Ointment, &c., with other ingredients.
Sugar	Shukker	Sukkaray	
Sugar Candy	Meesseree	Kulkundoo	

LIST OF MEDICINES, &c.—*(continued.)*

English.	Hindoostanee.	Tamil.	Uses.
Sugar Cane	Gunna	Kārumboo	
Sulphur	Genduk	Ghendagum	For mange with other ingredients both internally and externally.
Syrup	Shukkur Ka-shirap ...	Shukkaray pavoo ...	For keeping masses soft.
Sweet flag	Butch	Vassamboo	Ground well with water into a fine paste and rub well, for killing lice or fleas.
Tobacco	Thumbaukoo	Pooghie ellay... ...	For Flatulent colic, inject Tobacco smoke.
Turmeric	Huldie...	Mungil...	In cordial preparations.
Vinegar	Cirkah...	Kadi	Diluted for lotion in sprains.
Vitriol blue (Sulphate of Copper)	Neela totta	Toorushee	As Tonic in drachms doses daily, as Caustic either in powder in solution.
Vomica nux	Koochla	Yettie cottay	The nut when burnt (not to ashes) and powdered is useful in long standing Ulcers by covering it over.
Wax	Moom	Molakoo	
Zinc (Sulphate of) ...	Suffaid toota	Vullay tootum ...	Combined with Sugar of Lead, as lotion to sprains, &c., a drachm of the latter to a scruple of the former in a bucket of water.

THE VETERINARY PHARMACOPŒIA

FROM

"Forrest's Farrier and Horse Doctor."

THE following *formulæ for veterinary practice* have been compiled from the works of the most eminent veterinary writers of the present day, as Blaine, Clark, Laurence, Peel, White, etc.; and we can from our own experience also, confidently recommend the selection to the notice of agriculturists, and the owners of horses in general. It would be prudent for such as have many horses, and particularly for such as live at a distance from the assistance of an able veterinarian, to keep the more necessary articles by them in case of emergency: some vendors of horse drugs keep veterinary medicine chests: and where the compositions can be depended on, and the uncompounded drugs are genuine and good, one of these is a most convenient appendage to every stable.

The veterinary pharmacopœia for oxen, calves, and sheep has been included in the arrangement. Where any speciality occurs, or where distinct recipes are requisite, they have been carefully noticed; it will therefore only be necessary to be kept in mind, that with the exception of acrid substances, as mineral acids, etc., which no cattle can bear with equal impunity with the horse; the remedies prescribed require about the following proportions: A large ox will bear the proportions of a moderate sized horse; a moderate sized cow something less; a calf about a third of the quantity; and a sheep about a quarter, or at most a third of the proportions directed for a cow. It is also to be remarked, that the degrees in strength in the different recipes, are usually regulated by their numbers, the mildest standing first. Each Recipe is numbered and complete.

Alteratives.

1—Levigated Antimony, 2 drachms.
Cream of Tartar,
Flour of Sulphur, each ½ an ounce.

2—Cream of Tartar, Nitre, of each ½ an ounce.

3—Æthiop's mineral, Levigated Antimony,
Powdered Resin, each 3 drachms.

Give in a mash or in oats and bran, a little wetted, every night, or make into a ball with honey.

Tonic Alteratives.

1—Gentian, Aloes, Ginger,
Blue Vitriol, in powder, of each 1 drachm,
Oak Bark in powder, 6 drachms.

2—Winter's Bark in powder, 3 drachms,
Green Vitriol in powder, 1½ drachms,
Gentian in powder, 3 drachms.

Make either of these into a ball with honey, and give every morning.

3—White Vitriol, 1 drachm,
Ginger or Pimento, ground, 2 drachms,
Powdered Quassia ½ an ounce, Ale, 8 ounces.
Mix and give as a drink.

4—Arsenic 10 grains, Oatmeal 1 ounce.
Mix and give in a mash, or moisten oats nightly.

Astringent Mixture for Diarrhœa, Lax, or Scouring.

1—Powdered Ipecacuanha, 1 drachm,
Powdered Opium, ½ a drachm,
Prepared Chalk, 2 ounces, Boiled Starch, 1 pint.

2—Suet 4 ounces, boiled in Milk, 8 ounces,
Boiled Starch, 6 ounces, Powdered Alum, 1 drachm.

The following has been very strongly recommended in some cases, for the lax of horses and cattle.

3—Glauber's Salts, 2 ounces, Epsom Salts, 1 ounce,
Green Vitriol, 4 grains, Gruel, ½ a pint.

When the lax or scouring at all approaches to dysentery or molten grease, the following drink should be first given.

4—Castor Oil, 4 ounces,
Glauber's Salts, dissolved, 2 ounces,
Powdered Rhubarb, ½ a drachm,
Powdered Opium, 4 grains, Gruel, 1 pint.

Astringent Balls for Diabetes or Pissing Evil.

Catechu [Japan earth], ½ an ounce,
Alum, powdered, ½ a drachm,
Sugar of Lead, 10 grains,
Conserve of Roses, to make a ball.

Astringent Paste for Thrush, Foot-rot, Foul in the Foot, etc.

Prepared Calamine, Verdigris, of each ½ an ounce,
White Vitriol, Alum, of each ½ a drachm,
Tar, 3 ounces. Mix.

Astringent Washes for Cracks in the Heels, Wounds, etc.

1—Sugar of Lead, 2 drachms, White Vitriol, 1 drachm.
Strong infusion of Oak, or Elm Bark, 1 pint. Mix.

2—Green Vitriol, 1 drachm,
Infusion of Galls, half a pint.

Mix and wash the parts three times a day.

Powder for Cracks, etc.

3—Prepared Calamine, 1 ounce,
Fuller's Earth, powdered,
Pipe Clay, powdered, of each 2 ounces.

Mix and put within gauze, and dab the moist surfaces of the sores frequently.

Astringent Paste for Grease.

1—Prepared Calamine,
Tutty, powdered,
Charcoal, powdered, of each 2 ounces,
Yeast enough to make a paste.

2—To the above, if more strength be required, add of Alum and Verdigris each a drachm.

Astringent Wash for Grease.

3—Corrosive Sublimate, 2 drachms,
Spirit of Wine or Brandy, 1 ounce,
Soft Water, 10 ounces.

Rub the Sublimate in a mortar with the spirit till dissolved, then add the water. This is a strong preparation and has often proved successful in very bad cases of grease, which have resisted all the usual remedies.

Blisters. 1.—A general one.

Cantharides, powdered, 2 ounces,
Venice Turpentine, 2 ounces,
Resin, 2 ounces, Palm Oil or Lard. 2 lbs.

Melt the three latter articles, together, and when not too hot stir in the Spanish Flies.

2.—A Strong, Cheap Blister, but not proper to be used in Fevers or Inflammations, as of the Lungs, Bowels, etc.

Euphorbium, powdered, 1 ounce,
Oil of Vitriol, 2 scruples,
Spanish Flies, 6 ounces, Palm Oil or Lard,
Resin, of each 1 pound, Oil of Turpentine 3 ounces.

Melt the Resin with the Lard or Palm Oil. Having previously mixed the Oil of Vitriol with an ounce of the water gradually, as gradually add this mixture to the melted mass; which again set on a very slow fire for ten minutes more: afterwards remove the whole, and when beginning to cool, add the powders previously mixed together.

3.—A Mercurial Blister, for Splints, Spavins, and Ring-Bones.

Of either of the above 4 ounces,
Corrosive Sublimate finely powdered, ½ a drachm.

4.—Strong Liquid Blister.

Spanish Flies, in gross powder 1 ounce,
Oil of Origanum, 2 drachms,
Oil of Turpentine, 4 ounces. Olive Oil, 2 ounces.

Steep the Flies in the Turpentine three weeks, strain off and add the oil.

5.—Mild Liquid or Sweating Blister.

Of the above, 1 ounce,
Olive Oil or Goose-grease, 1½ ounces.

Clysters. 1.—A Laxative one.

Thin Gruel or Broth, 5 quarts,
Epsom or Common Salts, 6 ounces.

2.—Clyster for Gripes.

Mash 2 moderate sized Onions,
Pour over them Oil of Turpentine, 2 ounces,
Capsicum or Pepper, ½ an ounce,
Thin Gruel, 4 quarts.

3.—Nutritious Clyster.

Thick Gruel, 3 quarts, Strong sound Ale, 1 quart.

Or 4.—Stong Broth, 2 quarts,
Thickened Milk, 2 quarts.

5.—Astringent Clyster.

Tripe Liquor or Suet boiled in Milk, 3 pints,
Thick Starch, 2 pints, Laudanum, ½ an ounce.

Or 6.—Alum Whey, 1 quart, Boiled Starch, 2 quarts.

Cordial Balls.

Gentian powdered 4 ounces,
Ginger powdered, 2 ounces,
Coriander Seeds powdered, 4 ounces,
Caraway Seeds, 4 ounces,
Oil of Aniseed, ¼ of an ounce.

Make into a mass with Honey, Treacle, or Lard, and give an ounce and a half for a dose.

Chronic Cough Balls.

Calomel 1 scruple, Gum Ammoniacum,
Horse-radish, of each 2 drachms, Balsam of Tolu,
Squills, of each 1 drachm.

Beat all together, and make into a ball with Honey, and give every morning fasting.

2.—Drink for the same.

Tar Water, Lime Water, or each ½ a pint,
Tincture of Squills, ½ an ounce.

3.—Powder for the same.

Tartar Emetic, 2 drachms,
Powdered Foxglove, ½ a drachm,
Powdered Squill, ½ a drachm,
Calomel, 1 scruple, Nitre, 3 drachms.

Give every night in a malt mash.

Diuretic Balls.

Resin, yellow, 1 pound, Nitre, ½ a pound.
Horse Turpentine, ½ a pound,
Yellow Soap, ¼ of a pound.

Melt the Resin, Soap, and Turpentine over a slow fire; when cooling, add the Nitre. For a strong dose, an ounce and a half, for a mild one, an ounce. It should be kept in mind, *that mild diuretics are always equal to what is required:* and that strong diureties are always hurtful.

Diuretic Powders.

Yellow Resin, Powdered, 4 ounces,
Nitre, powdered, 8 ounces,
Cream of Tartar, powdered, 4 ounces.

Dose—6, 8, or 10 drs. nightly, which some horses will readily eat in a mash.

Urine Drink.

Glauber's Salts, 2 ounces. Nitre, 6 drachms.

Dissolve in a pint of warm water.

1.—Embrocations—Cooling for Inflammations.

Goulard's Extract, ½ an ounce.
Spirits of Wine or Brandy, 1 ounce,
Soft Water, 1 quart.

2.—Mindererus Spirit, 4 ounces, Water, 12 ounces.

For Sprains.

Bay Salt, bruised, ½ a pouud,
Crude Sal Ammoniac, 2 ounces,
Sugar of Lead, ¼ of an ounce,
Vinegar, 1½ pint, Water, 1 pint.

1.—For the Eyes.

Sugar of Lead, 1 drachm, White Vitriol, 2 scruples,
Water, 1 pint.

2—Brandy, 1 ounce, Infusion of Green Tea, 4 ounces,
Tincture of Opium, 2 drachms,
Infusion of Red Roses, 4 ounces.

3—Rose Water, 6 ounces,
Mindererus Spirit, 3 ounces.

4—Corrosive Sublimate, 4 grains, Alcohol, 1 ounce,
Lime Water, 1 pint.

5—Alum, powdered, 1 drachm,
Calomel, ½ a drachm.

Mix and insert a little at one corner of the eye. The custom of blowing it in alarms the horse.

1.—Fever Powders.

Tartar Emetic, 2 drachms, Nitre, 5 drachms.

2—Antimonial powder, 2 drachms,
Cream of Tartar, Nitre, of each 4 drachms.

3.—Fever Drink.

Sweet Spirit of Nitre, 1 ounce.
Mindererus Spirit, 6 ounces, Water, 4 ounces.

4.—Epidemic Fever Drink.

Sweet Spirit of Nitre, 1 ounce,
Simple Oxymel, 6 ounces,
Tartar Emetic, 3 drachms.

5.—Malignant Epidemic Fever.

Simple Oxymel, Mindererus Spirits,
Beer Yeast, of each 4 ounces,
Sweet Spirit of Nitre, 1 ounce.

Fumigations for Purifying Infected Stables, Sheds, etc.

Manganese, 2 ounces, Common Salt, 2 ounces,
Oil of Vitriol, 3 ounces, Water, 1 ounce.

Put the mixed Manganese and Salt into a basin; then, having before mixed the Vitriol and Water very gradually, pour them by means of tongs, or any thing that will enable you to stand at a sufficient distance, on the articles in the basin gradually. As soon as the fumes rise, retire and shut up the door close. Carbolic Acid is now used as a disinfectant.

Hoof Liquid.

Oil of Turpentine, 4 ounces,
Tar, 4 ounces, Whale Oil, 8 ounces.

This softens and toughens the hoofs extremely, when brushed over them night and morning.

Purging Medicines.

Balls—very mild.
Aloes, powdered, 6 drachms,
Oil of Turpentine, 1 drachm.

Mild.—Aloes, powdered, 8 drachms,
Oil of Turpentine, 1 drachm,

Strong.—Aloes, powdered, 10 drachms.
Oil of Turpentine, 1 drachm.

The Aloes may be beaten with Treacle to a mass adding, during the beating the Oil of Turpentine. All Spices, Cream of Tartar, Jalap, etc., are useless, and often hurtful additions.

Liquid Purge.

Epsom Salts, dissolved, 8 ounces,
Castor Oil, 4 ounces,
Watery Tincture of Aloes, 8 ounces.

Mix.—The Watery Tincture of Aloes is made by beating powdered Aloes with the yolk of egg, adding water by degrees, by these means half an ounce of Aloes may be suspended in 8 ounces of water, and such a purge is useful when a ball cannot be got down, as in partial locked jaw.

Scalding Mixture for Poll-Evil.

Corrosive Sublimate, finely powdered, 1 drachm,
Yellow Basilicon, 4 ounces.

Foot Stoppings.

Horse and Cow dung, each about 2 lbs.
Tar, ½ a pound.

Wash for Coring out, Destroying Fungus, or Proud Flesh, etc.

Lunar Caustic, 1 drachm, Water, 2 ounces.

Wash for Mange.

Corrosive Sublimate, 2 drachms,
Spirits of Wine or Brandy, 1 ounce,
Decoction of Tobacco,
Decoction of White Helebore, of each 1 pint.

Dissolve the Mercury in the spirits, and then add the decoctions.

1.—Ointment for Healing.

White Vitriol, powdered, ½ a drachm,
Turner's Cerate, 2 ounces, Lard, 4 ounces.

2.—For Digesting.

Turner's Cerate, 2 ounces, White Vitriol, 1 dra
Yellow Basilicon, 5 ounces.

For Mange.

Sulphur Vivum, 8 ounces,
Arsenic in powder, 2 drachms,
Mercurial Ointment, 2 ounces,
Turpentine, 2 ounces, Lard, 8 ounces.

Mix, and dress with every morning.

For Scab or Shab in Sheep, Mallenders and Sellenders in Horses, and Foul Blotches and Eruptions in Cattle in general.

Camphor, 1 drachm, Sugar of Lead, ½ a drachm,
Mercurial Ointment, 1 ounce.

CAUTION :—*Always keep your Bottles and Powders labelled with the proper names on them, this may save life.*